


SOUTHERN BLUEFIN TUNA (STN)

(Thunnus maccoyii)



1. FISHERY SUMMARY

Southern bluefin tuna were introduced into the QMS on 1 October 2004 under a single QMA, STN 1, with allowances, TACC, and TAC in Table 1.

Table 1: Recreational and Maori allowances, TACCS and TAC for southern bluefin tuna.

Fishstock	Recreational Allowance (t)	Maori customary Allowance (t)	Other mortality (t)	TACC (t)	TAC (t)
STN 1	4	1	2	413	420

Southern bluefin tuna were added to the Third Schedule of the 1996 Fisheries Act with a TAC set under s14 because a national allocation of southern bluefin tuna for New Zealand has been determined as part of an international agreement. The TAC applies to all New Zealand fisheries waters, and all waters beyond the outer boundary of the exclusive economic zone.

Southern bluefin tuna were also added to the Sixth Schedule of the 1996 Fisheries Act with the provision that:

“A person who is a New Zealand national fishing against New Zealand’s national allocation of southern bluefin tuna may return any southern bluefin tuna to the waters from which it was taken from if –

- (a) that southern bluefin tuna is likely to survive on return; and
- (b) the return takes place as soon as practicable after the southern bluefin tuna is taken”.

Management of southern bluefin tuna throughout its range is the responsibility of the Commission for Conservation of Southern Bluefin Tuna (CCSBT) of which New Zealand is a founding member. Current members of the CCSBT also include Australia, Japan, the Republic of Korea, and the Fishing Entity of Taiwan. The Republic of South Africa, the European Community, and the Philippines have Cooperating Non-member status. Indonesia have expressed interest in becoming members of the Commission. Determination of the global TAC and provision of a national allocation to New Zealand is carried out by the CCSBT. New Zealand was allocated a catch limit of 420 t (whole weight) at the 11th meeting of CCSBT in October 2004. This allocation has remained at this level since 1989.

In July 2006, the CCSBT Commission reviewed the results of two joint Australia / Japan reviews: the first was an assessment of the amount of southern bluefin tuna being sold through Japanese markets (referred to as the Market Review), and the second was an assessment of the potential for overcatch from the Australian surface fishery through the subsequent farming operations (referred to as the Farming Review).

The Market Review reported that quantities of southern bluefin tuna sold through the Japanese markets (back to the mid-1980s) were well in excess of the amount reported by Japan as domestic catch or imported from other countries (measured through the Trade Documentation Scheme), i.e. there were large volumes of unreported catch. The Market Review could not determine where the catch came from.

The Farming Review reported that while the catch in numbers from the surface fishery were probably well reported there was scope for biases in reported catch in weight due to two factors: (1) changes in the weight of fish between the time of capture and when the weight sample is taken; and (2) the sample of fish taken to estimate the mean weight of fish in the catch may not be representative (causing either negative or positive biases in the mean weight estimate).

The Farming Review was inconclusive. To remove doubt Australia has agreed to undertake a research program to address some of the issues raised in the Farming Review.

While Japan does not accept the findings of the Market review they have acknowledged some illegal catch during the 2005 fishing season and recently changed how they manage their fishery and accepted a cut in their allocated catch to 3000 t down from 6065 t. Current allocations are provided in Table 2 below.

Table 2: Allocated catches for Members, Cooperating Non-members and Observers.

	Allocated catch (t)
Australia	5265
Fishing Entity of Taiwan [#]	1140
Japan	3000
New Zealand	420
Republic of Korea [#]	1140
European Community	10
Indonesia	750
Philippines	45
South Africa	40

[#] The Fishing Entity of Taiwan and the Republic of Korea have both agreed to voluntarily limit their catches to 1000t.

The lack of acceptance of the findings of the two reviews has resulted in considerable uncertainty in the southern bluefin tuna science process as even the most fundamental data, i.e. a catch history, are not known and may be very different from previously thought. Further many of the indicators of stock status previously relied upon are now under question as they may be biased due to illegal activity.

This working group report has not been updated to reflect the findings of these two reviews, but in some places the possible impact of the reviews are noted.

1.1 Commercial fisheries

The Japanese distant water longline fleet began fishing for southern bluefin tuna in the New Zealand region in the late 1950s and continued after the declaration of New Zealand's EEZ in 1979 under a series of bilateral access agreements until 1995 (Table 4).

The domestic southern bluefin tuna fishery began with exploratory fishing by Watties in 1966 and Ferons Seafoods in 1969. Most of the catch was used for crayfish bait (reported landings began in 1972). During the 1980s the fishery developed further when substantial quantities of southern bluefin tuna were air freighted to Japan. Throughout the 1980s, small vessels handlining and trolling for

SOUTHERN BLUEFIN TUNA (STN)

southern bluefin tuna dominated the domestic fishery. Southern bluefin tuna were landed to a dedicated freezer vessel serving as a mother ship, or, ashore for the fresh chilled market in Japan. Longlining for southern bluefin tuna was introduced to the domestic fishery in the late 1980s under government encouragement and began in 1988 with the establishment of the New Zealand Japan Tuna Company Ltd. New Zealand owned and operated longliners, mostly smaller than 50 GRT, began fishing in 1991 for southern bluefin tuna (1 vessel). The number of domestic vessels targeting STN expanded throughout the 1990s and early 2000s prior to the introduction of STN into the QMS. Table 3 summarises southern bluefin landings in New Zealand waters since 1972.

Since 1991 surface longlines have been the predominant gear used to target southern bluefin tuna in the domestic fishery with 96% of all days fished using this method and only 4% using hand line (< 1% used trolling). This represents a major change from the 1980s when most fishing was by hand line.

Table 3: Reported domestic¹ and total² southern bluefin tuna landings (t) from 1972 to 2007 (calendar year).

Year	NZ Landings (t)	Total stock (t)	Year	NZ Landings (t)	Total stock (t)
1972	1	51 925	1990	529	13 870
1973	6	41 205	1991	164	13 691
1974	4	46 777	1992	279	14 217
1975	0	32 982	1993	217	14 344
1976	0	42 509	1994	277	13 154
1977	5	42 178	1995	436	13 637
1978	10	35 908	1996	139	16 356
1979	5	38 673	1997	334	16 076
1980	130	45 054	1998	337	17 776
1981	173	45 104	1999	461	19 529
1982	305	42 788	2000	380	15 475
1983	132	42 881	2001	358	16 027
1984	93	37 090	2002	450	15 240
1985	94	33 325	2003	389	14 060
1986	82	28 319	2004	393	13 623
1987	59	25 575	2005	264	16 146
1988	94	23 145	2006	238	11 850
1989	437	17 843	2007	378	11 540

¹ Domestic here includes catches from domestic vessels and Japanese vessels operating under charter agreement, i.e. all catch against the New Zealand allocation;

² These figures are likely underestimates as they do not incorporate the findings from the Market and Farming Reviews

Source: NZ data from Annual Reports on Fisheries, MAF data, NZ Fishing Industry Board Export data and LFRR data; Total stock from www.ccsbt.org.

In the few instances when the New Zealand allocation has been exceeded, the domestic catch limit has been reduced in the following year by an equivalent amount. Table 3 contrasts New Zealand STN catches with those from the entire stock. The low catches relative to other participants in the global fishery are due to New Zealand fisher's limited involvement historically rather than to local availability. Table 4 indicates that throughout most of the 1980s catches of STN up to two thousand tonnes were taken within the New Zealand EEZ.

Data on reported catch of southern bluefin tuna are available from the early 1950s. By 1960 catches had peaked at nearly 80 000 t, most taken on longline by Japan. From the 1960s through the mid 1970s, when Australia was expanding their domestic surface fisheries for southern bluefin tuna, total catches were in the range 40 000 to 60 000 t. From the mid 1970s through the mid 1980s catches were in the range 35 000 to 45 000 t. Catches declined from 33 325 t in 1985 to 13 869 t in 1990 and have fluctuated (average of 15 242 t per year) since that time (see Table 3). However, it should be noted that reported total catches are likely to be underestimates, at least after 1989, as they do not incorporate the findings from the Market and Farming Reviews. Despite this uncertainty the catches reported in 2007 (11 540 t) are the lowest estimated global catch for over 50 years.

Table 4: Reported catches or landings (t) of southern bluefin tuna by fleet and Fishing Year. NZ: New Zealand domestic and charter fleet, ET: catches by New Zealand flagged vessels outside these areas, JPNFL: Japanese foreign licensed vessels, KORFL: foreign licensed vessels from the Republic of Korea, LFRR: Estimated landings from Licensed Fish Receiver Returns, and MHR: Monthly Harvest Return Data.

Fish Yr	JPNFL	NZ	Total	LFRR/MHR	NZ ET
1979/80	7 374.7		7 374.7		
1980/81	5 910.8		5 910.8		
1981/82	3 146.6		3 146.6		
1982/83	1 854.7		1 854.7		
1983/84	1 734.7		1 734.7		
1984/85	1 974.9		1 974.9		
1985/86	1 535.7		1 535.7		
1986/87	1 863.1		1 863.1	59.9	
1987/88	1 059.0		1 059.0	94.0	
1988/89	751.1	284.3	1 035.5	437.0	
1989/90	812.4	379.1	1 191.5	529.3	
1990/91	780.5	93.4	873.9	164.6	
1991/92	549.1	248.9	798.1	279.1	
1992/93	232.9	126.6	359.5	216.4	
1993/94	0.0	287.3	287.3	277.0	
1994/95	37.3	358.0	395.2	435.3	
1995/96		141.8	141.8	140.5	
1996/97		331.8	331.8	333.5	
1997/98		330.8	330.8	331.5	
1998/99		438.1	438.1	457.9	
1999/00		378.3	378.3	381.3	
2000/01		366.0	366.0	366.4	
2001/02		468.3	468.3	452.3	
2002/03		405.7	405.7	391.7	0.0
2003/04		399.6	399.6	394.1	0.0
2004/05		272.1	272.1	263.3	0.0
2005/06		237.7	237.7	237.3	0.1
2006/07*		379	379	379	-

* - Southern bluefin tuna landings are not separated into within zone and ET for 2006/07 data

From 1960 to the 1990s catches by longline declined while surface fishery catches in Australian waters increased to reach its maximum level of 21 512 t in 1982 (equal to the longline catches of Japan). During the 1980s catches by both surface and longline fisheries declined but following dramatic TAC reductions in the late 1980s, catches stabilised. The main difference between gear types is that surface fisheries target juveniles (age-1 to age-3 year olds) while longline fisheries catch older juveniles and adults (age-4 year old up to age-40+). The surface fishery has comprised purse seine and pole-&-line vessels supported by aerial spotter planes that search out surface schools. The Australian surface fisheries that prior to 1990 were a mix of pole-&-line and purse seine vessels, have since the mid-1990s become almost exclusively a purse seine fishery. Whereas prior to 1990 surface fishery catches supplied canneries, since the mid-1990s these vessels catch juveniles for southern bluefin tuna farms where they are “on-grown” for the Japanese fresh fish market. In contrast to Australia, the fisheries of all other members, (including New Zealand) is based on longline. Historically New Zealand also supported handline and troll fisheries for STN, although these were small scale and targeted large adults.

Analysis of New Zealand catch data shows that most southern bluefin tuna are caught in FMA1, FMA2, FMA5 and FMA7. The northern FMAs (FMA1 and FMA2) that accounted for a small proportion of southern bluefin tuna before 1998 have in recent years accounted for about the same amount of southern bluefin tuna as the southern FMAs (FMA5 and FMA7).

This change in spatial distribution of catches can be attributed to the increase in domestic longline effort in the northern waters.

1.2 Recreational fisheries

There is no estimate available for recreational catch of southern bluefin tuna, although charter vessels based in Milford Sound are known to have targeted this species historically and take it as bycatch in the newly developed Pacific bluefin tuna fishery.

SOUTHERN BLUEFIN TUNA (STN)

1.3 Maori customary fisheries

Given that Maori knew of several oceanic fish species and missionaries reported that Maori regularly fished several miles from shore, it is possible that southern bluefin tuna were part of the catch of Maori prior to European settlement. It is clear that Maori trolled lures (for kahawai) that are very similar to those still used by Tahitian fishermen for small tunas and also used large baited hooks capable of catching large southern bluefin tuna. However, there is no Maori name for southern bluefin tuna. It is therefore, uncertain if Maori caught southern bluefin tuna. An estimate of the current customary catch is not available.

1.4 Illegal catch

There is no known illegal catch of southern bluefin tuna by New Zealand vessels in the EEZ or from the high seas. The recent review of the Japanese Market suggests very large illegal catch from the broader stock historically.

Recent actions by individual CCSBT members to improve monitoring, control, and surveillance measures for southern bluefin tuna fisheries is intended to halt the occurrence of unreported catch.

1.5 Other sources of mortality

Incidental catches of southern bluefin tuna appear to be limited to occasional small catches in trawl fisheries. Small catches of southern bluefin tuna have been reported as non-target catch (< 0.5 t and 2 t respectively), in trawl fisheries for hoki (*Macruronus novaezelandiae*) and arrow squid (*Notodarus* spp.). In addition there have been occasional anecdotal reports of southern bluefin being caught in trawl fisheries for southern blue whiting (*Micromesistius australis*) and jack mackerel (*Trachurus* spp.) in sub-Antarctic waters.

In addition to the limited trawl bycatch there is some discarding (usually as a result of shark damage) and loss before fish are landed that occurs in the longline fishery. The estimated overall incidental mortality rate from observed longline effort is 0.54% of the catch. Discard rates are 0.86% on average from observer data of which approximately 50% are discarded dead. Fish are also lost at the surface in the longline fishery during hauling, 1.47% on average from observer data, of which 95% are thought to escape alive. An allowance of 2 t has been made for other sources of mortality.

2. BIOLOGY

The age at which 50% of southern bluefin are mature is uncertain because of limited sampling of fish on the spawning ground off Java. Recent sampling of the Indonesian catch suggests that 50% age-at-maturity may be as high as 12 years, while interpretations of available data since 1994 have used 8 years and older fish as representing the adult portion of the stock in the population models.

As the growth rate has changed over the course of the fishery (see following section & Table 5) the size at maturity depends on when the fish was alive (prior to the 1970s, during the 1970s, or in the period since 1980), as well as which maturity ogive is used. A simple linear interpolation is assumed for the 1970s. Table 5 shows the range of sizes (cm) for southern bluefin tuna aged 8 to 12 years for the two von Bertalanffy growth models used.

Table 5: Differences in southern bluefin tuna size at ages 8 – 12 between the 1960s and 1980s (lengths in cm).

Age	1960s	1980s
8	138.2	147.0
9	144.6	152.7
10	150.2	157.6
11	155.1	161.6
12	159.4	165.0

Radiocarbon dating of otoliths has been used to determine that southern bluefin tuna live beyond 30 years of age and that individuals reaching asymptotic length may be 20 years or older.

The sex ratio of southern bluefin caught by longline in the EEZ has been monitored since 1987. The ratio of males to females is 1.2:1.0, and is statistically significantly different than 1:1.

The parameters of length:weight relationships for southern bluefin tuna based on linear regressions of greenweight versus fork length are in Table 6.

Table 6: Parameters of length/ weight relationship for southern bluefin tuna. $\ln(\text{Weight}) = b_1 \ln(\text{length}) - b_0$ (Weight in kg, length in cm).

	b_0	B_1
Male	-10.94	3.02
Female	-10.91	3.01
All	-10.93	3.02

The data used include all longline observer data for the period 1987 to 2000 from all vessels in the EEZ (n = 18 994).

CCSBT scientists have used two stanza Von Bertalanffy growth models since 1994:

$$l_t = L_\infty(1 - e^{-k_2(t-t_0)})(1 + e^{-\beta(t-t_0-\alpha)}) / (1 + e^{\beta\alpha})^{-(k_2-k_1)}, \text{ where } t \text{ is age in years.}$$

Table 7: von Bertalanffy growth parameters for southern bluefin tuna.

	L_∞	k_1	k_2	α	β	t_0
1960 von Bertalanffy	187.6	0.47	0.14	0.75	30	0.243
1980 von Bertalanffy	182	0.23	0.18	2.9	30	-0.35

While change in growth in the two periods (pre-1970 and post 1980) is significant and the impact of the change in growth on the results of population models substantial, the differences between the growth curves seem slight. The change in growth rate for juveniles and young adults has been attributed to a density dependant effect of over fishing.

No estimates of F and Z are presented because they are model dependent and because a range of models and modelling approaches are used with no agreement as to which model(s) is/are best. Prior to 1995 natural mortality rates were assumed to be constant and $M = 0.2$ was used. However, the results indicating that asymptotic size was reached at about 20 years and fish older than 30 years were still in the population, suggested that values of $M \geq 0.2$ were likely to be too high. Tagging results of juvenile's ages 1 to 3 years also suggests that M for these fish is high (possibly as high as $M = 0.4$), while M for fish of intermediate years is unknown. For these reasons M has been considered to be age-specific and represented by various M vectors. In the CCSBT stock assessments, a range of natural mortality vectors are now used.

A conversion factor of 1.15 is used for gilled and gutted southern bluefin tuna.

3. STOCKS AND AREAS

Southern bluefin tuna consist of a single stock primarily distributed between 30°S and 45°S, which is only known to spawn in the Indian Ocean south of Java.

Adults are broadly distributed in the South Atlantic, Indian and western South Pacific Oceans, especially in temperate latitudes while juveniles occur along the continental shelf of Western and South Australia and in high seas areas of the Indian Ocean. Southern bluefin tuna caught in the New Zealand EEZ appear to represent the easternmost extent of a stock whose centre is in the Indian Ocean.

SOUTHERN BLUEFIN TUNA (STN)

A large-scale electronic tagging programme, involving most members of the Commission, has been undertaken to provide better information on stock structure. The goal has been to tag smaller fish across the range of the stock. New Zealand has participated in this programme, having deployed 19 implantable tags in small fish in 2007. Fifteen larger SBT were tagged with pop-off tags as well, with 12 tags having reported data thus far. Of note, one of the tagged fish moved to the spawning ground south of Indonesia.

Electronic tagging of juvenile STN in the Great Australian Bight has shown that in recent years tagged juveniles have not been moving into the Tasman Sea. It is not known whether this is due to unfavourable environmental conditions or range contraction following the decline in the stock.

Two sources of information suggest that there may be 'sub-structure' within the broader STN stock, in particular the Tasman Sea. Tagging of adult STN within the Australian east coast tuna and billfish fishery suggests that STN may spend most of the years within the broader Tasman Sea region. An analysis of the length and age composition of catches from the New Zealand JV fleet showed that cohorts that were initially strong or weak did not change over time, e.g. if a particular year class was weak (or strong) when it initially recruited to the New Zealand fishery it did not become strong (or weak) over time.

4. STOCK ASSESSMENT

Determination of the status of the southern bluefin tuna stock is undertaken by the CCSBT Stock Assessment Group (CCSBT-SAG) and Scientific Committee (CCSBT-SC) and is based on an analysis of fishery indicators and the results from an operating model. Indicators considered included those relating to recent recruitment, spawning biomass, and vulnerable biomass and were based on catch at age data, CPUE data, and information from various surveys (e.g. acoustic and aerial sightings). There is no agreed stock assessment model, though results from an operating model of the STN population and fishery was considered in 2004 and updated annually since 2006.

In 2006 serious doubts were raised on the reliability of the total catch and the CPUE indicators from the Japanese longline fishery. A study of the volume of fish traded through the market showed a large under-reporting of catch had occurred over a number of years (market anomaly). A study of the Australian farming operations also suggested the total catch in this fishery had been under-estimated (farm anomaly). As these problems made routine stock assessment impossible a scenario modelling exercise was used to provide management advice. On this basis, the Commission set constant TAC levels for each country for 3 years (and for 5 years for Japan) that would only be reviewed under exceptional circumstances.

This section draws on the conclusions of the CCSBT-SAG and CCSBT-SC.

4.1 Estimates of fishery parameters and abundance

As part of the stock assessment, a range of fishery indicators that were independent of any stock assessment model were considered to provide support and/or additional information important to aspects of current stock status. The reviews of Japanese SBT market anomalies and Australian SBT farming anomalies raise serious doubts on the reliability of the catch and Japanese LL CPUE indicators, thus interpretation of many of the indicators is more difficult than in previous years.

(i) Recruitment

The indicators continue to support the previous evidence for poor recruitment in the 2000 and 2001 year class, and ongoing recruitment below the 1994-1998 levels. The size distribution in the NZ LL fishery (Figure 1) and the Japanese longline (LL) fishery continue to indicate poor 2000 and 2001 recruitments, and the aerial spotting survey and commercial spotting index are both consistent with a reduction in average recruitment below the 1994-1998 levels. The high fishing mortality rate estimates for age 3 and 4 from recent SRP tagging are also consistent with low recruitments in these years. Trends in year class strength in the Japanese LL fleet show poor strength of the 2000 and 2001

year classes, but recent data indicates an increase in juveniles after the 2002 year class. However, this indicator could be biased by catch anomalies.




Figure 1: Proportion at length for the charter fleet for 2001 to 2008. Data for 2008 is preliminary.
 Source: NZ Report to CCSBT (2008).

(ii) Spawning biomass

Reported catch rates of fish aged 12 and older in the Japanese LL continue to indicate a drop in spawning stock biomass in about 1995, but this is potentially impacted by catch anomalies. Since the

SOUTHERN BLUEFIN TUNA (STN)

Japanese LL CPUE is the primary indicator of stock abundance the potential anomalies make the spawning stock status less certain than last year. The increase in tonnage of Indonesian catch as well as the increase in proportion of SBT in the Indonesian catch was associated with a shift in the behaviour of the Indonesian fleet to target SBT south of the spawning ground. This change in behaviour complicates the interpretation of the age and size structure of catches from the spawning stock.

(iii) Vulnerable biomass

Reported Japanese LL CPUE of SBT for all ages combined suggests that the exploitable biomass for these gears has remained fairly constant during the past 10 years, though this level is low compared to historical values. Confidence in this indicator has diminished considerably due to the uncertainty associated with catch anomalies. Reported CPUE indicate increases in the CPUE of ages 8-11 since about 1992, but there is a slight decline in 2003 and 2004, with a slight increase in 2005. Reported CPUE of fish aged 4-7 has increased since the mid 1980s but has been declining in recent years.

For the New Zealand Charter fleet CPUE in the core part of the fishery (west coast South Island) averaged around 3 SBT per 1000 hooks over 1997-2002 (Figure 2). Associated with the lack of new recruitment, CPUE declined dramatically in 2003 and has stayed at these historically low levels in 2004 and 2005.




Figure 2: Catch per unit effort (number of SBT per thousand hooks) from the Charter fleet in Region 6 (west coast South Island) for all southern bluefin tuna (dashed line) and for fish greater than 10 years of age based on proportional ageing data (solid line, open symbols) and based on direct ageing data (solid line, solid symbols). Source: NZ report to CCSBT (2008).

4.2 Biomass estimates

Biomass estimates depend on the specific population model used and important factors relating to potential catch anomalies; hence there is no agreed set of biomass estimates for southern bluefin tuna.

4.3 Estimation of Maximum Constant Yield (MCY)

MCY has not been estimated.

4.4 Estimation of Current Annual Yield (CAY)

CAY has not been estimated.

4.5 Other yield estimates and stock assessment results

At the 2008 CCSBT-SAG meeting the operating model was run under a number of scenarios that are generally similar to those evaluated in 2006. The scenarios indicate that spawning stock biomass is still at a very low level (generally below 10% of pre-exploitation spawning stock biomass, a level at which recruitment may be at risk of further decline). This is well below the 1980 level and below the level that could produce maximum sustainable yield. Rebuilding the spawning stock biomass would almost certainly increase sustainable yield and provide security against unforeseen environmental events. Presently, however, there is no sign of spawning stock biomass rebuilding.

Recruitments in the last two decades are estimated to be well below the levels in the period 1950-1980. All scenarios suggest that recruitment in the 1990s fluctuated at a low level with no overall trend. Analysis of the average of all indicators suggest historically low recruitments from 1999-2002. The indicators suggest that 2004 and 2005 year classes are stronger and close to the average of the 1990s.

Consistent with the poor recruitment from 1999 to 2002, a gap in the size (and presumably age) composition is apparent. By inference this gap will lead to a further decline in spawning stock biomass in coming years.

There remain serious sources of concern from new and previous information including:

- a very low spawning stock;
- at least three poor recruitments in the recent past which will lead to a further decline in spawning stock biomass;
- a general decline in recruitment since about 1970, coincident with declining spawning stock sizes;
- increased exploitation rates, particularly on recent weak year classes;
- overall, exploitation has shifted towards younger ages, the abundances of which are poorly estimated;
- there is the potential for increasing exploitation rate with declining recruitment, which would pose a serious risk to rebuilding.

The CCSBT-SC noted that given the current reported catch and previously reported constant catch projections, the stock is likely to rebuild very slowly in the long term. However, there is about a 50:50 chance that the spawning stock will decrease over the next 6 years. The possibility of further decline is of concern.

4.6 Other factors

It is generally agreed that there is strong evidence for a stock recruit relationship in southern bluefin tuna with high recruitment and high parental biomass in early years, low recruitment and low parental biomass in later years. Despite continuing low parental stock and low recruitment, the parental stock appears to have been relatively stable over much of the 1990s. However, the most recent assessment indicates that there is now very little chance that current catches will allow the stock to rebuild to the 1980 level by 2020 and evidence of recent poor recruitment, even if they continue for a few years, may mean that current catches will not be sustainable.

5. STATUS OF THE STOCK

The IUCN listed southern bluefin tuna as critically endangered in 1996 based on the rate of population decline. The three original Parties to the CCSBT (Australia, Japan and New Zealand) have met to assess stock status regularly and in nearly every case have agreed that the parental stock was at historically low levels and that the parental stock should be rebuilt to the level seen in the 1980s. This level of parental biomass coincided with a period when recruitment (on average) had been stable.

The CCSBT- Scientific Committee in 2008 provided the following summary of the status of the stock:

“At the 2008 CCSBT-SAG meeting the operating model was run under a number of scenarios that are generally similar to those evaluated in 2006. The scenarios indicate that spawning

SOUTHERN BLUEFIN TUNA (STN)

stock biomass is still at a very low level (generally below 10% of pre-exploitation spawning stock biomass, a level at which recruitment may be at risk of further decline). This is well below the 1980 level and below the level that could produce maximum sustainable yield. Rebuilding the spawning stock biomass would almost certainly increase sustainable yield and provide security against unforeseen environmental events. Presently, however, there is no sign of spawning stock biomass rebuilding.

Recruitments in the last two decades are estimated to be well below the levels in the period 1950-1980. All scenarios suggest that recruitment in the 1990s fluctuated at a low level with no overall trend. Analysis of the average of all indicators suggest historically low recruitments from 1999-2002. The indicators suggest that 2004 and 2005 year classes are stronger and close to the average of the 1990s.

Consistent with the poor recruitment from 1999 to 2002, a gap in the size (and presumably age) composition is apparent. By inference this gap will lead to a further decline in spawning stock biomass in coming years.”

The most recent assessment was undertaken in 2005 and covered the entire stock. On a regional level there are concerns relating to the current status of this stock and the level of fishing effort. The stock is presently well below the level necessary to produce the maximum sustainable yield. The global catch limit has been set at a level considered to reduce the possibility of further decline in the stock and hopefully lead to rebuild.

6. FOR FURTHER INFORMATION

- Anon. 2001. Report of the sixth meeting of the scientific committee, Tokyo, Japan, 28–31 August 2001.
- Anon. 2004. Report of the Fifth Meeting of the Stock Assessment Group, 6-11 September 2004, Jeju, Republic of Korea
- Anon. 2004. Report of the Ninth Meeting of the Scientific Committee, 13-16 September 2004, Jeju, Republic of Korea.
- Anon. 2005. Report of the Sixth Meeting of the Stock Assessment Group, 29 August - 3 September 2005, Taipei, Taiwan
- Anon. 2005. Report of the Tenth Meeting of the Scientific Committee, 9 September 2005, Narita, Japan.
- Anon. 2006. Report of the Seventh Meeting of the Stock Assessment Group, 4-11 September 2004, Tokyo, Japan
- Anon. 2006. Report of the Eleventh Meeting of the Scientific Committee, 12-15 September 2006, Tokyo, Japan.
- Best, E. 1929. Fishing methods and devices of the Maori. *Dominion Museum Bulletin* no 12.
- Caton, A. (ed.). 1991. Review of aspects of southern bluefin tuna biology, population and fisheries. I-ATTC Special Report 7: 181–350.
- Harley, S. J., and Kendrick, T. 2006. Annual Review of National SBT Fisheries for the Scientific Committee. CCSBT-ESC/0609/SBT Fisheries: New Zealand.
- Hufflett, C. 2000. Southern bluefin tuna. Unpublished report available from Solander Fisheries, Nelson.
- Kalish, J.M.; Johnston, J.M.; Gunn, J.S.; Clear, N.P. 1996. Use of bomb radiocarbon chronometer to determine age of southern bluefin tuna (*Thunnus maccoyii*). *Marine Ecology Progress Series* 143: 1–8.
- Leigh, G.M.; Hearn, W.S. 2000. Changes in growth of juvenile southern bluefin tuna (*Thunnus maccoyii*): an analysis of length–frequency data from the Australian fishery. *Marine and Freshwater Research* 51: 143–154.
- Murray, T.; Richardson, K.M. 2001. Trends in the New Zealand southern bluefin tuna fishery. CCSBT-SC/0108/SBT Fisheries – New Zealand.
- Murray, T.; Richardson, K.; Griggs, L. 2002. New Zealand tuna fisheries, 1991–2000. *N.Z. Draft Fisheries Assessment Report*.
- Strickland, R.R. 1990. Nga tini a Tangaroa, a Maori-English, English-Maori dictionary of fish names. NZ Fisheries Occasional Publication no. 5.