


## COCKLES (COC 3) Otago Peninsula

(*Austrovenus stutchburyi*)  
Tuaki



### 1. FISHERY SUMMARY

COC 3 was introduced into the Quota Management System in October 2002 with a TAC of 1500 t; comprising of a customary allowance of 10 t, a recreational allowance of 10 t, an allowance for other fishing related mortality of 10 t, and a TACC of 1470 t. Historical catch limits can be seen in Table 1.

#### 1.1 Commercial fisheries

Cockles are present at various locations around the Otago Peninsula but are only commercially fished from Papanui Inlet, Waitati Inlet, and Otago Harbour (under a current special permit). Commercial fishing in Papanui and Waitati Inlets began in 1983. A limit of 104 t was in effect for Papanui and Waitati Inlets combined from 1986–87 until 1991–92. From 1992–93 to 1998–99, the catch limits were 90 t for Papanui Inlet and 252 t for Waitati Inlet. In April 2000, the catch limits were increased to 427 t for Papanui Inlet and 746 t for Waitati Inlet. In 2002 when cockles entered the QMS spatial restrictions upon harvest within COC 3 were removed. Commercial landings from Papanui and Waitati Inlets are shown in Table 1. Since August 2009 cockles have been taken from Otago Harbour under a special permit in order to investigate the ecosystem effects of commercial cockle harvesting in this location. This permit states no explicit limit to the tonnage able to be taken but does delimit the area where harvest will be taken and presently expires on the 31<sup>st</sup> of August 2012<sup>1</sup>

In 1992, 35 mm shell length was the minimum size for commercial cockles. However, commercial fishers currently target  $\geq 28$  mm cockles, therefore 28 mm is used as the effective minimum size in yield calculations. CPUE data are available for this fishery, but have not been analysed.

#### 1.2 Recreational fisheries

Cockles are taken by recreational fishers in many areas of New Zealand. The recreational fishery is harvested entirely by hand digging. Relatively large cockles are preferred.

Amateur harvest levels in FMA 3 were estimated by telephone and diary surveys in 1993–94 (Teirney *et al.* 1997), 1996 (Bradford 1998) and 2000 (Boyd & Reilly 2002), Table 2. Harvest weights are estimated using an assumed mean weight of 25 g (for cockles  $>30$  mm). In 2004, the Marine Recreational Fisheries Technical Working Group reviewed the harvest estimates of these surveys and concluded that the 1993–94 and 1996 estimates were unreliable due to a methodological error. While the same error did not apply to the 1999–00 and 2000–01 surveys, it was considered the estimates

<sup>1</sup> This permit is able to be extended or revoked before this date.


## COCKLES (COC 3)

may still be very inaccurate. No recreational harvest estimates specific to the COC 3 commercial fishery areas are available.

**Table 1: Reported landings (t) of cockles from Papanui and Waitati Inlets, Otago, combined (FMA 3), from 1986–87 to 2011–12 based on Licensed Fish Receiver Returns (LFRR). Catch splits are provided by Southern Clams Ltd and are partially from Stewart (2005). N/A = Not Applicable [Continued on other page].**

Year	Papanui catch (t)	Papanui limit (t)	Waitati catch (t)	Waitati limit (t)	Otago Harbour catch (t)	Total catch (t)	Total limit (t)
1986–87	14	–	–	–	–	14	104
1987–88	8	–	–	–	–	8	104
1988–89	5	–	–	–	–	5	104
1989–90	25	–	–	–	–	25	104
1990–91	90	–	16	–	–	106	104
1991–92	90	–	14	–	–	104	104
1992–93	90	90	92	252	–	182	342
1993–94	90	90	109	252	–	199	342
1994–95	90	90	252	252	–	342	342
1995–96	90	90	252	252	–	342	342
1996–97	90	90	252	252	–	342	342
1997–98	90	90	252	252	–	342	342
1998–99	90	90	293	252	–	383	342
1999–00	118	427	434	746	–	552	1 273
2000–01	90	427	606	746	–	696	1 273
2001–02	49	N/A	591	N/A	–	640	1 273
2002–03	52	N/A	717	N/A	–	767	1 470
2003–04	73	N/A	689	N/A	–	762	1 470
2004–05	91	N/A	709	N/A	–	800	1 470
2005–06	68	N/A	870	N/A	–	943	1 470
2006–07	0*	N/A	907	N/A	–	907	1 470
2007–08	–	N/A	760	N/A	–	760	1 470
2008–09	–	N/A	751	N/A	24	775	1 470
2009–10	–	N/A	379	N/A	441	820	1 470
2010–11	–	N/A	240	N/A	596	836	1 470
2011–12	–	N/A	358	N/A	437	795	1 470

\*No catches have been taken from Papanui Inlet since 2006–07 because of water quality problems.



**Figure 1: Historical landings and TACC for COC 3 (Otago). QMA data from 2003-04 to present.**

### 1.3 Customary non-commercial fisheries

Many intertidal bivalves, including cockles, are very important to Maori as traditional food, particularly to Huirapa and Otakou Maori in the Otago area. Tangata tiaki issue customary harvest permits for cockles in Otago. The number of cockles harvested under customary permits is given in Table 3, and is likely to be an underestimate of customary harvest.

**Table 2: Estimated numbers of cockles harvested by recreational fishers in FMA 3, and the corresponding harvest tonnage. Figures were extracted from a telephone and diary survey in 1993–94, and the national recreational diary surveys in 1996 and 2000.**

Fishstock	Survey	Harvest (N)	% CV	Harvest (t)
FMA 3	1993–94 South	106 000	51	2.7
FMA 3	1996	144 000	–	3.6
FMA3	2000	1 476 000	45	36.9

On 1 October 2010, on the recommendation of the Taiāpure Committee, the Minister of Fisheries introduced new regulations for the East Otago Taiāpure<sup>2</sup>. These included a new amateur daily bag limit of 50 for shellfish, including cockles, and a ban on the commercial take of cockles from any part of the Taiāpure, except for the existing sanitation areas within Waitati Inlet. The new regulations reflect the Committee’s concern about fishing pressure on shellfish stocks, including cockles, within the Taiāpure.

A long-running time series of surveys suggest that there are no sustainability concerns in terms of cockles within the Taiāpure. However, they do indicate a shift in some beds towards smaller size classes of cockle. Larger cockles are preferred by both customary and recreational fishers. The Committee hopes that reducing the bag limit and limiting the spatial extent of commercial harvest will lead to an increase in the number of large cockles.

**Table 3: Number of cockles harvested under customary fishing permits.**

Year	Number of cockles
1998	750
1999	0
2000	1 109
2001	1 090
2002	0
2003	2 750
2004	4 390
2005	5 699

#### 1.4 Illegal catch

No quantitative information is available on the magnitude of illegal catch but it is thought to be insignificant.

#### 1.5 Other sources of mortality

No quantitative information is available on the magnitude of other sources of mortality. It has been suggested that some harvesting implements, such as brooms, rakes, “hand-sorters”, bedsprings and “quick-feeds” cause some incidental mortality, particularly of small cockles, but this proposition has not been scientifically investigated. High-grading of cockles is also practised, with smaller sized cockles being returned to the beds. The mortality from this activity is unknown, but is likely to be low.

## 2. STOCKS AND AREAS

Each inlet is assumed to be an independent fishery within the stock.

<sup>2</sup> The Kati Huirapa Runanga ki Puketeraki application for a taiāpure-local fishery was gazetted as the East Otago Taiāpure-Local Fishery in 1999. A management committee, made up of representatives from the Runanga and various recreational, environmental, commercial, community and scientific groups, was appointed in 2001.

### 3. STOCK ASSESSMENT

Stock assessments for Papanui Inlet and Waitati Inlet have been conducted using absolute biomass surveys, yield-per-recruit analyses, and Method 1 for estimating CAY (Annala *et al.* 2003). Breen *et al.* (1999) also estimated biomasses and yields for Otago Harbour and Purakanui. Stewart (2005, 2008a) estimated biomass and yields for Papanui and Waitati Inlets in 2004 and Waitati Inlet in 2007.

#### 3.1 Estimates of fishery parameters and abundance

A project to estimate growth and mortality in Papanui and Waitati Inlets, Purakanui and Otago Harbour was undertaken in the late 1990s. Notched clams did not exhibit significant growth when recovered after one year, and modes in the length frequency distributions did not shift when measured over four sampling periods within a year (Breen *et al.* 1999).

In 2004 and 2007 yield-per-recruit modelling was conducted for Papanui and Waitati inlets separately (Stewart 2005, 2008a). The most recent parameters used in this modelling are detailed in Table 2 of the cockle introductory section. Estimates of  $F_{0.1}$  from these studies are given in Table 4 below. Exploitation rate is below 7% for both Waitati and Papanui Inlet (Figure 2) and is unable to be calculated for Otago harbour.


**Table 4: Estimates of fishery parameters (recruitment to this fishery is at  $\geq 28\text{mm}$ )**

$M$	$F_{0.1}$ 2004	$F_{0.1}$ 2007
0.2	0.2321	0.2899
0.3	0.3412	0.3863
0.4	0.4767	0.5537

**Exploitation rate % (for cockles  $\leq 30$  mm across each entire inlet)\***

Year	Papanui	Waitati
1998	2	0
2002	1	5
2004	2	6
2007	0	7

\* This measure is likely to overestimate exploitation as harvest occurs down to a size limit of 28mm.



**Figure 2: Exploitation rate as calculated by landings divided by biomass ( $\geq 19$  mm) from whole inlets.**

### 3.2 Biomass estimates

Biomass surveys have been undertaken periodically in COC 3 since 1984. The methods for the calculation of biomass have changed over time<sup>3</sup> which means that comparison of biomass values between times of different calculation methodologies should be done cautiously.

The Spawning stock biomass ( $\geq 19$  mm shell length) has been stable around the level of virgin biomass in Waitati Inlet (Table 5, Figure 3). In Papanui Inlet the spawning stock biomass ( $\geq 19$  mm shell length) has shown a trend of gradual decline from 1984 until 2004, when it was at 78% of virgin biomass. The recruited biomass ( $\geq 30$  mm shell length) in the sanitation areas (beds 1804 and 1805) in Otago Harbour decreased prior to the start of harvesting in 2008.

**Table 5: Current ( $\pm 95\%$  CI) and previous biomass estimates from COC 3\*.**

Papanui Inlet								
Size Class	1984	1992	1998	2002	2004	2004	2004	
					Total inlet	Commercial area		
>2 to 18 mm (juveniles)	65	139	33	17 $\pm$ 1.7	36 $\pm$ 2.2		13 $\pm$ 1.3	
19 – 34 mm (adults)	3 705	3 721	3 435	1 970 $\pm$ 192	2 415 $\pm$ 151		825 $\pm$ 88	
$\geq 35$ mm	2 370	1 706	2 231	2 579 $\pm$ 252	2 301 $\pm$ 273		1 847 $\pm$ 208	
$\geq 30$ mm			3 990.2	3 860 $\pm$ 365	3 677 $\pm$ 367		2 420 $\pm$ 271	
Total (t)	6 140	5 567	5 699	4 565 $\pm$ 424	4 752 $\pm$ 425		2 685 $\pm$ 298	
Waitati Inlet**.								
Size Class	1984	1992	1998	2002	2004	2004	2007	2007
				Total Inlet	Commercial area	Total Inlet	Commercial area	
>2 to 18 mm (juveniles)	619	1 210	304	153 $\pm$ 20	257 $\pm$ 14	77 $\pm$ 4	335 $\pm$ 26	102 $\pm$ 7.5
19 to 34 mm (adults)	7 614	5 198	8 519	6 653 $\pm$ 652	7 272 $\pm$ 403	2 735 $\pm$ 129	7 673 $\pm$ 591	1 284 $\pm$ 95***
$\geq 35$ mm	3 844	4 620	4 381	4 298 $\pm$ 298	4 535 $\pm$ 508	3 872 $\pm$ 384	3 941 $\pm$ 462	
$\geq 30$ mm			7 235	7 183 $\pm$ 463	7 993 $\pm$ 720	5 612 $\pm$ 681	7 107 $\pm$ 548	4 726 $\pm$ 352
Total (t)	12 080	11 027	13 204	11 103 $\pm$ 848	12 064 $\pm$ 925	6 685 $\pm$ 517	11 948 $\pm$ 921	6 112 $\pm$ 456
			1998	2008				
Purkaunui Inlet ( $\geq 30$ mm)			1 825					
Otago Harbour ( $\geq 30$ mm)			32 975					
Otago Harbour (sanitation area 1804, $\geq 30$ mm)			8 901	5 473				
Otago Harbour (sanitation area 1805, $\geq 30$ mm)			5 546	3 526				

\*Wildish 1984; Stewart *et al.* 1992; Breen *et al.* 1999; Wing *et al.* 2002; Stewart, 2005; Stewart 2008a, Stewart 2008b. Area of current commercial beds, Papanui Inlet = 815,811 m<sup>2</sup>. \*\*Area of current commercial beds, Waitati Inlet = 943,986 m<sup>2</sup>. \*\*\* = this value is only for  $\geq 19$  mm to <30 mm cockles.

---

Wildish (1984a and b) and Stewart *et al.* (1992) separated cockles by sieving into three size classes. Breen *et al.* (1999) measured random samples of cockles from each inlet to calculate length-weight relationships. The first method only allows estimation of biomass from predetermined size classes. By calculating size structure of populations using length to weight data a more flexible approach is allowed where data can be matched to current commercial needs as well as to future survey results. The 1998 survey used random samples from each inlet to calculate length to weight relationships (Breen *et al.* 1999). This method was once again used in the 2002 survey (Wing *et al.* 2002). In the 2004 and 2007 surveys random samples from each shellfish bed were weighed and their longest axis measured (Stewart 2005, 2008). These data were then used to generate length to weight relationships

## COCKLES (COC 3)




Figure 3: Biomass as a proportion of  $B_0$  for Waitati and Papanui Inlets, this is estimated from biomass >19mm.

### 3.3 Yield estimates and projections

Estimates of  $MCY$  are given in Table 6.

Table 6: Estimates of  $MCY(t)$  for COC 3 generated using Method 1 (Annala *et al.* 2003) an average biomass  $\geq 30$  mm as  $B_0$  and the 2007 estimate of  $F_{0.1}$ . This calculation is likely to underestimate the true  $MCY$ .

Location	M	1998	2002	2004	2007
Waitati Inlet	0.2	1049	1045	1083	1070
Waitati Inlet	0.3	1397	1392	1443	1425
Waitati Inlet	0.4	2003	1996	2068	2043
Waitati Inlet (commercial)	0.2			813	749
Waitati Inlet (commercial)	0.3			1084	998
Waitati Inlet (commercial)	0.4			1554	1431
Papanui Inlet	0.2	289	280	266	
Papanui Inlet	0.3	385	373	355	
Papanui Inlet	0.4	552	534	509	
Papanui Inlet (commercial)	0.2			175	
Papanui Inlet (commercial)	0.3			234	
Papanui Inlet (commercial)	0.4			335	

For Waitati Inlet,  $CAY$  was estimated (Table 7) using Method 1 ( $CAY = (F_{0.1}/Z) (1 - \exp(-Z))B_{BEG}$ ) (Annala *et al.* 2003) and biomass estimates at different times.  $CAY$  has been estimated at times for both the entire inlet area and a subset area where the commercial fishery has been operating for the past several years. This approach assumes that, between the start of the fishing year and when the biomass survey is started, productivity and catch cancel each other.

**Table 7: CAY estimates (*t*) for COC 3. WI = Waitati Inlet, PI = Papanui Inlet, WIc and PIc are estimates for commercial areas only,  $B_{beg}$  = Projected biomass at the beginning of the fishing year.**

Year	<i>M</i>	$F_{0.1}$	$\geq$ SL (mm)	WI		WIc		PI		PIc		Reference
				$B_{beg}$	CAY	$B_{beg}$	CAY	$B_{beg}$	CAY	$B_{beg}$	CAY	
2007	0.2	0.2899	28	8378	1920	5261	1206					Stewart 2008a
2007	0.3	0.3863	28	8378	2342	5261	1471					Stewart 2008a
2007	0.4	0.5537	28	8378	2990	5261	1878					Stewart 2008a
2007	0.2	0.2899	30	7106	1629	4725	1083					Stewart 2008a
2007	0.3	0.3863	30	7106	1986	4725	1321					Stewart 2008a
2007	0.4	0.5537	30	7106	2536	4725	1686					Stewart 2008a
2004	0.2	0.2321	30	9399	1771	6081	1146	4119	776	2454	462	Stewart 2005
2004	0.3	0.3412	30	9399	2367	6081	1532	4119	1038	2454	618	Stewart 2005
2004	0.4	0.4767	30	9399	2984	6081	1930	4119	1308	2454	779	Stewart 2005
2002	0.2	0.2017	30	7183	1193	5364	891	3860	641	2322	386	Wing <i>et al.</i> 2002
2002	0.3	0.3015	30	7183	1627	5364	1215	3860	874	2322	526	Wing <i>et al.</i> 2002
2002	0.4	0.3956	30	7183	1960	5364	1464	3860	1053	2322	634	Wing <i>et al.</i> 2002
1999	0.2	0.258	30	7235	1498			3990	826			Breen <i>et al.</i> 1999
1999	0.3	0.357	30	7235	1848			3990	1019			Breen <i>et al.</i> 1999
1999	0.4	0.457	30	7235	2221			3990	1225			Breen <i>et al.</i> 1999

### 3.4 Other factors

Commercial, customary and recreational fishers target different sized cockles. Biomass and yield estimates will differ for different sizes of recruitment to the fishery. Maori and recreational fishers prefer larger cockles (>45 mm shell length and greater) whereas commercial fishers currently prefer cockles of around 28–34 mm. Estimates of yields have been estimated for size of recruitment at  $\geq 28$  mm; however, these estimates do not consider multiple fisheries preferring different sized cockles. Depending on the management approach taken in the future in COC 3, the appropriateness of the current methods to estimate yield may need to be reviewed.

The yield estimates use information from yield-per-recruit analyses that assume constant recruitment, and constant growth and mortality rates. Yield estimates will be improved when growth, mortality and recruitment variation are better known.

As cockles become sexually mature at around 18 mm, using a size of recruitment of 30 mm should provide some protection against egg overfishing under most circumstances. Certainly the increase in the biomass of small cockles (>2 to 18 mm) seen in both inlets in 2004 suggests that the very poor recruitment observed by Wing *et al.* (2002) may have been due to natural variability, and supports the conjecture that significant recruitment might occur only sporadically in the Otago fishery, as suggested by John Jillett (*pers. comm.*) and Breen *et al.* (1999). The possibility that fishing has an effect on recruitment remains an unknown.

In other cockle fisheries it has been shown that recruitment of juvenile cockles can be reduced by the removal of a large proportion of adult cockles from a given area of substrate. This would suggest that there is some optimal level of adult biomass to facilitate recruitment, although its value is not known. To date it has not been determined whether the cockles being targeted by commercial harvesting in the Otago fishery comprise the bulk of the spawning stock or if disturbance of the cockle beds is influencing settlement.

The distribution of very small size classes (2 to 10 mm) across the various beds is variable and no consistent differences exist for this size of shellfish between commercial and non-commercial beds (Stewart 2008a). A comparison of the size/frequency histograms with fishing history for each bed would be a worthwhile exercise and may reveal more. The fact that the relationship between spawning stock and recruitment in this fishery is poorly understood remains a concern.

The very slight decrease in biomass recorded in the Stewart (2008a) survey suggests that the current level of harvest is sustainable. What is not known is if the decrease in biomass is the beginning of a long-term trend or simply the result of natural variability.

## COCKLES (COC 3)

The impacts of the illegal catch, the Maori traditional catch and incidental handling mortality are unknown, although illegal catch is thought to be insignificant. The impacts of the recreational fishery are probably minor compared with those from the commercial fishery.

### 4. STATUS OF THE STOCKS

#### Stock structure assumptions

Each inlet is assessed separately.

- COC 3

<b>Stock Status</b>																																				
Year of Most Recent Assessment	2007																																			
Assessment Runs Presented	Survey biomass estimate for $\geq 19$ mm shell length																																			
Reference Points	Target: Not defined, but $B_{MSY}$ assumed Soft Limit: 20% $B_0$ Hard Limit: 10% $B_0$																																			
Status in relation to Target	Likely ( $> 60\%$ ) to be at or above the target																																			
Status in relation to Limits	Unlikely ( $< 40\%$ ) to be below both soft and hard limits																																			
<b>Historical Stock Status Trajectory and Current Status</b>																																				
<table border="1"> <caption>Estimated data from the Historical Stock Status Trajectory and Current Status graph</caption> <thead> <tr> <th>Year</th> <th>Waitati Inlet (%)</th> <th>Papanui Inlet (%)</th> <th>Soft Limit (%)</th> <th>Hard Limit (%)</th> </tr> </thead> <tbody> <tr> <td>1984</td> <td>100</td> <td>100</td> <td>20</td> <td>10</td> </tr> <tr> <td>1992</td> <td>85</td> <td>85</td> <td>20</td> <td>10</td> </tr> <tr> <td>1998</td> <td>115</td> <td>95</td> <td>20</td> <td>10</td> </tr> <tr> <td>2002</td> <td>95</td> <td>75</td> <td>20</td> <td>10</td> </tr> <tr> <td>2004</td> <td>105</td> <td>78</td> <td>20</td> <td>10</td> </tr> <tr> <td>2006</td> <td>100</td> <td>-</td> <td>20</td> <td>10</td> </tr> </tbody> </table> <p>Biomass as a proportion of <math>B_0</math> for Waitati and Papanui Inlets, this is estimated from biomass <math>&gt; 19</math> mm.</p>		Year	Waitati Inlet (%)	Papanui Inlet (%)	Soft Limit (%)	Hard Limit (%)	1984	100	100	20	10	1992	85	85	20	10	1998	115	95	20	10	2002	95	75	20	10	2004	105	78	20	10	2006	100	-	20	10
Year	Waitati Inlet (%)	Papanui Inlet (%)	Soft Limit (%)	Hard Limit (%)																																
1984	100	100	20	10																																
1992	85	85	20	10																																
1998	115	95	20	10																																
2002	95	75	20	10																																
2004	105	78	20	10																																
2006	100	-	20	10																																

<b>Fishery and Stock Trends</b>	
Recent Trend in Biomass or Proxy	Biomass at Waitati Inlet has been stable and has never decreased below 85% of $B_0$ . At Papanui Inlet biomass generally decreased to approximately 70% of $B_0$ in 2004 but little commercial catch has come out of this inlet since.
Recent Trend in Fishing Mortality or Proxy	Exploitation rate has never exceeded 2% for Papanui Inlet and has increased in Waitati Inlet to just over 6%. Exploitation rate is unable to be calculated for Otago Harbour. It is Very Unlikely ( $< 10\%$ ) that overfishing is occurring.



	<p>Exploitation rate as calculated by landings divided by biomass (<math>\geq 19</math> mm) from whole inlets.</p>
Other Abundance Indices	Recruited biomass in the two currently harvested beds in Otago Harbour has declined between 1998 and 2008, prior to the start of harvesting.
Trends in Other Relevant Indicators or Variables	-

<b>Projections and Prognosis</b>	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing decline below Limits	Fishing at present levels is Very Unlikely (< 10%) to cause declines below soft or hard limits
<b>Assessment Methodology</b>	
Assessment Type	Level 2: Partial quantitative stock assessment
Assessment Method	Absolute biomass estimates from quadrant surveys
Main data inputs	Abundance and length frequency information
Period of Assessment	Latest assessment: 2007      Next assessment: unknown
Changes to Model Structure and Assumptions	-
Major Sources of Uncertainty	-

<b>Qualifying Comments</b>
Water quality issues have influenced the amount of time when cockles can be harvested from Papanui Inlet in recent years.

<b>Fishery Interactions</b>
-

**Table 8: Summary of yields, catch limits, and reported landings (t) of COC 3 for the most recent fishing year.**

<u>Fishstock</u>	<u>MCY&gt;28mm SL</u>	<u>CAY&gt;28mm SL</u>	2011-12 <u>Actual TACC</u>	2011-12 <u>Reported Landings</u>
COC 3	N/A	N/A	1470	795

## 5. FOR FURTHER INFORMATION

- Annala J.H., Sullivan K.J. 1996. Report from the mid-year Fishery Assessment Plenary, November 1996: stock assessments and yield estimates. Ministry for Primary Industries.
- Annala J.H., Sullivan K.J., O'Brien C.J., Smith N.W.McL., Grayling S.M. 2003. Report from the Fishery Assessment Plenary, May 2003: stock assessments and yield estimates. Ministry for Primary Industries.
- Boyd R.O., Gowing L., Reilly J.L. 2004. 2000-2001 national marine recreational fishing survey: diary results and harvest estimates. Final Research Report for Ministry of Fisheries project REC2000/03. 81 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Boyd R.O., Reilly J.L. 2002. 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries project REC98/03. 28 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Bradford E. 1998. Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p.

### COCKLES (COC 3)

- Breen P.A., Carbines G.C., Kendrick T.H. 1999. Stock assessment of cockles in Papanui and Waitati Inlets, Otago Harbour, and Purakanui, Otago. Final Report for the Ministry of Fisheries research project COC9701 dated July 1999.
- Cameron T. 1997. Aspects of growth and condition in *Austrovenus stutchburyi* (Finlay 19270 (Bivalvia: Veneridae) at Waitati Inlet: Influence of shore height and distance from mouth. Unpublished MSc thesis, University of Otago, Dunedin, New Zealand.
- Cranfield H.J., Micheal K.P., Francis R.I.C.C. 1996. Growth rates of five species of subtidal clam on a beach in the South Island, New Zealand. *Marine and Freshwater Research* 47: 773-784.
- Francis R.I.C.C. 1984. An adaptive strategy for stratified random trawl surveys. *New Zealand Journal of Marine and Freshwater Research* 18: 59-71.
- Hilborn R., Walters C.J. 1992. *Quantitative Fisheries Stock Assessment: Choice, dynamics and uncertainty*. Chapman and Hall, London.
- Irwin C.R. 1999. The effects of harvesting on the reproductive and population biology of the New Zealand Littleneck Clam (*Austrovenus stutchburyi*) in Waitati Inlet. Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- McKinnon J. 1996. Studies of the age, growth and shell increment patterns in the New Zealand cockle (*Austrovenus stutchburyi*). Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- Manly B.F.J., Akroyd J.M., Walshe K.A.R. 2002. Two-phase stratified random surveys on multiple populations at multiple locations. *New Zealand Journal of Marine and Freshwater Research* 36: 581-591.
- Morton J., Miller M. 1973. *The New Zealand Sea Shore*. Collins, Auckland. 653 p.
- Stewart B., Keogh J., Fletcher D., Mladenov P. 1992. Biomass survey of the New Zealand littleneck clam (*Chione stutchburyi*) in Papanui and Waitati Inlets, Otago during 1991/1992. Marine Science and Aquaculture research Centre, University of Otago, Dunedin, New Zealand. 37 p.
- Stewart B. 2004. Warrington treated Sewage Discharge: Waitati Inlet Ecological Survey, 2004. Prepared for DCC. 21 p.
- Stewart B. 2005. Stock assessment of Cockles (*Austrovenus stutchburyi*) in Papanui and Waitati Inlets, Otago, 2004. Final Research Report for the Ministry of Fisheries Research Project COC2004/02. 54 p.
- Stewart B. 2008a. Stock assessment of clams (*Austrovenus stutchburyi*) in Waitati Inlet, Otago, 2007. Final Research Report for Southern Clams Ltd. 24 p.
- Stewart B.G. 2008b. Clam (*Austrovenus stutchburyi*) resource and habitat survey in Otago Harbour (COC 3), Otago, 2008. Report prepared for Southern Clams Ltd. By Ryder Consulting.
- Sukhatme P.V. 1954. *Sampling theory of surveys: with applications*. New Delhi : Indian Society of Agricultural Statistics. 49 p.
- Teirney L.D., Kilner A.R., Millar R.E., Bradford E., Bell J.D. 1997. Estimation of recreational catch from 1991/92 to 1993/94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p.
- Wildish K. 1984a. The cockle resource in the Otago region: 1. An initial study of the New Zealand cockle (*Chione stutchburyi*) resource in Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries. Now Ministry for Primary Industries.
- Wildish K. 1984b. The cockle resource in the Otago region: further analysis of results from the survey of *Chione stutchburyi* (the New Zealand cockle) populations at Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries.
- Wing S., Irwin C., Granger G. 2002. Biomass survey and yield estimates for the New Zealand littleneck clam *Austrovenus stutchburyi* in Papanui and Waitati Inlets, Otago. Final Report for the Ministry of Fisheries Research Project COC2001/02. 52 p.
- Zar J.H. 1996. *Biostatistical Analysis*. Third Edition. Prentice Hall International Inc.