


RIBALDO (RIB)*(Mora moro)***1. FISHERY SUMMARY****1.1 Commercial fisheries**

In New Zealand ribaldo is caught mainly on bottom longlines and as a bycatch of trawling. Up to 4500 t were reported in 1977 by Japanese and Korean longline vessels target fishing for ling on the Chatham Rise and east coast of the South Island in the 1970s. Since 1982-83, overall reported catch has been mainly from the Chatham Rise and east coast South Island (QMAs 3 & 4) but has declined somewhat from these areas since being introduced into the QMS in the 1998-99 fishing year. Since entering the QMS, a similar decline in reported ribaldo catch is seen in other QMAs with the exception of RIB 7 where reported catches have been increasing. RIB 7 now has the highest reported catch of any QMA. The reasons for these changes in catch levels are not well understood as ribaldo is mainly taken as bycatch. Levels of discarding and unreported catch are likely to have changed with the introduction of ribaldo into the QMS. Ribaldo are caught throughout the New Zealand Exclusive Economic Zone by a variety of fishing methods in different target fisheries but mainly as bycatch in bottom trawls targeting hoki (*Macruronus novaezelandiae*), hake (*Merluccius australis*) and ling (*Genypterus blacodes*) and bottom longlines for ling.

There is no seasonality of catch other than on the west coast South Island which is related to target fishing of hoki and hake during the winter spawning season. Catches by Japanese and Korean longliners in the mid 1970s are shown in Table 1. Landings from 1982-83 onwards are shown in Table 2, while Figure 1 shows the historical landings and TACC values for the main RIB stocks.

Table 1: Japanese and Korean longline catch (t) of ribaldo (“deep-sea cod”¹) from New Zealand waters, probably mostly Chatham Rise and east coast South island, by calendar year from 1975-77.

Year	1975	1976	1977
Japan	2 417	4 920	4 283
Korea	-	-	2 86

1. Reported as “cods” but considered to be mainly ribaldo. The Korean fleet began fishing in April 1977.

Ribaldo was introduced into the QMS from 1 October 1998, no customary, recreational or other mortality allowances have been given. Historical catch limits up to the most recent fishing year (2011-12) are shown in Table 2. TACCs were increased from 1 October 2006 in RIB 6 to 231 t and in RIB 7 to 330 t. In these stocks landings were above the TACC for a number of years and the TACCs have

been increased to the average of the previous 7 years plus an additional 10%. Current levels of reported catch are well below TACCs apart from in RIB 7 where they have continued to exceed the TACC, with the exception of the most recent fishing year.

Table 2: Reported landings (t) of ribaldo by QMA for fishing years 1983-84 to 2011-12 and TACCs (t). QMA 10 has no catches and a TACC of 0. Total includes catches from outside the NZ EEZ.

	QMA 1		QMA 2		QMA 3		QMA 4		QMA 5	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1982-83	0		8		15		33		111	
1983-84	0		3		24		21		68	
1984-85	0		4		17		61		21	
1985-86	1		1		26		13		35	
1986-87	4		1		44		20		41	
1987-88	19		4		65		31		56	
1988-89	1		2		33		41		6	
1989-90	8		9		23		28		6	
1990-91	15		15		177		119		34	
1991-92	95		40		160		169		73	
1992-93	131		54		217		228		67	
1993-94	87		70		217		186		23	
1994-95	116		136		437		303		68	
1995-96	121		168		286		253		26	
1996-97	114		188		365		843		64	
1997-98	78		122		141		375		80	
1998-99	24	121	55	176	161	394	290	357	71	52
1999-00	22	121	89	176	264	394	347	357	80	52
2000-01	5	121	107	176	269	394	306	357	78	52
2001-02	7	121	53	176	198	394	370	357	62	52
2002-03	12	121	98	176	211	394	183	357	50	52
2003-04	12	121	120	176	175	394	299	357	50	52
2004-05	28	121	127	176	156	394	379	357	44	52
2005-06	49	121	137	176	126	394	202	357	47	52
2006-07	39	121	125	176	149	394	312	357	49	52
2007-08	53	121	135	176	134	394	173	357	43	52
2008-09	45	121	74	176	216	394	216	357	31	52
2009-10	28	121	63	176	213	394	162	357	27	52
2010-11	42	121	67	176	348	394	137	357	30	52
2011-12	29	121	27	176	174	394	304	357	32	52

	QMA 6		QMA 7		QMA 8		QMA 9		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1982-83	0		58		0		0		225	
1983-84	1		25		0		0		142	
1984-85	13		18		0		0		134	
1985-86	2		37		0		0		115	
1986-87	10		6		0		0		126	
1987-88	12		68		0		0		255	
1988-89	6		69		1		10		169	
1989-90	13		21		0		0		108	
1990-91	106		55		0		0		521	
1991-92	98		40		0		0		675	
1992-93	96		106		0		0		899	
1993-94	92		42		1		0		718	
1994-95	122		39		2		6		1 231	
1995-96	109		62		0		0		1 025	
1996-97	158		77		1		0		1 824	
1997-98	262		110		1		1		1 214	
1998-99	223	124	243	55	1	1	0	2	1 081	1 282
1999-00	237	124	300	55	< 1	1	< 1	2	1 359	1 282
2000-01	191	124	275	55	< 1	1	< 1	2	1 242	1 282
2001-02	322	124	254	55	0	1	< 1	2	1 311	1 282
2002-03	172	124	338	55	< 1	1	1	2	1 209	1 282
2003-04	205	124	364	55	< 1	1	2	2	1 302	1 282
2004-05	105	124	307	55	< 1	1	2	2	1 240	1 282
2005-06	62	124	336	55	0	1	4	2	1 018	1 282
2006-07	61	231	404	330	0	1	9	2	1 162	1 664
2007-08	80	231	356	330	< 1	1	14	2	992	1 664
2008-09	63	231	456	330	< 1	1	10	2	1 111	1 664
2009-10	104	231	137	330	< 1	1	21	2	755	1 664
2010-11	67	231	198	330	3	1	20	2	913	1 664
2011-12	76	231	177	330	3	1	12	21	835	1 683

RIBALDO (RIB)




Figure 1: Historical landings and TACC for the seven main RIB stocks. From top to bottom: RIB1 (Auckland East), RIB2 (Central East) and RIB3 (South East Coast). [Continued on next page].




Figure 1 [Continued]: Historical landings and TACC for the seven main RIB stocks. From top to bottom: RIB4 (South East Chatham Rise), RIB5 (Southland) and RIB6 (Sub-Antarctic). [Continued on next page].

RIBALDO (RIB)




Figure 1 [Continued]: Historical landings and TACC. Above: RIB7 (Challenger).

In RIB 1, ribaldo are taken as bycatch primarily in the ling and to lesser extent bluenose bottom longline fisheries. There is also some direct targeting of ribaldo by bottom longline. In RIB 2, ribaldo are taken as bycatch primarily in the ling and bluenose bottom longline fisheries and to a lesser extent the hoki and orange roughy bottom trawl fisheries. There is also some direct targeting of ribaldo by bottom longline. In RIB 8 very small catches (< 1 t) of ribaldo are reported mainly in bottom trawl and bottom longline fisheries for a variety of target species. There is no reported targeting of ribaldo in RIB 8 and since entering the QMS in the 1998-99 fishing year the TACC of one tonne has never been caught. In RIB 9 very small amounts of ribaldo are taken as bycatch in orange roughy, cardinal and alfonsino target trawl fisheries and in the ling bottom longline fishery. In all areas, a variety of other fishing methods and target fisheries also report catching ribaldo but only in negligible amounts. The majority of the ribaldo catch is taken in RIB 3-7. Fisheries interactions for these areas are described in the Status of the Stocks tables in Section 5.

1.2 Recreational fisheries

There are no known recreational fisheries for ribaldo.

1.3 Customary non-commercial fisheries

There is no known customary non-commercial fishing for ribaldo.

1.4 Illegal catch

Estimates of illegal catch are not available.

1.5 Other sources of mortality

There is no quantitative information on the level of other sources of mortality.

2. BIOLOGY

Ribaldo is known from the North Atlantic Ocean from Iceland to West Africa, the western Mediterranean Sea, the Indian Ocean south of Madagascar and the Pacific Ocean from Australia, New Zealand and Chile. In New Zealand it is widespread and has been caught by research trawl at depths of about 200-1300 m. It appears to be most common at 500-1000 m. The relatively high catch by bottom longline suggests that it favours rough bottom habitats.

Ribaldo reaches fork lengths (FL) of about 75 cm and 65 cm for females and males respectively. Most research trawls have caught fish ranging from 30 to 70 cm FL. The 50% length at sexual maturity has been estimated at 45 cm total length for New Zealand ribaldo (O'Driscoll *et al.* 2003). Analyses of

data on female gonad development, collected by the Ministry of Fisheries Observer Programme, indicates a winter/early spring spawning season. Fish do not appear to form large spawning aggregations. Locations at which spawning fish have been observed are the upper North Island (extending outside the EEZ), north-east and west Chatham Rise, the area between the Snares and Auckland Islands shelves, and west coast of the South Island. Early life history is largely unknown but a few individuals less than 10 cm FL were captured in plankton nets in the upper 200 m of the water column over bottom depths of about 1000 m at the south west end of Chatham Rise. The distribution of juveniles < 28 cm is similar to that of observed spawning females. Juveniles up to 35 cm have been observed in all fished areas of the EEZ except for the Bounty Islands.

Ageing by zone counts of otoliths has been validated using radiometric techniques (Sutton *et al.* 2010) using ribaldo caught on Chatham Rise trawl surveys by *Tangaroa* from 2001 to 2005. Maximum observed ages were 37 and 39 years for females and males respectively. Von Bertalanffy growth parameters are presented in Table 3, estimates of natural mortality (*M*) are presented in Table 4 and length-weight parameters in Table 5.

Ribaldo are caught in low numbers both in research trawl surveys and in observed commercial fisheries making tracking of cohorts by length frequencies difficult. Collection of otoliths is required to construct catch-at-age series for any fishing areas. Analyses of trawl survey and observer data has shown that biomass of females is usually greater than that of males on the Chatham Rise although sex ratios are about 1:1. In the SubAntarctic the biomass and numbers of females are significantly greater than males, often over 10:1. Sex ratios elsewhere in the EEZ are less clear.

Table 3: Von Bertalanffy growth parameter values for ribaldo. Source: Sutton *et al.* 2010.

Von Bertalanffy growth parameters			
	<i>K</i>	<i>t</i> ₀	<i>L</i> _∞
RIB 3 & 4 females	0.135	0.221	67.526
RIB 3 & 4 males	0.072	-5.246	61.444
RIB 3 & 4 combined sexes	0.14	-0.287	60.47

Table 4: Estimates of natural mortality (*M*). Source: Sutton *et al.* 2010.

	Females	Males
Natural mortality (<i>M</i>)	0.106	0.112

Table 5: Length-weight parameter values for ribaldo.

Fishstock			Estimate		Source
Weight = a(length) ^b (Weight in g, length in cm total length)					
		Females		Males	
		a	b	a	b
RIB 3 & 4	0.0037	3.27	0.0053	3.18	Sutton <i>et al.</i> (2010)
RIB 5 & 6	-	-	-	-	
		Sexes combined			
		a	b		
RIB 3 & 4		0.004289	3.237753		Sutton <i>et al.</i> (2010)
RIB 5 & 6		0.0039	3.15		Bagley <i>et al.</i> (unpublished data)

3. STOCKS AND AREAS

It is not known whether different regional stocks of ribaldo occur in New Zealand waters but it is possible that there are separate stocks based on natural bathymetric boundaries. The Working Group had previously agreed on five fishstocks based on the four main fishing areas plus the Kermadec area, i.e., the east coast of the North Island (QMAs 1 and 2), Chatham Rise and east coast South Island (QMAs 3 and 4), Southland and Sub-Antarctic (QMAs 5 and 6), the west coast of New Zealand (QMAs 7, 8 and 9) and QMA 10. A review of all available information in 2010 indicated that the main fishing areas are still as found previously. The review also indicated spawning activity in all areas, except RIB 8 and RIB 10 (for which there is no information). This is not inconsistent with the

RIBALDO (RIB)

management of the fishery by the current 10 FMAs. Highly skewed sex ratios in the SubAntarctic have unknown implications for stock structure.

4. STOCK ASSESSMENT

The Middle Depths Working Group agreed in February 2011 that relative biomass estimates of ribaldo from middle depth trawl surveys on the Chatham Rise and the SubAntarctic were suitable for monitoring major changes in ribaldo abundance for RIB 3 & 4 and RIB 5 & 6 respectively. Standardised CPUE indices from the spawning hoki and hake target fisheries in RIB 7 were not accepted although this could possibly be revisited in future using daily processed catch of ribaldo instead of estimated catch at the tow by tow level. There are no stock monitoring indices available for RIB 1, 2, 8 & 9

4.1 Estimates of fishery parameters and abundance

No estimates of fishery parameters have been made. Analyses of research trawl survey abundance estimates on the Chatham Rise and Sub-Antarctic suggest they may be suitable for monitoring major changes in abundance. While the Sub-Antarctic survey does not fully cover the depth range of ribaldo the Chatham Rise survey has been extended down to 1300 m since 2010. Biomass from the deep strata (800-1300 m) accounted for 23% of the total ribaldo biomass in 2010.

Dunn (2006) described the fishery up to 2002-03 but found that interpreting CPUE data was difficult because ribaldo are taken mainly as by-catch of other fisheries and the CPUE may reflect changes in those fisheries rather than ribaldo abundance. MacGibbon & Hurst (2011) carried out a standardised trawl CPUE analysis of hoki and hake in RIB 7 from 1999-2009. The index was problematic for the same reasons listed above. The trend showed a slight decline from 1999 to 2002 and was fairly steady thereafter.

4.2 Biomass estimates

Estimates of biomass are given in Table 6.

4.3 Yield estimates and projections

MCY cannot be estimated.

CAY cannot be estimated.

4.5 Other yield estimates and stock assessment results

No information is available.

Table 6: Biomass indices (t) and coefficients of variation (CV) of ribaldo from *Tanagroa* trawl surveys (Assumptions: areal availability, vertical availability and vulnerability = 1). NB: estimates are for the core strata only for the respective time series.

Vessel	Trip code	Date	Biomass (t)	%CV
<u>Chatham Rise (RIB 3 & 4)</u>				
<i>Tangaroa</i>	TAN9106	Dec 91-Feb 92	417	12.2
	TAN9212	Dec 92-Feb 93	336	17.2
	TAN9401	Jan-94	602	10.8
	TAN9501	Jan-Feb 95	406	19.7
	TAN9601	Dec 95-Jan 96	470	18.2
	TAN9701	Jan-Jan 97	333	21.3
	TAN9801	Jan-Jan 98	510	14.3
	TAN9901	Jan-Jan 99	395	18
	TAN0001	Dec 99-Jan 00	387	20.8
	TAN0101	Dec 00-Jan 01	762	18.3
	TAN0201	Dec 01-Jan 02	417	13.2
	TAN0301	Dec 02-Jan 03	455	18.1
	TAN0401	Dec 03-Jan 04	535	15.6
	TAN0501	Dec 04-Jan 05	491	14.2
	TAN0601	Dec 05-Jan 06	313	16.9
	TAN0701	Dec 06-Jan 07	380	15
	TAN0801	Dec 07-Jan 08	479	14.3
	TAN0901	Dec 08-Jan 09	463	12.7
	TAN1001	Jan-10	416	19.9
	TAN1101	Jan-11	396	16.7
	TAN1201	Jan-12	469	14.6
	TAN1301	Jan-13	428	15.7
<u>SubAntarctic (summer, RIB 5 & 6)</u>				
<i>Tangaroa</i>	TAN9105	Nov-Dec 91	1035	11.2
	TAN9211	Nov-Dec 92	389	18.6
	TAN9310	Nov-Dec 93	996	12.8
	TAN0012	Nov-Dec 00	873	14
	TAN0118	Nov-Dec 01	1 017	17.2
	TAN0219	Nov-Dec 02	656	17.5
	TAN0317	Nov-Dec 03	653	18.9
	TAN0414	Nov-Dec 04	951	16.5
	TAN0515	Nov-Dec 05	721	14.6
	TAN0617	Nov-Dec 06	780	16.4
	TAN0714	Nov-Dec 07	1 062	13.5
	TAN0813	Nov-Dec 08	658	18
	TAN0911	Nov-Dec 09	1 056	13.4
	TAN1117	Nov-Dec 11	1 017	17.2
	TAN1215	Nov-Dec 12	787	16.7
<i>Tangaroa</i>	TAN9204	Apr-May 92	768	17.1
	TAN9304	May-Jun 93	1 162	15.1
	TAN9605	Mar-Apr 96	989	16.7
	TAN9805	Apr-May 98	837	14.2

RIBALDO (RIB)




Figure 2: Doorspread biomass estimates of ribaldo by sex from the Chatham Rise 1991 to 2011 (upper) and Sub-Antarctic 1991 to 1993 and 2000 to 2009 (lower), from *Tangaroa* trawl surveys.

5. STATUS OF THE STOCKS

• RIB 1, 2, 7, 8 and 9

There are no accepted stock monitoring indices available for RIB 1, 2, 7, 8 & 9.

• RIB 3 & 4

Stock Status	
Year of Most Recent Assessment	2011
Reference Points	Target: Not established but 40% B_0 assumed Soft Limit: 20% B_0

	Hard Limit: 10% B_0
Status in relation to Target	Unknown
Status in relation to Limits	Unlikely (< 40%) to be below soft limit Unlikely (< 40%) to be below hard limit
Historical Stock Status Trajectory and Current Status	
<p>Doorspread biomass estimates of ribaldo (error bars are \pm two standard deviations) from the Chatham Rise, from <i>Tangaroa</i> surveys from 1991 to 2011.</p>	

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The relative biomass index of ribaldo from summer middle depth trawl surveys of the Chatham Rise is relatively flat. Precision is generally good in this time series (< 20%). Although numbers of individual ribaldo caught are low the Working Group considered this index to be suitable to monitor major trends in this stock.
Recent Trend in Fishing Mortality or Proxy	-

Projections and Prognosis	
Stock Projections or Prognosis	Stock size is Likely (> 60%) to remain near current levels under current catches.
Probability of Current Catch or TACC causing decline below Limits	Soft limit: Unlikely (< 40%) for current catches Hard limit: Unlikely (< 40%) for current catches

Assessment Methodology	
Assessment Type	Level 2: Partial quantitative stock assessment
Assessment Method	Evaluation of agreed trawl survey indices thought to index RIB 3 & 4 abundance
Main data inputs	Data collected on trawl surveys
Period of Assessment	Latest assessment: 2011 Next assessment: 2012
Changes to Model Structure and Assumptions	-
Major Sources of Uncertainty	- Low numbers of individuals caught on trawl surveys; proportion of biomass deeper than 800 m (not surveyed in core strata)
Qualifying Comments	
-	

RIBALDO (RIB)

Fishery Interactions


In RIB 3 & 4, ribaldo are taken as bycatch primarily in the ling and hoki bottom trawl fisheries and ling bottom longline fishery.

• RIB 5 & 6

Stock Status

Year of Most Recent Assessment	2011
Reference Points	Target: Not established but 40% B_0 assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0
Status in relation to Target	Unknown
Status in relation to Limits	Unlikely (< 40%) to be below soft limit Unlikely (< 40%) to be below hard limit

Historical Stock Status Trajectory and Current Status



Doorspread biomass estimates of ribaldo (error bars are \pm two standard deviations) from the Sub-Antarctic, from *Tangaroa* surveys from 1991 to 1993, and 2000 to 2009.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	Relative biomass estimates of ribaldo from summer middle depth surveys of the SubAntarctic show a relatively flat index. CVs are consistently low in this time series (< 20%). Although numbers of individual ribaldo caught are low the Working Group considered this index to be suitable to monitor major trends in this stock.
----------------------------------	--

Recent Trend in Fishing Mortality or Proxy	Unknown
--	---------

Projections and Prognosis

Stock Projections or Prognosis	Stock size is Likely (> 60%) to remain near current levels under current catches and TACCs.
Probability of Current Catch or TACC causing decline below Limits	Soft limit: Unknown Hard limit: Unlikely (< 40%)

Assessment Methodology

Assessment Type	Level 2: Partial quantitative stock assessment
Assessment Method	Evaluation of agreed trawl survey indices thought to index RIB 5 & 6 abundance
Main data inputs	Data collected on trawl surveys

Period of Assessment	Latest assessment: 2011	Next assessment: 2014
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	Low numbers of individuals caught on trawl surveys; proportion of biomass deeper than 800 m (not surveyed in core strata); and unknown implications of highly skewed sex ratios (females usually make up > 90% of biomass) for stock structure. Observer data also shows skewed sex ratios in favour of females.	

Qualifying Comments
-

Fishery Interactions
In RIB 5 & 6, ribaldo are mainly caught as bycatch in hoki and ling bottom trawl fisheries and ling bottom longline fisheries.

TACCs and reported landings for the 2011-12 fishing year are summarised in Table 7.

Table 7: Summary of TACCs (t) and reported landings (t) of ribaldo for the most recent fishing year.

Fishstock		QMA	2011-12 Actual TACC	2011-12 Estimated landings
RIB 1	Auckland (East)	1	121	29
RIB 2	Central (East)	2	176	27
RIB 3	South-east (Coast)	3	394	174
RIB 4	South-east (Chatham)	4	357	304
RIB 5	Southland	5	52	32
RIB 6	Sub-Antarctic	6	231	76
RIB 7	Challenger	7	330	177
RIB 8	Central (West)	8	1	3
RIB 9	Auckland (West)	9	21	12
RIB 10	Kermadec	10	0	0
Total			1 683	835

6. FOR FURTHER INFORMATION

- Bagley N.W., O'Driscoll R.L. 2012. Trawl survey of middle depth species in the Southland and Sub-Antarctic areas, November–December 2009 (TAN0911). *New Zealand Fisheries Assessment Report 2012/05*. 70p.
- Cohen D.M., Inada T., Iwamoto T., Scialabba N. 1990. FAO species catalogue. Vol. 10. Gadiform fishes of the world (Order Gadiformes). An annotated and illustrated catalogue of cods, hakes, grenadiers and other gadiform fishes known to date. Food and Agriculture Organisation Fisheries Synopsis. No. 125, Vol. 10. Rome, FAO. 442p.
- Dunn M.R. 2006. Descriptive and catch per unit effort analyses for New Zealand ribaldo fisheries for the fishing years 1988-78 to 2002-03> New Zealand Fisheries Assessment Document 2006/22. 55p.
- Elder R.D., Taylor J.L. (Comps.) 1979. Prospects and problems for New Zealand's demersal fisheries. Proceedings of the Demersal Fisheries Conference October 1978. Fisheries Research Division Occasional Publication No. 19. 123p.
- MacGibbon D.J., Hurst R.J. 2011. Fishery characterisation and standardised CPUE analyses for ribaldo, *Mora moro*, (Risso, 1810) (Moridae), 1989-90 to 2008-09. New Zealand Fisheries Assessment Report 2011/XX
- McMillan P.J., Hart A.C. 1998. Summary of biology and commercial landings, and a stock assessment of ribaldo *Mora moro* (Risso, 1810), in New Zealand waters. New Zealand Fisheries Assessment Research Document 1998/9. 16p.
- O'Driscoll R.L., Booth J.D., Bagley N.W., Anderson O.F., Griggs L.H., Stevenson M.L., Francis M.P. 2003. Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand deepwater fish, pelagic fish, and invertebrates. NIWA Technical Report 119. 377 p.
- Sutton C.P., Tracey D.M., Andrews A.H., Hart A.C., MacGibbon D.J. 2010. Validated age and growth of ribaldo (*Mora moro*). *New Zealand Fisheries Assessment Report 2010/24*. 26 p.