


SCHOOL SHARK (SCH)

(*Galeorhinus galeus*)
Tupere, Tope, Makohuarau

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

This moderate-sized shark has supported a variety of fisheries around New Zealand from the early 1940s onwards. Landings rose steeply from the late 1970s until 1983 (Table 1), with the intensification of setnets targeting this and other species, and a general decline in availability of other, previously more desirable, coastal species. However, because of the earlier discarding and under-reporting, this recorded rise in landings does not reflect an equal rise in catches. After a small decline in 1984-85, catches decreased by about 50% from 1986 onwards because of reduced quotas within the QMS (Table 2). From 1987-88 to 1991-92 total reported landings were around 2200-2500 t. In 1995-96 total landings increased markedly to 3387 t and the total TACC (3107 t) was exceeded for the first time. Landings have remained around the TACC level since 1995-96. TACCs for SCH 3, 5, 7 & 8 were increased by between 5% (SCH 5) and 20% (the remainder) under AMP management in October 2004. From the 1 October 2007 the TACC for SCH 1 was increased to 689 t, at that time a TAC was set for the first time at 893 t with 102 t, 68 t and 34 t being allocated to customary, recreational and other sources of motility respectively. In 2004 SCH 3, 5, 7 & 8 were allocated an equal recreational and customary non-commercial catch of 48 t, 7 t, 58 t, and 21 t respectively and other sources of mortality were allocated 19 t, 37 t, 32 t, and 26 t respectively. All AMP programmes ended on 30th September 2009. School shark were added to the 6th schedule on the 1st of January 2013, this means that school shark that are alive and likely to survive can be released. Table 1 shows the historical landings and TACC values for the main SCH stocks.

Table 1: Reported domestic landings (t) of school shark from 1948 to 1983.

| Year | Landings | Year | Landings | Year | Landings | Year | Landings |
|------|----------|------|----------|------|----------|------|----------|
| 1948 | 75 | 1957 | 301 | 1966 | 316 | 1975 | 518 |
| 1949 | 124 | 1958 | 323 | 1967 | 376 | 1976 | 914 |
| 1950 | 147 | 1959 | 304 | 1968 | 360 | 1977 | 1 231 |
| 1951 | 157 | 1960 | 308 | 1969 | 390 | 1978 | 161 |
| 1952 | 179 | 1961 | 362 | 1970 | 450 | 1979 | 481 |
| 1953 | 142 | 1962 | 354 | 1971 | 597 | 1980 | 1 788 |
| 1954 | 185 | 1963 | 380 | 1972 | 335 | 1981 | 2 716 |
| 1955 | 180 | 1964 | 342 | 1973 | 400 | 1982 | 2 965 |
| 1956 | 164 | 1965 | 359 | 1974 | 459 | 1983 | 3 918 |

Source: MPI data.

During the period of high landings in the mid 1980s setnetting was the main fishing method, providing about half the total catch, with lining accounting for one-third of the catch, and trawling the remainder. There were large regional variations.

Small amounts of school shark are also caught by the foreign charter tuna longliners fishing offshore in the EEZ to well beyond the shelf edge.

The Banks Peninsula Marine Mammal Sanctuary was established in 1988 by the Department of Conservation under the Marine Mammal Protection Act 1978, for the purpose of protecting Hector's dolphins. The sanctuary extends 4 nautical miles from the coast from Sumner Head in the north to the Rakaia River mouth in the south. Prior to 1 October 2008, no setnets were allowed within the sanctuary between 1 November and the end of February. For the remainder of the year, setnets were allowed; but could only be set from an hour after sunrise to an hour before sunset, be no more than 30 metres long, with only one net per boat which was required to remain tied to the net while it was set.

Voluntary setnet closures were implemented by the SEFMC from 1 October 2000 to protect nursery grounds for rig and elephantfish and to reduce interactions between commercial setnets and Hector's dolphins in shallow waters. The closed area extended from the southernmost end of the Banks Peninsula Marine Mammal Sanctuary to the northern bank of the mouth of the Waitaki River. This area was closed permanently for a distance of 1 nautical mile offshore and for 4 nautical miles offshore for the period 1 October to 31 January.

From 1 October 2008, a suite of regulations intended to protect Maui's and Hector's dolphins was implemented for all of New Zealand by the Minister of Fisheries.

For SCH 1, there have been two recent changes to the management regulations affecting setnet fisheries which take school shark off the west coast of the North Island. The first was a closure to setnet fishing from Maunganui Bluff to Pariokariwa Point for a distance of 4 nautical miles on 1 October 2003. This closure was extended by the Minister to 7 nautical miles on 1 October 2008. An appeal was made by affected fishers who were granted interim relief by the High Court, allowing setnet fishing beyond 4 nautical miles during daylight hours between 1 October to 24 December.

For SCH 3, commercial and recreational set netting was banned in most areas to 4 nautical miles offshore of the east coast of the South Island, extending from Cape Jackson in the Marlborough Sounds to Slope Point in the Catlins. Some exceptions were allowed, including an exemption for commercial and recreational set netting to only one nautical mile offshore around the Kaikoura Canyon, and permitting setnetting in most harbours, estuaries, river mouths, lagoons and inlets except for the Avon-Heathcote Estuary, Lyttelton Harbour, Akaroa Harbour and Timaru Harbour. In addition, trawl gear within 2 nautical miles of shore was restricted to flatfish nets with defined low headline heights.

For SCH 5, commercial and recreational setnetting was banned in most areas to 4 nautical miles offshore, extending from Slope Point in the Catlins to Sandhill Point east of Fiordland and in all of Te Waewae Bay. An exemption which permitted setnetting in harbours, estuaries and inlets was allowed. In addition, trawl gear within 2 nautical miles of shore was restricted to flatfish nets with defined low headline heights.

For SCH 7, both commercial and recreational setnetting were banned to 2 nautical miles offshore, with the recreational closure effective for the entire year and the commercial closure restricted to the period 1 December to the end of February. The closed area extends from Awarua Point north of Fiordland to the tip of Cape Farewell at the top of the South Island. There is no equivalent closure in SCH 8, with the southern limit of the Maui's dolphin closure beginning north of New Plymouth at Pariokariwa Point. There have been two recent changes to the management regulations affecting setnet fisheries which take school shark off the west coast of the North Island.

1.2 Recreational fisheries

Although school shark is a listed gamefish and is regularly caught by recreational fishers, it is not considered to be a particularly desirable target species at the present time. Recreational catch records have been obtained from diary surveys undertaken in 1991-94, 1996 and 1999-00 (Tables 3 and 4).

SCHOOL SHARK (SCH)

Table 2: Reported landings (t) of school shark by Fishstock from 1983-84 to 2011-12 and actual TACCs (t) from 1986-87 to 2011-12. QMS data from 1986-present.

| Fishstock FMA (s) | SCH 1 1 & 9 | | SCH 2 2 | | SCH 3 3 | | SCH 4 4 | | SCH 5 5 & 6 | |
|----------------------|----------------|------|------------|------|------------|------|------------|------|----------------|------|
| | Landings | TACC | Landings | TACC | Landings | TACC | Landings | TACC | Landings | TACC |
| 1983-84* | 1 087 | - | 298 | - | 630 | - | 8 | - | 792 | - |
| 1984-85* | 861 | - | 237 | - | 505 | - | 12 | - | 995 | - |
| 1985-86* | 787 | - | 214 | - | 370 | - | 23 | - | 647 | - |
| 1986-87 | 418 | 560 | 137 | 160 | 283 | 270 | 19 | 200 | 382 | 610 |
| 1987-88 | 530 | 604 | 123 | 168 | 320 | 289 | 22 | 200 | 529 | 613 |
| 1988-89 | 483 | 624 | 134 | 188 | 222 | 294 | 25 | 200 | 494 | 615 |
| 1989-90 | 585 | 652 | 154 | 197 | 272 | 305 | 27 | 235 | 450 | 635 |
| 1990-91 | 559 | 664 | 139 | 198 | 227 | 318 | 21 | 239 | 480 | 649 |
| 1991-92 | 596 | 664 | 161 | 198 | 264 | 318 | 34 | 239 | 612 | 686 |
| 1992-93 | 820 | 664 | 202 | 199 | 220 | 320 | 38 | 239 | 593 | 686 |
| 1993-94 | 658 | 667 | 156 | 199 | 202 | 322 | 41 | 239 | 624 | 686 |
| 1994-95 | 658 | 668 | 159 | 199 | 237 | 322 | 86 | 239 | 656 | 694 |
| 1995-96 | 804 | 668 | 212 | 199 | 296 | 322 | 229 | 239 | 690 | 694 |
| 1996-97 | 793 | 668 | 228 | 199 | 290 | 322 | 179 | 239 | 662 | 694 |
| 1997-98 | 764 | 668 | 214 | 199 | 270 | 322 | 127 | 239 | 623 | 694 |
| 1998-99 | 783 | 668 | 275 | 199 | 331 | 322 | 100 | 239 | 714 | 694 |
| 1999-00 | 820 | 668 | 250 | 199 | 341 | 322 | 97 | 239 | 706 | 694 |
| 2000-01 | 799 | 668 | 178 | 199 | 364 | 322 | 100 | 239 | 724 | 694 |
| 2001-02 | 691 | 668 | 208 | 199 | 324 | 322 | 93 | 239 | 673 | 708 |
| 2002-03 | 689 | 668 | 225 | 199 | 410 | 322 | 130 | 239 | 746 | 708 |
| 2003-04 | 758 | 668 | 187 | 199 | 323 | 322 | 149 | 239 | 727 | 708 |
| 2004-05 | 694 | 668 | 201 | 199 | 424 | 387 | 206 | 239 | 743 | 743 |
| 2005-06 | 634 | 668 | 177 | 199 | 325 | 387 | 183 | 239 | 712 | 743 |
| 2006-07 | 661 | 668 | 200 | 199 | 376 | 387 | 88 | 239 | 738 | 743 |
| 2007-08 | 708 | 689 | 227 | 199 | 345 | 387 | 133 | 239 | 781 | 743 |
| 2008-09 | 713 | 689 | 232 | 199 | 364 | 387 | 145 | 239 | 741 | 743 |
| 2009-10 | 589 | 689 | 213 | 199 | 426 | 387 | 191 | 239 | 784 | 743 |
| 2010-11 | 777 | 689 | 187 | 199 | 366 | 387 | 174 | 239 | 701 | 743 |
| 2011-12 | 689 | 689 | 188 | 199 | 351 | 387 | 201 | 239 | 729 | 743 |

| Fishstock FMA (s) | SCH 7 7 | | SCH 8 8 | | SCH 10 10 | | Total | |
|----------------------|------------|------|------------|------|--------------|------|-----------|-------|
| | Landings | TACC | Landings | TACC | Landings | TACC | Landings§ | TACC |
| 1983-84* | 1 039 | - | 694 | - | 0 | - | 4 776 | - |
| 1984-85* | 1 030 | - | 698 | - | 0 | - | 4 501 | - |
| 1985-86* | 851 | - | 652 | - | 0 | - | 3 717 | - |
| 1986-87 | 454 | 470 | 229 | 310 | 0 | 10 | 1 946 | 2 590 |
| 1987-88 | 515 | 500 | 374 | 345 | 0 | 10 | 2 367 | 2 729 |
| 1988-89 | 532 | 522 | 419 | 433 | 0 | 10 | 2 309 | 2 886 |
| 1989-90 | 516 | 524 | 371 | 438 | 0 | 10 | 2 377 | 2 996 |
| 1990-91 | 420 | 531 | 369 | 441 | 0 | 10 | 2 215 | 3 050 |
| 1991-92 | 431 | 531 | 409 | 441 | 0 | 10 | 2 508 | 3 086 |
| 1992-93 | 482 | 531 | 484 | 441 | 0 | 10 | 2 839 | 3 089 |
| 1993-94 | 473 | 531 | 448 | 441 | 0 | 10 | 2 603 | 3 093 |
| 1994-95 | 370 | 534 | 417 | 441 | 0 | 10 | 2 583 | 3 105 |
| 1995-96 | 635 | 534 | 521 | 441 | 0 | 10 | 3 387 | 3 107 |
| 1995-96 | 542 | 534 | 459 | 441 | 0 | 10 | 3 153 | 3 107 |
| 1997-98 | 471 | 534 | 447 | 441 | 0 | 10 | 2 917 | 3 107 |
| 1998-99 | 681 | 534 | 533 | 441 | 0 | 10 | 3 421 | 3 107 |
| 1999-00 | 639 | 534 | 469 | 441 | 0 | 10 | 3 324 | 3 107 |
| 2000-01 | 576 | 534 | 453 | 441 | 0 | 10 | 3 193 | 3 107 |
| 2001-02 | 501 | 534 | 449 | 441 | 0 | 10 | 2 913 | 3 121 |
| 2002-03 | 512 | 534 | 448 | 441 | 0 | 10 | 3 161 | 3 121 |
| 2003-04 | 574 | 534 | 405 | 441 | 0 | 10 | 3 124 | 3 121 |
| 2004-05 | 546 | 641 | 554 | 529 | 0 | 10 | 3 368 | 3 416 |
| 2005-06 | 568 | 641 | 503 | 529 | 0 | 10 | 3 102 | 3 416 |
| 2006-07 | 583 | 641 | 534 | 529 | 0 | 10 | 3 180 | 3 416 |
| 2007-08 | 606 | 641 | 497 | 529 | 0 | 10 | 3 299 | 3 437 |
| 2008-09 | 694 | 641 | 588 | 529 | 0 | 10 | 3 477 | 3 437 |
| 2009-10 | 605 | 641 | 460 | 529 | 0 | 10 | 3 268 | 3 436 |
| 2010-11 | 677 | 641 | 587 | 529 | 0 | 10 | 3 469 | 3 436 |
| 2011-12 | 612 | 641 | 506 | 529 | 0 | 10 | 3 276 | 3 436 |

* FSU data. § Includes landings from unknown areas before 1986-87.




Figure 1: Historical landings and TACC for the seven main SCH stocks. From top to bottom: SCH1 (Auckland East), SCH2 (Central East) and SCH3 (South East coast). [Continued on next page].

SCHOOL SHARK (SCH)




Figure 1 [Continued]: Historical landings and TACC for the seven main SCH stocks. From top to bottom: SCH4 (South East Chatham Rise), SCH5 (Southland) and SCH7 (Challenger). [Continued on next page].




Figure 1 [Continued]: Historical landings and TACC for the seven main SCH stocks. Above: SCH8 (Central Egmont).

The Recreational Technical Working Group recommends that the harvest estimates from the diary surveys should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. Relative comparisons may be possible between stocks within these surveys.

Table 3: Estimated number and weight of school sharks harvested by recreational fishers relative to Fishstock and survey. Surveys were carried out in different years in the Ministry of Fisheries regions: South in 1991-92, Central in 1992-93 and North in 1993-94 (Teirney *et al.* 1997).

| Fishstock | Survey | Total | | Survey harvest (t) |
|-----------|---------|--------|--------|--------------------|
| | | Number | CV (%) | |
| SCH 1 | North | 17 000 | 24 | 10-170 |
| SCH 1 | Central | 1 000 | - | 0-10 |
| SCH 2 | Central | 13 000 | 27 | 25-45 |
| SCH 3 | South | 6 000 | 33 | 15-35 |
| SCH 5 | South | 1 000 | - | 0-10 |
| SCH 7 | Central | 9 000 | 84 | 10-35 |
| SCH 7 | South | 3 000 | - | 5-15 |
| SCH 8 | Central | 7 000 | 45 | 10-30 |

Table 4: Estimates of annual number and weight of school shark harvested by recreational fishers from national diary surveys in 1996 (Bradford 1998) and Dec1999-Nov 2000 (Boyd & Reilly 2005). The mean weights used to convert numbers to catch weight are considered the best available estimates. Estimated harvest is also presented as a range to reflect the uncertainty in the point estimates.

| Fishstock | Number caught | CV (%) | Estimated harvest range (t) | Point estimate (t) |
|-----------|---------------|--------|-----------------------------|--------------------|
| 1996 | | | | |
| SCH 1 | 23 000 | 17 | 35-55 | 46 |
| SCH 2 | 5 000 | - | - | - |
| SCH 3 | 3 000 | - | - | - |
| SCH 5 | 1 000 | - | - | - |
| SCH 7 | 8 000 | 24 | 5-25 | 16 |
| SCH 8 | 11 000 | 22 | 15-25 | 21 |
| 1999-00 | | | | |
| SCH 1 | 27 000 | 42 | 38-93 | 66 |
| SCH 2 | 7 000 | 30 | 13-24 | 18 |
| SCH 3 | 19 000 | 46 | 26-70 | 48 |
| SCH 5 | 3 000 | 66 | 2-11 | 7 |
| SCH 7 | 23 000 | 56 | 26-91 | 58 |
| SCH 8 | 3 000 | 55 | 4-13 | 8 |

1.3 Customary non-commercial fisheries

Maori fishers made extensive use of school shark in pre-European times for food, oil, and skin. There is no quantitative information on the current level of customary non-commercial take.

1.4 Illegal catch

There is no quantifiable information on the level of illegal catch. There is an unknown amount of unreported offshore trawl and pelagic longline catch of school shark, either landed (under another name, or in "mixed") or discarded.

1.5 Other sources of mortality

There is an unknown discarded bycatch of juvenile, mainly first-year, school shark taken in harbour and bay setnets. Quantitative information is not available on the level of other sources of mortality.

2. BIOLOGY

School sharks are distributed across the shelf, generally being inshore in summer and offshore in winter. They extend in smaller numbers near the seafloor down the upper continental slope, to at least 600 m. The capture of school sharks by tuna longliners shows that their distribution extends well offshore, up to 180 nm off the South Island, and 400 nm off northern New Zealand towards the Kermadec Islands. They feed predominantly on small fish and cephalopods (octopus and squid).

Growth rates have not been estimated for New Zealand fish, but in Australia and South America school sharks are slow growing and long-lived (Grant *et al.* 1979, Olsen 1984, Peres & Vooren 1991). They are difficult to age by conventional methods, but up to 45 vertebral rings can be counted. Growth is fastest for the first few years, slows appreciably between 5 and 15 years, and is negligible at older ages, particularly after 20. Results from an Australian long-term tag recovery suggest a maximum age of at least 50 years. Age-at-maturity has been estimated at 12-17 years for males and 13 to 15 years for females (Francis & Mulligan 1998). The size range of commercially caught maturing and adult school shark is 90-170 cm total length (TL), with a broad mode at 110-130 cm TL, which varies with area, season and depth.

Breeding is not annual; it has generally been assumed to be biennial, but recent work on a Brazilian stock suggests that females have a 3-year cycle (Peres & Vooren 1991). Fecundity (pup number) increases from 5-10 in small females to over 40 in the largest. Mating is believed to occur in deep water, probably in winter. Release of pups occurs during spring and early summer (November-January), apparently earlier in the north of the country than in the south. Nursery grounds include harbours, shallow bays and sheltered coasts. The pups remain in the shallow nursery grounds during their first one or two years and subsequently disperse across the shelf. The geographic location of the most important pupping and nursery grounds in New Zealand is not known.

Table 5: Estimates of biological parameters for school shark.

| Fishstock | Estimate | | Source |
|---|---------------------|------|--|
| <u>1. Weight = a (length)^b (Weight in g, length in cm fork length)</u> | | | |
| | Both sexes combined | | |
| | a | b | |
| SCH 1 | 0.0003 | 3.58 | McGregor (unpub.) |
| SCH 3 | 0.0035 | 3.08 | McGregor (unpub.) |
| SCH 5 | 0.0181 | 2.72 | McGregor (unpub.) |
| SCH 5 | 0.0068 | 2.94 | Hurst <i>et al.</i> (1990) |
| SCH 7 | 0.0061 | 2.94 | Blackwell (unpub.) |
| SCH 8 | 0.0104 | 2.84 | Blackwell (unpub.) |
| <u>2. Estimate of <i>M</i> for Australia</u> | | | |
| | 0.1 | | Grant <i>et al.</i> (1979), Olsen (1984) |

The combination of late maturity, slow growth, and low fecundity gives a low overall productivity. In Australia, *M* has been estimated as 0.1.

New Zealand tagging studies have shown that school shark may move considerable distances, including trans-Tasman migrations (for details see the 1995 Plenary Report).

Biological parameters relevant to stock assessment are shown in Table 5.

3. STOCKS AND AREAS

Information relevant to determining school shark stock structure in New Zealand was reviewed in 2009 (Smith 2009, Blackwell & Francis 2010, Francis 2010). Primarily based on the tagging evidence, there is probably a single biological stock in the New Zealand EEZ. Genetic, biological, fishery and tagging data were all considered, but the evidence for the existence of distinct biological stocks is poor. Some differences were found in CPUE trends between OMAs, but stock separation at the QMA level seems unlikely, and the CPUE differences may have resulted from processes acting below the stock level, such as localised exploitation of different sexes or different size classes of sharks. An apparent lack of juvenile school shark nursery areas in SCH 4 and SCH 5 suggests that these Fishstocks are not distinct, but are instead maintained by recruitment from other QMAs.

The most useful source of information was an opportunistic tagging programme undertaken mainly on research trawlers since 1985 (Hurst *et al.* 1999). However most tag releases were made around the South Island so little information is provided for North Island school shark. Female school shark were slightly more mobile than males, with higher proportions of the former moving to non-adjacent QMAs and to Australia. About 30% of school shark recaptures were reported from outside the release QMA within a year of release, and this was maintained in the second year after release. After 2-5 years at liberty about 60% of recaptured school sharks (both sexes) were reported from outside the release QMA. After more than 5 years at liberty, 8% of males and 19% of females were recaptured from Australia. A large proportion of tagged school sharks moved outside the QMA of release within 5 years, and a significant proportion eventually moved to Australia. These trends in apparent movement are consistent across two decades of tagging. The relative importance of various breeding grounds around New Zealand (e.g., aggregations of breeding females in Kaipara Harbour) and whether females return to the area in which they were born are unknown.

The current stock management units are a precautionary measure to spread fishing effort; amalgamation of all QMAs into one QMA for the whole EEZ could create unacceptable risks to stock sustainability.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

Fishery characterisations and CPUE analyses for SCH 1, SCH2, SCH3, SCH 5, SCH 7 and SCH 8 were undertaken in 2010 as part of the review of AMP stocks. Although SCH 1 and SCH 2 are not AMP stocks they were included by Industry to obtain a better understanding of the status of New Zealand school shark.

SCH 1

SCH 1 are primarily taken by bottom trawl while targeting tarakihi and snapper, with smaller catches when targeting trevally and red gurnard. The bottom longline SCH 1 fishery is primarily directed at school shark, with hapuku and snapper being other important targets. The setnet fishery is also primarily targeted at school shark, with some targeting of rig, trevally, gurnard and snapper.

The previously accepted indices for SCH 1 were based on bottom longline snapper (1E) and a bottom trawl mixed (1W) catches. The 2010 assessment explored a wide range of alternative fishery definitions and the AMP FAWG accepted indices based on SN and BLL catches on both the east and west coasts. These indices were based on Generalized Linear Models of positive catches with log normal error distribution. Models of bottom trawl catch were not explored.

SCHOOL SHARK (SCH)

Standardised CPUE abundance indices for SCH 1 show different trends west and east of North Cape (Fig. 2).

SCH 1 W

Discounting the last two years of the analysis (2007/08 and 2008/09) for setnet (which are poorly estimated), the SN and BLL indices for SCH 1W are flat, indicating no change in abundance over the past 20 years (Fig. 2). Analysis of the spatial distribution of catches revealed that the BLL catches were concentrated around North Cape and the SN catches were mostly made in the North Taranaki Bight; near the SCH 8 boundary line. The SN index shows very high uncertainty over the last two years of the analysis (2007/08 and 2008/09), potentially being affected by recent setnet closures on the west coast. There are now only two vessels in this west coast SN fishery and the index may become unreliable in future.

SCH 1 E

Since 1998-99 the SCH 1E index shows an increasing trend to above the long-term average, peaking in 2003-04, and then dropping to just above the average by 2005-06 and remained at about that level to 2008/09 (Fig. 2).




Figure 2: [left panel] comparison of the two SCH 1W standardised series: SCH 1W_SN(SHK+GUR) and SCH 1W_BLL(44inc); [right panel] comparison of the two SCH 1E standardised series: SCH 1E_SN(3in4+SNA+TRE) and SCH 1E_BLL(MIX). (Each series is scaled so that the geometric mean = 1) Starr & Kendrick (2010a).

SCH 2

SCH 2 are caught primarily in the bottom trawl fishery (46%) targeting tarakihi, hoki, gemfish and gurnard; and the bottom longline fishery (30%) targeting school shark, ling, hapuku/bass and bluenose. Sixteen per cent of the catch is taken in setnet targeting school shark, blue warehou and blue moki.

The 2010 analyses used setnet and bottom longline (no bottom trawl index was attempted), based on a broader range of target fisheries than previously. The previous assessment used tarakihi bottom trawl index.

Two indices were considered for SCH 2 in 2010: one based on setnet catches with a range of target (SN[MIX]) and the other based on bottom longline catches, also with a range of targets. These two indices present conflicting trends, the setnet index generally increasing over the series and the bottom longline index decreasing steadily (Fig. 3). The AMP FAWG noted particular concerns with the bluenose targeted bottom longline index, related to suggestions of a steady shift towards mid-water targeting of bluenose. There is a substantial correspondence between the standardised setnet index for SCH 2 with setnet and bottom longline indices for SCH 1E, which together indicate a slow but steady increase in CPUE to 2005-06, levelling off since then.




Figure 3: Comparison of the lognormal indices from the two fisheries operating in SCH 2: a) SN[MIX]: mixed target species; b) BLL[MIX]: mixed target species Starr & Kendrick (2010b).

SCH 3

SCH 3 is predominantly caught in the setnet fishery (56%) targeting school shark and rig, with some targeting of spiny dogfish and tarakihi; and in the bottom trawl fishery (36%) targeting red cod, with some targeting of flatfish, barracouta and tarakihi. Mixed targeted bottom longline takes 7% of the catch.

The mixed shark target SN(SHK) standardised CPUE is the accepted index of abundance for SCH 3. The 2010 CPUE analysis is an update of the shark-targeted setnet CPUE analysis conducted in 2003 and 2007, with no extension to other target species or other model changes. This index shows a sharp decline of almost 60% from a peak in 1989-90 to its lowest point over the 20 year series in 1992-93 (Fig. 4). Thereafter the index shows a steady and continual increase through to 2003-04 / 2004-05 to a level about 10% above the long-term average and about 40% above the lowest level, fluctuating around this level thereafter.




Figure 4: Lognormal SCH 3_SN(SHK) standardised indices with core vessel criteria of at least 10 trips in a minimum of 6 years) (series is scaled so that the geometric mean = 1). From Starr *et al.* (2010a).

SCHOOL SHARK (SCH)

Biomass estimates

Biomass in the core strata (30–400 m) for the east coast South Island trawl survey is variable, but generally higher in recent years compared with the 1990s (Figure 5). Coefficients of variation are variable ranging from 18 to 42% (mean 25%), but overall are low to medium. The additional biomass captured in the 10–30 m depth range accounted for only 2.5% and 6% of the biomass in the core plus shallow strata (10–400 m) for 2007 and 2012, respectively.

Length frequency distributions

School shark are most common in 30–100 m with a tendency for the smallest cohorts to be in the shallower depth ranges for the east coast South Island trawl survey (Figure 6). The three modes at 35, 50, and 60 cm are all pre-recruited school shark and correspond to ages of 0+, 1+, and 2+. The survey appears to be monitoring pre-recruited cohorts 0+, 1+, 2+ (and possibly a few more older cohorts) reasonably well, but not the recruited school shark size distribution. Plots of time series length frequency distributions are spiky because of the low numbers caught, but the size range is reasonably consistent among surveys. The addition of the 10–30 m depth range has changed the shape of the length frequency distribution slightly.




Figure 5: School shark total biomass and 95% confidence intervals for the all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007 and 2012.

Table 6: Relative biomass indices (t) and coefficients of variation (CV) for school shark for the east coast South Island (ECSI) – winter, survey area*. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16 and 17). – , not measured; NA, not applicable.

| Region | Fishstock | Year | Trip number | Total Biomass estimate | CV (%) | Total Biomass estimate | CV (%) |
|---------------|-----------|------|-------------|------------------------|--------|------------------------|--------|
| ECSI (winter) | SCH | | | 30-400m | | 10-400m | |
| | | 1991 | KAH9105 | 100 | 30 | - | - |
| | | 1992 | KAH9205 | 104 | 21 | - | - |
| | | 1993 | KAH9306 | 369 | 42 | - | - |
| | | 1994 | KAH9406 | 155 | 36 | - | - |
| | | 1996 | KAH9608 | 202 | 18 | - | - |
| | | 2007 | KAH0705 | 538 | 22 | 522 | 21 |
| | | 2008 | KAH0806 | 411 | 20 | - | - |
| | | 2009 | KAH0905 | 254 | 18 | - | - |
| | | 2012 | KAH1207 | 292 | 20 | 310 | 19 |




Figure 6: Scaled length frequency distributions for school shark in core strata (30–400 m) for all nine ECSI winter surveys. The length distribution is also shown in the 10–30 m depth strata for the 2007 and 2012 surveys overlaid in red for species with many length classes, otherwise in light grey (not stacked). Population estimates are for the core strata only. n, number of fish measured; no., population number; c.v., coefficient of variation.

SCHOOL SHARK (SCH)

SCH 5

SCH 5 is almost entirely caught in the school shark targeted setnet fishery (83%), with some minor targeting of rig. Seven percent is taken by bottom trawl primarily targeting stargazer and squid, and 5% by bottom longline primarily targeting hapuku/bass and ling.

The targeted SN(SCH) standardised CPUE index is the accepted indicator of SCH 5 abundance. The 2011 CPUE analysis is an update of previous analyses conducted in 2003, 2007 and 2010, with no substantial changes to the fishery definitions or standardisation models. The index fluctuated around long-term average levels through to 2005-06 (Fig. 7). Thereafter the index declines to slightly below average levels over 2006-07/2007-08, and then steeply to about half average levels in 2008-09. The index is considered to be less reliably estimated in the final year, due to changes in fleet size and structure.




Figure 7: Lognormal SCH 5_SN(2011) standardised index and the index from the 2010 assessment SCH5_SN(2010). From Starr and Kendrick (2011).

There is close correspondence in the declining indices for SCH 5 and SCH 7, except in the final year. Both indices monitor mature fish caught around Southland and the WCSI, raising some concern about the declines in both these areas.

SCH 7

SCH 7 are caught about one-third each by setnet targeting school shark, rig and spiny dogfish; bottom longline targeting school shark, hapuku/bass and ling; and bottom trawl targeting barracouta, tarakihi, flatfish, hoki, red cod and others.

The mixed shark target SHK7_SN(2011) standardised CPUE index is the accepted indicator of SCH 7 abundance. The 2011 CPUE analysis updates previous analyses conducted in 2003, 2007 and 2010, with no substantial changes to the fishery definitions or standardisation models. The index remained stable around long-term average levels over 1989-90 to 1995-96 and then increased to a peak about 50% above average levels in 1999-00, then declined steadily to its lowest value over the 20 year period by 2007-08 (Figure 8). The index increased in 2008-09 and then decreased in the final year to below the long-term mean.

There is close correspondence in the declining indices for SCH 5 and SCH 7, except in the final two years where SCH 5 and SCH 7 vary inversely. Both indices monitor mature fish caught around Southland and the WCSI, raising some concern about the declines in both these areas.




Figure 8: Lognormal indices from the setnet target shark CPUE series for SCH 7 SCH7_SN(2011) and the index from the 2010 assessment SCH7_SN(2010). From Starr & Kendrick (2011).

SCH 8

SCH 8 are caught mainly (66%) by setnet targeting school shark and rig; and by bottom longline (22%) targeting school shark and hapuku/bass. Ten percent is caught by bottom trawl targeting gurnard, tarakihi and trevally.

The mixed shark target SCH8_SN(2011) standardised CPUE index is the accepted indicator of SCH 8 abundance. The 2011 CPUE analysis is an update of previous analyses conducted in 2003, 2007 and 2010, with no substantial changes to the fishery definitions or standardisation models. The index remains flat at the long-term average, apart from a drop to lower levels over 1997-98 to 2000-01 (Figure 9). The Working Group concluded that the SCH 8 index showed no change in abundance over the series. There was an inverse relationship between the SCH 7 and SCH 8 indices over this period, suggesting a possible shift in stock distribution between these areas.




Figure 9: Lognormal indices from the setnet target shark CPUE series for SCH 8 SCH8_SN(2011) and the index from the 2010 assessment SCH8_SN(2010). From Starr and Kendrick (2011).

SCHOOL SHARK (SCH)

SCH overview

SCH are mainly caught in setnet fisheries targeting sharks (school shark, rig, elephantfish and spiny dogfish); in bottom trawl fisheries targeting red cod, tarakihi, gurnard and snapper and others; and in bottom longline fisheries targeting school shark, hapuku/bass and ling.

In SCH 3, 5, 7 and 8, CPUE indices have been conducted using the same, or similar, models since entry into AMPs. In some areas, additional target species have been added to fishery definitions, particularly for bottom longline indices. New analyses were developed for SCH 1 and 2. Bottom trawl indices previously produced for SCH 1 and 2 were not updated in 2010.

There is a close similarity in trends in the indices for 1E, 2 and 3; SCH 5 and SCH 7; and SCH 8 and 1W. The indices show higher recent CPUE for SCH 1E, 2 and 3; stable CPUE for SCH 1W and 8; and declining CPUE for SCH 5 and 7. The Working Group noted that SCH 5 and 7 have accounted for 41% of the SCH catch over the past 20 years, and are the areas in which the highest proportion of mature fish are caught. SCH 1E, 2 and 3, have accounted for 26% of the SCH catch over the past 20 years. Areas 1W and 8 have accounted for 30% of the catch.

Recent setnet closures have potentially compromised the continuity of setnet indices for SCH 1W, 3, 5 and 7.

4.2 Yield estimates and projections

The estimates of *MCY* are no longer considered valid.

Current biomass cannot be estimated, so *CAY* cannot be determined.

4.3 Other factors

In Australia, recruitment overfishing has occurred to such an extent that the stock is considered seriously threatened and a series of conservative management measures (TAC reductions) have been progressively imposed between 1996 and 2007 (Wilson *et al.* 2008). The Australian modelling work indicates that the stock is overfished. Wilson *et al.* (2008) noted that the stock had been in an overfished state and overfishing was occurring from 1992 to 2004. While the stock was still listed as overfished since then, they are uncertain as to whether overfishing is still occurring.

The most important conclusion from this for New Zealand is that fishing pressure on large mature females should be minimised to maintain the productivity of this species.

5. STATUS OF THE STOCKS

Stock Structure Assumptions

SCH are known from tagging studies to be highly mobile, moving between the North and South Islands, and as far as Australia. From the tagging evidence, there is probably a single biological SCH stock in the New Zealand EEZ. However, differences in average modal length and CPUE trends between FMAs indicate that movement between areas may be viscous, and that components of the stock may aggregate in different areas. Larger females predominate in catches around Southland and west coast of the South Island. Therefore, the current stock management units are a precautionary measure to spread fishing effort and mortality across components of the stock.

• SCH 1

| Stock Status | |
|--------------------------------|---|
| Year of Most Recent Assessment | 2010 (Fishery characterisation and CPUE standardisation) |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |

| | |
|--|---|
| Status in relation to Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |
| Historical Stock Status Trajectory and Current Status | |
| | |
| <p>Left: Comparison of the two SCH 1W standardised series: SCH 1W_SN(SHK+GUR) and SCH 1W_BLL(44inc); Right: The SN (SHK+SNA+TRE) index for SCH 1E.</p> | |

| | |
|----------------------------------|--|
| Fishery and Stock Trends | |
| Recent Trend in Biomass or Proxy | <p>Standardised CPUE abundance indices for SCH 1 show different trends west and east of North Cape. Discounting the last two years for setnet (which are poorly estimated), the SN and BLL indices for SCH 1W are flat, indicating no change in abundance over the past 20 years.</p> <p>The index for SCH 1E shows higher than long-term average abundance since 1999-00. From 1999-00 the SCH 1E index shows an increasing trend to above the long-term average, peaking in 2003-04, and then dropping to just above the average by 2006-07.</p> |

| | |
|--|--|
| Recent Trend in Fishing Mortality or Proxy | Overfishing is Unlikely (< 40%) to be occurring. |
|--|--|

| | |
|---|---|
| Projections and Prognosis | |
| Stock Projections or Prognosis | <p>SCH 1E: Stock size is Likely (> 60%) to remain near current levels or increase under current catches and TACCs.</p> <p>SCH 1W: Stock size is Likely (> 60%) to remain near current levels under current catches and TACCs.</p> |
| Probability of Current Catch or TACC causing decline below Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |

| | |
|--|--|
| Assessment Methodology | |
| Assessment Type | Level 2 - Standardised CPUE abundance index |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 1 abundance |
| Main data inputs | - Catch and effort data derived from the Ministry of Fisheries catch returns |
| Period of Assessment | Latest assessment: 2010 Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The previously accepted indices were based on bottom longline snapper (1E) and a bottom trawl mixed (1W). This assessment explored a wide range of alternative fishery definitions. Four credible indices were selected: setnet (SN) and bottom longline (BLL) on both the east and west coasts. |
| Major Sources of Uncertainty | - Setnet closures have jeopardised the continuity of the west coast setnet index in recent years. The BLL(W) index is considered to index the top of the North Island and lacks data. |

SCHOOL SHARK (SCH)

Qualifying Comments

Recent setnet closures designed to protect Maui's dolphin have affected setnet fisheries which take school shark off the west coast of the North Island. These closures have resulted in changes in fleet deployment and jeopardised the setnet indices.

Fishery Interactions


SCH 1 are primarily taken by bottom trawl while targeting tarakihi and snapper, with smaller catches when targeting trevally and red gurnard. The bottom longline SCH 1 fishery is primarily directed at school shark, with hapuku and snapper being other important targets. The setnet fishery is also primarily targeted at school shark, with some targeting of rig, trevally, gurnard and snapper. The bottom pair trawl fishery is almost entirely directed at snapper and trevally, with tarakihi becoming more important in recent years. In the setnet fisheries there is a risk of incidental capture of seabirds, Maui's dolphins on the west coast, other dolphins and New Zealand fur seals.

• SCH 2

Stock Status

| | |
|--------------------------------|---|
| Year of Most Recent Assessment | 2010 (Fishery characterisation and CPUE standardisation) |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |

Historical Stock Status Trajectory and Current Status



The lognormal index for SCH 2: SN[MIX]: mixed target species (scaled so that the geometric mean = 1).

Fishery and Stock Trends

| | |
|--|--|
| Recent Trend in Biomass or Proxy | The CPUE index generally increases over the series. There is a substantial correspondence between the standardised SN index for SCH 2 with SN and BLL indices for SCH 1E, which together indicate a slow but steady increase in CPUE to 2005-06, levelling off since then. |
| Recent Trend in Fishing Mortality or Proxy | Overfishing is Unlikely (< 40%) to be occurring. |

Projections and Prognosis

| | |
|--------------------------------|---|
| Stock Projections or Prognosis | Correspondence between SN indices for SCH 1E, SCH 2 and SCH 3 indicates that. SCH 2 stock size is Likely to remain near current levels or increase under current catches and TACCs. |
|--------------------------------|---|

| | |
|---|---|
| Probability of Current Catch or TACC causing decline below Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |
|---|---|

| Assessment Methodology | |
|--|--|
| Assessment Type | Level 2 - Standardised CPUE abundance index |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 2 abundance |
| Main data inputs | - Catch and effort data derived from the Ministry of Fisheries compulsory catch reporting |
| Period of Assessment | Latest assessment: 2010 Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The previous assessment used tarakihi bottom trawl index. The 2010 analyses used setnet and bottom longline (no bottom trawl index was attempted), based on a broader range of target fisheries than previously. |
| Major Sources of Uncertainty | - |


| Qualifying Comments | |
|--|--|
| There have been no regulation changes affecting SCH 2 in recent years. | |

| Fishery Interactions | |
|---|--|
| SCH 2 are caught primarily in the bottom trawl fishery (46%) targeting tarakihi, hoki, gemfish and gurnard; and the bottom longline fishery (30%) targeting school shark, ling, hapuku/bass and bluenose. 16% of the catch is taken in setnet targeting school shark, blue warehou and blue moki. | |

- **SCH 3**

| Stock Status | |
|--------------------------------|---|
| Year of Most Recent Assessment | 2010 (Fishery characterisation and CPUE standardisation) |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |

Historical Stock Status Trajectory and Current Status



Lognormal SCH 3_SN(SHK) standardised indices with core vessel criteria of at least 10 trips in a minimum of 6 years) (series is scaled so that the geometric mean = 1).

Fishery and Stock Trends

| | |
|--|--|
| Recent Trend in Biomass or Proxy | The mixed shark target SN(SHK) standardised CPUE is the accepted index of abundance. This index shows a sharp decline of almost 60% from a peak in 1989-90 to its lowest point over the 20 year series in 1992-93. Thereafter the index shows a steady and continual increase through to 2003-04 to 2004-05 to a level about 10% above the long-term average and about 40% above the lowest level, fluctuating around this level thereafter. |
| Recent Trend in Fishing Mortality or Proxy | Overfishing is Unlikely (< 40%) to be occurring. |

Projections and Prognosis

| | |
|---|---|
| Stock Projections or Prognosis | Quantitative stock projections are unavailable. The long period of increase in the SN(SHK) index for SCH 3 since 1992-93, over a period when catches have increased steadily from about 200t to an average of 366t over the recent five years, indicates that stock size is Likely to remain near current levels or increase under current catches. |
| Probability of Current Catch or TACC causing decline below Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |

Assessment Methodology

| | |
|--|---|
| Assessment Type | Level 2: Standardised CPUE abundance index, and a review of length data. |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 3 abundance |
| Main data inputs | - Catch and effort data derived from the Ministry of Fisheries reporting requirements - Length frequency data summarised from logbooks compiled under the industry Adaptive Management Programme |
| Period of Assessment | Latest assessment: 2010 Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The 2010 CPUE analysis is an update of the shark-targeted setnet CPUE analysis conducted in 2003 and 2007, with no |

| | |
|------------------------------|--|
| | extension to other target species or other model changes. |
| Major Sources of Uncertainty | - Recent setnet closures have affected fleet distribution patterns, potentially jeopardising setnet indices in this area. These changes may have contributed to the strong fluctuations in the SCH 3 SN indices in recent years. |

Qualifying Comments

Like other setnet abundance indices, the SCH 3 setnet indices have been affected, and possibly compromised, by setnet closures.

Fishery Interactions

SCH 3 is predominantly caught in the setnet fishery (56%) targeting school shark and rig, with some targeting of spiny dogfish and tarakihi; and in the bottom trawl fishery (36%) targeting red cod, with some targeting of flatfish, barracouta and tarakihi. Mixed targeted bottom longline takes 7% of the catch. In the setnet fisheries there is a risk of incidental capture of seabirds, Hector's dolphins, other dolphins and New Zealand fur seals. There is a risk of incidental capture of sea lions from Otago Peninsula south.


- **SCH 4**

The status of SCH 4 relative to B_{MSY} is unknown.

- **SCH 5**

| | |
|--------------------------------|---|
| Stock Status | |
| Year of Most Recent Assessment | 2011 (Fishery characterisation and CPUE standardisation) |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) to be below |

Historical Stock Status Trajectory and Current Status



Lognormal SCH 5_SCH5_SN(2011) standardised index.

SCHOOL SHARK (SCH)

| Fishery and Stock Trends | |
|--|---|
| Recent Trend in Biomass or Proxy | The targeted SCH5_SN(2011) CPUE index fluctuated around long-term average levels through to 2005-06. Thereafter the index declined to slightly below average levels over 2006-07 / 2007-08, and then steeply to about half average levels in 2008-09. The value of the index in the most recent year has increased but is still below the long-term mean. |
| Recent Trend in Fishing Mortality or Proxy | Overfishing is About as Likely as Not (40-60%) to be occurring. |

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | <p>The SCH5_SN(2011) abundance index has declined steadily since 2005-06, reaching the lowest level over the 20 year period of the index in 2008-09, at about half of long-term average levels.</p> <p>This gives rise to concern that current catches, and the current TACC, may not be sustainable. While the most recent data point is above that of 2010, it is still below the long-term mean. The working group therefore concluded that the SCH 5 stock is Likely to decline under current catches and TACCs.</p> |
| Probability of Current Catch or TACC causing decline below Limits | <p>Soft Limit: Unknown</p> <p>Hard Limit: Unknown</p> |

| Assessment Methodology | |
|--|--|
| Assessment Type | Level 2 - Standardised CPUE abundance index, and a review of length data |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 5 abundance |
| Main data inputs | <ul style="list-style-type: none"> - Catch and effort data derived from the Ministry of Fisheries catch reporting - Length frequency data summarised from logbooks compiled under the industry Adaptive Management Programme |
| Period of Assessment | Latest assessment: 2011 Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The 2011 CPUE analysis is an update of previous analyses conducted in 2003, 2007 and 2010, with no substantial changes to the fishery definitions or standardisation models. |
| Major Sources of Uncertainty | - Recent setnet closures have affected fleet distribution patterns, potentially jeopardising setnet indices in this area. |

| Qualifying Comments |
|---|
| Concerns regarding the status of this stock are prompted by the decline in CPUE from the early 2000s. There is close correspondence in the indices for SCH 5 and SCH 7. Both indices monitor mature fish caught around Southland and the WCSI, raising some concern for both these areas. |

| Fishery Interactions |
|---|
| SCH 5 is almost entirely caught in the school shark targeted setnet fishery (83%), with some minor targeting of rig. Seven percent is taken by bottom trawl primarily targeting stargazer and squid, and 5% by bottom longline primarily targeting hapuku/bass and ling. In the setnet fisheries there is a risk of incidental capture of seabirds, white pointer sharks, Hector's dolphins, other dolphins and New Zealand fur seals. There is a risk of incidental capture of sea lions from Otago Peninsula south. |

- SCH 7

| Stock Status | |
|---|---|
| Year of Most Recent Assessment | 2011 |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) to be below |
| Historical Stock Status Trajectory and Current Status | |
| <p style="text-align: center;">SCH 7</p> <p style="text-align: center;">Relative CPUE Indices</p> <p style="text-align: center;">Fishing Year</p> <p style="text-align: center;">— SCH7_SN(2011) - - - SCH7_SN(2010)</p> | |
| Lognormal indices from the setnet target shark CPUE series for SCH 7. | |

| Fishery and Stock Trends | |
|---|--|
| Recent Trend in Biomass or Proxy | The mixed shark target SHK7_SN(2011) standardised CPUE index remained stable around long-term average levels over 1989-90 to 1995-96 and then increased to a peak about 50% above average levels in 1999-00, then declined steadily to its lowest value over the 20 year period by 2007-08. There was a sharp increase in the final year; however, the working group considered the last data point to be less reliably estimated. |
| Recent Trend in Fishing Mortality or Proxy | Overfishing is About as Likely as Not (40-60%) to be occurring. |
| Projections and Prognosis | |
| Stock Projections or Prognosis | The SHK7_SN(2011) abundance index declined steadily from 1999-00 to its lowest level over the 20 year period of the index in 2007-08. The Working Group concluded that the SCH 7 Fishstock stock size is Likely to decline under current catches and TACCs. |
| Probability of Current Catch or TACC causing decline below Limits | Soft Limit: Unknown Hard Limit: Unknown |

| Assessment Methodology | |
|-------------------------------|--|
| Assessment Type | Level 2 - Standardised CPUE abundance index, and review of length frequency data |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 7 abundance |

SCHOOL SHARK (SCH)

| | | |
|---|--|-----------------------|
| Main data inputs | <ul style="list-style-type: none"> - Catch and effort data derived from the Ministry of Fisheries catch reporting - Length frequency data summarised from logbooks compiled under the industry Adaptive Management Programme | |
| Period of Assessment | Latest assessment: 2011 | Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The 2011 CPUE analysis is an update of previous analyses conducted in 2003, 2007 and 2010, with no substantial changes to the fishery definitions or standardisation models. | |
| Major Sources of Uncertainty | <ul style="list-style-type: none"> - The fleet distribution has changed in the last 2-3 years. - Furthermore, setnet closures have potentially compromised the setnet indices in the last two years. | |
| Qualifying Comments | | |
| Concerns regarding the status of this stock are prompted by the decline in CPUE from the early 2000s. There is close correspondence in the indices for SCH 5 and SCH 7. Both indices monitor mature fish caught around Southland and the WCSI, raising some concern for both these areas. | | |

Fishery Interactions

SCH 7 are caught about one-third each by setnet targeting school shark, rig and spiny dogfish; bottom longline targeting school shark, hapuku/bass and ling; and bottom trawl targeting barracuda, tarakihi, flatfish, hoki, red cod and others. In the setnet fisheries there is a risk of incidental capture of seabirds, dolphins and New Zealand fur seals.

• **SCH 8**

| | |
|--|---|
| Stock Status | |
| Year of Most Recent Assessment | 2011 |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft limit: Unknown Hard Limit: Unlikely (< 40%) to be below |
| Historical Stock Status Trajectory and Current Status | |
| <p style="text-align: center;"> SCH8_SN(2011) SCH8_SN(2010) </p> | |
| <p>Lognormal indices from the setnet target shark CPUE series for SCH 8</p> | |

| Fishery and Stock Trends | |
|--|--|
| Recent Trend in Biomass or Proxy | The SCH8SN(2011) index remains flat at the long-term average, apart from a drop to lower levels over 1997-98 to 2000-01. The Working Group concluded that the SCH 8 index showed no change in abundance over the series. There is an inverse relationship between the SCH 7 and SCH 8 indices over this period, suggesting a possible shift in stock distribution between these areas. |
| Recent Trend in Fishing Mortality or Proxy | Overfishing is Unlikely (< 40%) to be occurring. |

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | The lack of any trend in the SCH 8 CPUE index indicates the stock size is Likely to remain near current levels under current catches and TACCs. |
| Probability of Current Catch or TACC causing decline below Limits | Soft Limit: Unknown Hard Limit: Unlikely (< 40%) |
| Assessment Methodology | |
| Assessment Type | Level 2 - Standardised CPUE abundance index, and review of length frequency data |
| Assessment Method | Evaluation of agreed standardised CPUE indices thought to index SCH 8 abundance |
| Main data inputs | - Catch and effort data derived from the Ministry of Fisheries catch reporting - Length frequency data summarised from logbooks compiled under the industry Adaptive Management Programme |
| Period of Assessment | Latest assessment: 2011 Next assessment: 2013 |
| Changes to Model Structure and Assumptions | The 2010 CPUE analysis are updates of previous analyses conducted in 2003 and 2007, with no substantial changes to the fishery definitions or standardisation models. |
| Major Sources of Uncertainty | - |

| Qualifying Comments |
|----------------------------|
| - |

| Fishery Interactions |
|---|
| SCH 8 are caught mainly (66%) by setnet targeting school shark and rig; and by bottom longline (22%) targeting school shark and hapuku/bass. Ten percent is caught by bottom trawl targeting gurnard, tarakihi and trevally. In the setnet fisheries there is a risk of incidental capture of seabirds, Maui's dolphins other dolphins and New Zealand fur seals. |

- **Combined SCH Stocks**


School shark are believed to be a single biological stock around the North and South Islands. It may therefore be appropriate for management responses to be consistent across areas broader than single QMAs.

| Stock Status | |
|--------------------------------|---|
| Year of Most Recent Assessment | 2010; 2011 for SCH 5, 7 and 8 |
| Reference Points | Target: Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 |
| Status in relation to Target | Unknown |
| Status in relation to Limits | Soft limit: Unknown Hard Limit: Unlikely (< 40%) |


SCHOOL SHARK (SCH)

Historical Stock Status Trajectory and Current Status

SCH 1E, 2 and 3




Comparison of three lognormal indices from a) the SCH 1E setnet series: [SCH 1E_SN(3in4+SNA=TRE)]; b) a setnet series from SCH 2: [SCH2_SN(MIX)]; c) a setnet series from SCH 3: [SCH3_SN(SHK)]



Comparison of the selected SCH 5 [SCH5_SN(SCH)] series with the equivalent series selected for the SCH 7 [SCH7_SN(SHK)] CPUE evaluation. Confidence intervals only shown for one series.

SCH 8 and 1W



Comparison of the selected SCH 8 [SCH8_SN(SHK)] series with the equivalent series selected for the SCH1W [SCH1W_SN(SHK+GUR)] CPUE evaluation.

Fishery and Stock Trends

| | |
|--|--|
| Recent Trend in Biomass or Proxy | There is a close similarity in trends in the indices for 1E, 2 and 3; SCH 5 and SCH 7; and SCH 8 and 1W. The indices show an increase or higher recent CPUE for SCH 1E, 2 and 3; and stable CPUE for SCH 1W and 8; SCH 5 and 7 show CPUE declining. The Working Group noted that SCH 5 and 7 have accounted for 41% of the SCH catch over the past 20 years, and are the areas in which the largest females are caught. SCH 1E, 2 and 3, have accounted for 26% of the SCH catch over the past 20 years. Areas 1W and 8 have accounted for 30% of the catch. |
| Recent Trend in Fishing Mortality or Proxy | Varies among FMAs |

Projections and Prognosis

| | |
|---|---|
| Stock Projections or Prognosis | <ul style="list-style-type: none"> •SCH 1E, 2, 3: Stock size is Likely to remain near current levels or increase under current catches and TACCs. •SCH 1W, 8: Stock size is Likely to remain near current levels under current catches and TACCs. •SCH 5&7: Stock size is Likely to decline under current catches and TACCs. |
| Probability of Current Catch or TACC causing decline below Limits | Combined stocks: Soft Limit: Unknown Hard Limit: Varies among FMAs |

Assessment Methodology

| | |
|-------------------|--|
| Assessment Type | Level 2 - Standardised CPUE abundance index, and length frequency analysis |
| Assessment Method | Evaluation of a range of agreed standardised CPUE indices |

SCHOOL SHARK (SCH)

| | | |
|--|---|--|
| | thought to index abundance of the SCH fishstocks in each FMA Comparison of length frequencies from SCH 3, 5, 7 and 8 | |
| Main data inputs | <ul style="list-style-type: none"> - Catch and effort data derived from the Ministry of Fisheries catch reporting - Length frequency data summarised from logbooks compiled under the industry Adaptive Management Programme | |
| Period of Assessment | Latest assessment: 2010 | Next assessment: 2011 (SCH 5 and 7) and 2013 (SCH 1, 2, 3 and 8) |
| Changes to Model Structure and Assumptions | In SCH 3, 5, 7 and 8, CPUE indices have been conducted using the same, or similar, models since entry into AMPs. In some areas, additional target species have been added to fishery definitions, particularly for bottom longline indices. New analyses were developed for SCH 1 and 2. Bottom trawl indices previously produced for SCH 1 and 2 were not updated in 2010. | |
| Major Sources of Uncertainty | - Recent setnet closures have potentially compromised the continuity of setnet indices for SCH 1W, 3, 5 and 7. | |

Qualifying Comments

See individual Fishstock Status of Stocks summaries.

Fishery Interactions

SCH are predominantly caught in setnet fisheries targeting sharks (school shark, rig, elephantfish and spiny dogfish); in bottom trawl fisheries targeting red cod, tarakihi, gurnard and snapper and others; and in bottom longline fisheries targeting school shark, hapuku/bass and ling. On the east coast on the North Island in the setnet fisheries there is a risk of incidental capture of seabirds, dolphins and New Zealand fur seals. On the east coast of the South Island in the setnet fisheries there is a risk of incidental capture of seabirds, Hector's dolphins, other dolphins and New Zealand fur seals. There is a risk of incidental capture of sea lions from Otago Peninsula south. On the west coast of the North Island in the setnet fisheries there is a risk of incidental capture of seabirds, Maui's dolphins, other dolphins and New Zealand fur seals.

Yield estimates, reported landings and TACCs for the 2011-12 fishing year are summarised in Table 7.

Table 7: Summary of yield estimates (t), TACCs (t) and reported landings (t) of school shark for the most recent fishing year.

| Fishstock | QMA | MCY Estimates | 2011-12 Actual TACC | 2011-12 Reported Landings |
|-----------------------------------|-------|---------------|---------------------|---------------------------|
| SCH 1 Auckland (East) (West) | 1 & 9 | | 689 | 689 |
| SCH 2 Central (East) | 2 | | 199 | 188 |
| SCH 3 South-east (Coast) | 3 | | 387 | 351 |
| SCH 4 South-east (Chatham) | 4 | | 239 | 201 |
| SCH 5 Southland and Sub-Antarctic | 5 & 6 | | 743 | 729 |
| SCH 7 Challenger | 7 | | 641 | 612 |
| SCH 8 Central (West) | 8 | | 529 | 506 |
| SCH 10 Kermadec | 10 | | 10 | 0 |
| Total | | | 3 436 | 3 276 |

6. FOR FURTHER INFORMATION

- Anon 1990. Management meets industry. Papers from the Southern Shark Fishery seminars held in Victor Harbour, Phillip Island and Hobart, October 1989.
- Ayers D., Paul L.J., Sanders B.M. 2004. Estimation of catch per unit effort analyses for school shark (*Galeorhinus galeus*) from bycatch and target fisheries in New Zealand, 1989-90 to 2001-02. New Zealand Fisheries Assessment Report 2006/26. 121p.
- Blackwell R.G., Francis M.P. in press. Review of life-history and fishery characteristics of New Zealand rig and school shark. *New Zealand fisheries assessment report No. 2009/xx*. 38 p.
- Boyd R.O., Reilly J.L. 2005. 1999-2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report.

- Bradford E. 1998. Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16. 27p.
- Bradford E. 2001. Standardised catch rate indices for New Zealand school shark, *Galeorhinus galeus*, in New Zealand, 1989-90 to 1998-99. New Zealand Fisheries Assessment Report 2001/33. 75p.
- Campbell D., Battaglene T., Pascoe S. 1991. Management options for the Southern Shark Fishery - an economic analysis. Australian Bureau of Agricultural and Resource Economics Discussion Paper 91.12. 43p.
- Coutin P., Bruce B., Paul L. 1992. New Zealand school sharks cross the Tasman Sea. Australian Fisheries 51(3): 24-25.
- Francis M.P. 1998. New Zealand shark fisheries: development, size and management. Marine and Freshwater Research 49: 579-591.
- Francis M.P., Mulligan K.P. 1998. Age and growth of New Zealand school shark, *Galeorhinus galeus*. New Zealand Journal of Marine and Freshwater Research 32(3): 427-440.
- Francis M.P. in press. Movement of tagged rig and school shark among QMAs, and implications for stock management boundaries. *New Zealand fisheries assessment report No. 2009/xx*. 22 p.
- Grant C.J., Sandland R.L., Olsen A.M. 1979. Estimation of growth, mortality and yield per recruit of the Australian school shark, *Galeorhinus australis* (Macleay), from tag recoveries. Australian Journal of Marine and Freshwater Research 30(5): 625-637.
- Hurst R.J., Bagley N.W., McGregor G.A., Francis M.P. 1999. Movements of the New Zealand school shark, *Galeorhinus galeus*, from tag returns. New Zealand Journal of Marine and Freshwater Research 33(1): 29-48.
- Hurst R.J., Bagley N.W., Uozumi Y. 1990. New Zealand-Japan trawl survey of shelf and upper slope species off southern New Zealand, June 1986. N.Z. Fisheries Technical Report No. 18. 50p.
- Hurst R.J., Bagley N.W., McGregor G.A., Francis M.P. (1999). Movements of the New Zealand school shark, *Galeorhinus galeus*, from tag returns. *New Zealand journal of marine and freshwater research* 33: 29-48.
- Livingston M.E., Uozumi Y., Berben P.H. 1991. Abundance, distribution, and spawning condition of hoki and other mid-slope fish on the Chatham Rise, July 1986. New Zealand Fisheries Technical Report No. 25. 47p.
- Lydon G.J., Middleton D.A.J., Starr P.J. 2006. Performance of the SCH 7 and SCH 8 Logbook Programmes. AMP-WG-06-06. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Olsen A.M. 1984. Synopsis of biological data on the school shark, *Galeorhinus australis* (Macleay 1881). FAO Fisheries Synopsis No. 139. 42p.
- Paul L.J. 1988. School shark. New Zealand Fisheries Assessment Research Document 1988/27. 32p.
- Paul L.J. 1991. Overseas travel report: "Sharks Down Under" conference, Taronga Zoo, Sydney, February 1991. MAF Fisheries Greta Point Internal Report No. 176. 137 p. (Draft report held in MAF Fisheries Greta Point library, Wellington).
- Paul L.J., Saunders B. 2001. A description of the commercial fishery for school shark, *Galeorhinus galeus*, in New Zealand, 1945 to 1999. New Zealand Fisheries Assessment Report 2001/32. 63p.
- Peres M.B., Vooren C.M. 1991. Sexual development, reproductive cycle, and fecundity of the school shark *Galeorhinus galeus* off southern Brazil. Fishery Bulletin, U.S. 89(4): 655-667.
- Seafood Industry Council (SeaFIC). 2003a. SCH 3 Adaptive Management Programme Proposal for the 2004-05 Fishing year. 50p. (Unpublished report held at the Seafood Industry Council, Wellington).
- Seafood Industry Council (SeaFIC). 2003b. SCH 5 Adaptive Management Programme Proposal for the 2004-05 Fishing year. 49p. (Unpublished report held at the Seafood Industry Council, Wellington).
- Seafood Industry Council (SeaFIC). 2003c. SCH 7 Adaptive Management Programme Proposal for the 2004-05 Fishing year. 42p. (Unpublished report held at the Seafood Industry Council, Wellington).
- Seafood Industry Council (SeaFIC). 2003d. SCH 8 Adaptive Management Programme Proposal for the 2004-05 Fishing year. 42p. (Unpublished report held at the Seafood Industry Council, Wellington).
- Smith P.J. (2009). Review of genetic studies of rig and school shark. *Final research report for Ministry of Fisheries research project No. INS200803*. 16 p.
- Starr P.J. 2005. CPUE indices for proper, *Polyprion* spp., when targeted and as a bycatch in four New Zealand fisheries, 1990-2003. New Zealand Fisheries Assessment Report 2005/51. 29p.
- Starr P.J., Kendrick T.H., Lydon G.J., and Bentley N. 2007a. Report to the Adaptive Management Programme Fishery Assessment Working Group: Two year review of the SCH 3 Adaptive Management Programme. AMP-WG-07-08. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Starr P.J., Kendrick T.H., Lydon G.J., Bentley N. 2007b. Report to the Adaptive Management Programme Fishery Assessment Working Group: Two year review of the SCH 5 Adaptive Management Programme. AMP-WG-07-09. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Starr P.J., Kendrick T.H., Lydon G.J., Bentley N. 2007c. Report to the Adaptive Management Programme Fishery Assessment Working Group: Two year review of the SCH 7 Adaptive Management Programme. AMP-WG-07-15. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Starr P.J., Kendrick T.H., Lydon G.J., Bentley N. 2007d. Report to the Adaptive Management Programme Fishery Assessment Working Group: Two year review of the SCH 8 Adaptive Management Programme. AMP-WG-07-16. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Starr P.J., Kendrick T.H., Bentley N. 2010. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for BYX 1. Document 2010/04-v3, 86 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3871.aspx>)
- Starr P.J., Kendrick T.H. 2010a. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation and CPUE analysis for SCH 1. Document 2010/05-v3, 85 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3872.aspx>)
- Starr P.J., Kendrick T.H. 2010b. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation and CPUE analysis for SCH 2. Document 2010/06-v2, 64 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3873.aspx>)
- Starr P.J., Kendrick T.H., Bentley, N. 2010a. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 3. Document 2010/07-v2, 62 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3874.aspx>)
- Starr P.J., Kendrick T.H., Bentley N. 2010b. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 5. Document 2010/08-v2, 65 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3875.aspx>)
- Starr P.J., Kendrick T.H., Bentley N. 2010c. Report to the Adaptive Management Programme Fishery Assessment Working Group: Characterisation, CPUE analysis and logbook data for SCH 7 and SCH 8. Document 2010/09-v2, 149 p. (Unpublished document held by the Ministry of Fisheries, Wellington, N.Z.) (<http://cs.fish.govt.nz/forums/thread/3876.aspx>)
- Teirney L.D., Kilner A.R., Millar R.E., Bradford E., Bell J.D. 1997. Estimation of recreational catch from 1991-92 to 1993-94. New Zealand Fisheries Assessment Research Document 1997/15. 43p.
- Wilson D., Curtotti R., Begg G., Phillips K. (eds) 2008. *Fishery Status Reports, 2008: status of fish stocks and fisheries managed by the Australian Government*, Bureau of Rural Sciences & Australian Bureau of Agricultural and Resource Economics, Canberra. 395p.