


**BARRACOUTA (BAR)***(Thyrsites atun)*

Manga, maka

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Barracouta are caught in coastal waters around mainland New Zealand, The Snares and Chatham Islands, down to about 400 m and have been managed under the Quota Management System since 1 October 1986. Catches by New Zealand vessels increased significantly in the late 1960s and total annual catch peaked at about 47 000 t in 1977, with the addition of foreign vessels around New Zealand. Between 1983–84 and 2013–14, catches fluctuated between 18 000 and 29 000 t per annum (Table 3), at an average 25 000 t. Figure 1 shows the historical landings and TACC values for the main BAR stocks.

**Table 1: Reported landings (t) by nationality from 1977 to 1987–88.**

Fishing Year	New Zealand		Foreign			Total (FSU)	Total (QMS)
	Domestic	Chartered	Japan	Korea	USSR		
1977	4 697	0	34 357	8 109	0	47 163	-
1978–79	5 335	58	4 781	2 481	0	12 655	-
1979–80	7 748	6 679	4 339	3 879	47	22 922	-
1980–81	10 058	4 995	4 227	15	60	19 355	-
1981–82	12 055	11 077	2 813	373	0	26 328	-
1982–83	10 814	7 110	1 746	1 888	31	21 589	-
1983–83*	7 763	2 961	803	1 115	0	12 642	-
1983–84	12 390	10 226	1 786	4 355	0	28 757	-
1984–85	7 869	10 425	1 430	5 252	0	24 976	-
1985–86	8 427	7 865	1 371	815	0	18 478	-
1986–87	9 829	13 732	1 575	742	0	25 878	27 660†
1987–88	9 335	12 077	896	609	0	22 971	26 607†

\* 6 month changeover in fishing years.

† The discrepancies between QMS and FSU total landings are due to under-reporting to the FSU.

Over 99% of the recorded catch is taken by trawlers. Major target fisheries have been developed on spring spawning aggregations (Chatham Islands, Stewart Island, west coast South Island and northern and central east coast South Island) as well as on summer feeding aggregations, particularly around The Snares and on the east coast of the South Island. Barracouta also comprise a significant proportion of the bycatch in the west coast North Island jack mackerel and The Snares squid fisheries. Catches have increased in recent years in BAR 1 to the level of the TAC, but have dropped in BAR 4

in the last three years. The TACC in BAR 5 was reduced from 9282 t to 7470 t on 1 October 1998 with a 2 t customary and 3 t recreational allocation and a TAC of 7475 t. Recent catches have fluctuated about the new TAC in this fishery. In BAR 7 the catch limit was exceeded from 2004–05 to 2006–07 (catches nearly reached 15 000 t in 2006–07), but catch has decreased since, to well below the TAC.

**Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.**

Year	BAR 1	BAR 4	BAR 5	BAR 7	Year	BAR 1	BAR 4	BAR 5	BAR 7
1931-32	4	0	0	0	1957	163	0	20	80
1932-33	55	0	0	77	1958	146	0	15	78
1933-34	5	0	1	0	1959	139	0	18	71
1934-35	36	0	0	52	1960	117	0	13	90
1935-36	1	0	0	0	1961	187	0	22	68
1936-37	26	0	0	35	1962	104	0	25	44
1937-38	21	0	0	26	1963	63	0	4	20
1938-39	91	0	22	55	1964	66	0	4	21
1939-40	107	0	27	50	1965	111	0	1	76
1940-41	153	0	53	30	1966	62	0	1	116
1941-42	212	0	86	17	1967	53	0	1	178
1942-43	371	0	151	20	1968	10 113	0	3	1 196
1943-44	192	0	79	7	1969	8 499	0	2	5 756
1944	247	0	97	50	1970	12 984	0	2	3 960
1945	306	0	114	32	1971	11 327	0	191	4 006
1946	391	0	125	63	1972	29 307	2	86	3 487
1947	590	0	213	45	1973	14 856	0	79	4 698
1948	466	0	172	27	1974	23 420	0	106	9 028
1949	425	0	169	40	1975	8 985	0	855	6 257
1950	430	0	153	76	1976	19 124	5	495	6 795
1951	266	0	95	47	1977	69 81	9 095	2 041	33 266
1952	190	0	56	68	1978	6 833	17	1 162	6 918
1953	202	0	41	77	1979	6 474	4 057	3 380	5 263
1954	166	0	35	38	1980	5 649	1 854	7 867	5 146
1955	139	0	14	58	1981	6 993	2 030	8 311	11 141
1956	165	0	16	45	1982	5 393	787	6 909	7 064

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

**Table 3: Reported landings (t) of barracouta by Fishstock from 1983–84 to 2012–13 and actual TACCs (t) from 1986–87 to 2013–14. QMS data from 1986-present.**

Fishstock FMAs	BAR 1		BAR 4		BAR 5		BAR 7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	7 805	-	1 743	-	11 291	-	7 222	-
1984–85*	5 442	-	1 909	-	12 487	-	4 425	-
1985–86*	5 395	-	1 509	-	6 380	-	4 536	-
1986–87	8 877	8 510	3 084	3 010	7 653	9 010	8 046	10 510
1987–88	9 256	8 837	1 775	3 010	6 457	9 011	9 117	10 603
1988–89	5 838	9 426	946	3 010	5 323	9 011	8 071	10 702
1989–90	9 209	9 841	1 349	3 016	5 960	9 282	7 050	10 925
1990–91	9 401	9 957	1 399	3 016	8 817	9 282	7 138	10 925
1991–92	6 733	9 957	1 156	3 016	6 897	9 282	7 326	10 925
1992–93	9 032	9 969	2 251	3 016	7 019	9 282	10 141	10 925
1993–94	7 299	9 969	606	3 016	3 410	9 282	8 030	10 925
1994–95	10 023	9 969	331	3 016	2 645	9 282	9 345	10 925
1995–96	11 252	9 969	2 234	3 016	4 255	9 282	8 593	10 925
1996–97	11 873	11 000	1 081	3 016	2 839	9 282	10 203	10 925
1997–98	11 543	11 000	1 966	3 016	6 167	9 282	8 717	10 925
1998–99	9 229	11 000	459	3 016	7 302	7 470	4 427	10 925
1999–00	10 032	11 000	1 911	3 016	6 205	7 470	3 288	10 925
2000–01	7 118	11 000	2 122	3 016	6 101	7 470	6 890	10 925
2001–02	6 900	11 000	1 160	3 019	5 883	7 470	7 655	11 173
2002–03	7 595	11 000	573	3 019	7 843	7 470	9 025	11 173
2003–04	5 949	11 000	477	3 019	6 919	7 470	9 114	11 173
2004–05	6 085	11 000	98	3 019	8 593	7 470	12 156	11 173
2005–06	7 030	11 000	687	3 019	9 479	7 470	10 685	11 173
2006–07	5 351	11 000	3 233	3 019	6 334	7 470	14 699	11 173
2007–08	5 987	11 000	2 975	3 019	8 561	7 470	10 451	11 173


## BARRACOUTA (BAR)

**Table 3 [continued]**

Fishstock FMAs	BAR 1		BAR 4		BAR 5		BAR 7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
2008-09	8 861	11 000	968	3 019	7 659	7 470	8 955	11 173
2009-10	10 635	11 000	1 223	3 019	6 951	7 470	9 642	11 173
2010-11	11 420	11 000	1 190	3 019	8 201	7 470	6 129	11 173
2011-12	9 305	11 000	1 423	3 019	7 071	7 470	8 643	11 173
2012-13	9 740	11 000	706	3 019	7 931	7 470	6 897	11 173
2013-14	11 309	11 000	1 4832	3 019	6 886	7 470	6 637	11 173

Fishstock FMAs	BAR 10		Total	
	Landings	TACC	Landings	TACC
1983-84*	0	-	28 061	-
1984-85*	0	-	24 263	-
1985-86*	0	-	17 820	-
1986-87	0	10	27 660	31 050
1987-88	0	10	26 605	31 471
1988-89	0	10	20 178	32 159
1989-90	0	10	23 568	33 073
1990-91	0	10	26 755	33 190
1991-92	0	10	22 212	33 190
1992-93	0	10	28 443	33 202
1993-94	0	10	19 345	33 202
1994-95	0	10	22 345	33 202
1995-96	0	10	26 334	33 202
1996-97	0	10	25 996	34 233
1997-98	0	10	28 393	34 233
1998-99	0	10	21 417	32 421
1999-00	0	10	21 436	32 421
2000-01	0	10	22 231	32 421
2001-02	0	10	21 598	32 672
2002-03	0	10	25 036	32 672
2003-04	0	10	22 459	32 672
2004-05	0	10	26 919	32 672
2005-06	0	10	27 881	32 672
2006-07	0	10	29 617	32 672
2007-08	0	10	27 968	32 672
2008-09	0	10	26 443	32 672
2009-10	0	10	28 451	32 672
2010-11	0	10	26 937	32 672
2011-12	0	10	26 442	32 672
2012-13	0	10	24 973	32 672
2013-14	0	10	26 313	32 672

\* FSU data.



**Figure 1: Reported commercial landings and TACC for the four main BAR stocks. BAR 1 (Auckland East), [Continued on next page].**






Figure 1: [Continued] Reported commercial landings and TACC for the four main BAR stocks. From top to bottom: BAR 4 (Chatham Rise), and BAR 5 (Southland), BAR7 (Challenger).

## BARRACOUTA (BAR)

### 1.2 Recreational fisheries

Barracouta are commonly encountered by recreational fishers in New Zealand, more frequently in the southern half of BAR 7 and BAR 1. Barracouta are typically harvested as bait for other fishing rather than for consumption. They are predominantly taken on rod and reel (97.9%) with a small proportion taken by net methods (1.7%). The catch is taken predominantly from boat (95.5%) with a small proportion from land based fishers (4.5%).

#### 1.2.1 Management controls

The main method used to manage recreational harvests of barracouta is daily bag limits. General spatial and method restrictions also apply. Fishers can take up to 30 barracouta as part of their combined daily bag limit in the Fiordland and Southland Fishery Management Areas. There is currently no bag limit in place in the other Fishery management Areas.

#### 1.2.2 Estimates of recreational harvest

There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and, offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for barracouta were calculated using an offsite approach, the offsite regional telephone and diary survey approach. Estimates for 1996 came from a national telephone and diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2005). The harvest estimates provided by these telephone diary surveys (Table 3) are no longer considered reliable.

In response to the cost and scale challenges associated with onsite methods, in particular the difficulties in sampling other than trailer boat fisheries, offsite approaches to estimating recreational fisheries harvest have been revisited. This led to the development and implementation of a national panel survey for the 2011–12 fishing year (Wynne-Jones et al 2014). The panel survey used face-to-face interviews of a random sample of New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and catch information collected in standardised phone interviews. Note that the national panel survey estimate does not include recreational harvest taken under s111 general approvals. Recreational catch estimates from the national panel survey are given in Table 4.

**Table 4: Recreational harvest estimates for barracouta stocks. Early surveys were carried out in different years in the regions: South in 1991–92, Central in 1992–93, and North in 1993–94 (Teirney et al 1997). The estimated Fishstock harvest is indicative in these surveys and made by combining estimates from the different years. Some early survey harvests are presented as a range to reflect the considerable uncertainty in the estimates. The telephone/diary surveys ran from December to November but are denoted by the January calendar year. The national panel survey ran through the October to September fishing year but is denoted by the January calendar year. A mean weight of 2.14kg was used for the national panel survey.**

Fishstock	Survey	Total		
		Number	CV	Survey harvest (t)
BAR 1	1992	27 000	47%	30–90
BAR 7	1992	2 100	44%	-
BAR 1	1993	17 000	22%	25–35
BAR 7	1993	15 600	24%	25–35
BAR 1	1996	68 000	8%	160–190
BAR 7	1996	74 000	15%	160–220
BAR 1	2000	156 000	35%	182–377
BAR 5	2000	2 000	51%	2–7
BAR 7	2000	35 000	28%	68–120
BAR 1	2012	22 224		47.7
BAR 5	2012	666		1.4
BAR 7	2012	16 743		35.9
All combined	2012	39 652	18%	85.05

### 1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial take is not available.

### 1.4 Illegal catch

Quantitative information on the level of illegal catch is not available.

### 1.5 Other sources of mortality

There may have been considerable amounts of barracouta discarded prior to the QMS, either because of quota restrictions under the deepwater policy, low value, or undesirable small size fish. There is also likely to be some mortality associated with escapement from trawl nets. Some discarding may also have occurred in BAR 1 because of the lack of quota availability and the high deemed value in relation to the low value of the fish.

## 2. BIOLOGY

Barracouta spawn mainly in late-winter/spring (August–September) on the east and west coasts of both of the main islands, and in late spring (November–December) in Southland and in the Chatham Islands. Some spawning activity may also extend into summer/autumn. Sexual maturity is reached at about 50–60 cm fork length (FL) at about 2–3 years of age.

Juvenile barracouta have been recorded from inshore areas (less than 100 m) all around New Zealand and the Chatham Islands, although they appear to be less common on the west coast of the South Island. Adult fish are found down to about 400 m depth. Tagged barracouta have moved considerable distances to spawn (up to 500 nautical miles).

No age data is available for the period prior to the onset of commercial fishing, which developed rapidly from 1968. Ageing studies carried out in the mid-1970s showed that the maximum age rarely exceeded 10 years. Data have been validated for fish up to 3 years old by following modal progressions over time.

$M$  was estimated using the equation  $M = \log_e 100/\text{maximum age}$ , where maximum age is the age to which 1% of the population survives in an unexploited stock. Using 10 years for the maximum age suggests an  $M$  of up to 0.46. The effect of fishing on age structure prior to the mid-1970s is unknown, but  $M$  is unlikely to be less than 0.3, which has been assumed in previous stock assessments.

Biological parameters relevant to the stock assessment are shown in Table 5.

**Table 5: Estimates of biological parameters.**

Fishstock	Estimate				Source
<u>1. Natural mortality (<math>M</math>)</u>					
All-both sexes	Less than 0.46 $M = 0.30$ considered best estimate for all areas for both sexes				Hurst (unpub. data)
<u>2. Weight = <math>a(\text{length})^b</math> (Weight in g, length in cm fork length).</u>					
	Females		Males		
	a	b	a	b	
BAR 4	0.0074	2.94	0.0117	2.82	Hurst & Bagley (1992)
BAR 5	0.0075	2090	0.0075	2.90	Hurst & Bagley (1992)
<u>3. Von Bertalanffy growth parameters</u>					
	Both sexes				
	$K$	$t_0$	$L_\infty$		
Tasmania	0.45	0.166	91.17	(unconstrained)	Grant et al (1978)
	0.42	-0.25	91.01	(constrained, $t_0$ fixed)	

### 3. STOCKS AND AREAS

There are no new data which would alter the stock boundaries given in previous assessment documents.

Four barracouta management areas were established in 1983, based on knowledge at the time: EEZ areas E + F, G + H, B + C and D. Stock boundaries are not well understood, but the Chatham Islands stock is probably separate. However, there may be some overlap between mainland stock management areas as currently defined from analysis of tagging data, commercial fishery data, biological data (i.e., length frequencies, otoliths, parasites, spawning areas and seasons) and from seasonal relative biomass estimates. In particular, it appears that there is considerable overlap of Southland fish with other areas, probably the west coast of the South Island and possibly the east coast as well. However, there is not enough data at this stage to alter the existing stock boundaries.

### 4. STOCK ASSESSMENT

There are no stock assessments available for any barracouta stocks and TACCs have remained constant in all stocks since 2001-02. For BAR 4, 5 and 7, McGregor (in prep.) characterised the fisheries and estimated CPUE indices for the fisheries on the WCNI and WCSI (BAR 7) and the southern Snares fishery (BAR 5). In BAR 4 the fishery has been highly variable and no standardised analysis is possible.

A time series of trawl surveys was carried out in the Southland area (QMA 5) in February–March from 1993 to 1996 using *Tangaroa* (Table 6). Trawl surveys on the east and west coasts of the South Island in autumn using *Kaharoa* may help interpretation of trends in biomass around the South Island. The long time series of trawl surveys on the Chatham Rise (deeper than 200m) and Sub-Antarctic (deeper than 300m) using *Tangaroa* are not considered to adequately survey the preferred depth range of barracouta.

#### 4.1 BAR 1 Auckland (E), Central (E), South-East (Coast)

##### 4.1.1 Estimates of fishery parameters and abundance

The results from trawl surveys carried out during the mid 1980s (sometimes from a variety of different vessels) were used to provide an approximate estimate of minimum absolute biomass. This approach required an assumption about catchability to convert the trawl survey catches to estimates of absolute biomass. This method is now considered obsolete and the estimates of absolute biomass have not been included.


##### 4.1.2 Biomass estimates

The ECSI winter surveys from 1991 to 1996 in 30–400 m were replaced by summer trawl surveys (1996–97 to 2000–01) which also included the 10–30 m depth range, but these were discontinued after the fifth in the annual time series because of the extreme fluctuations in catchability between surveys (Francis et al. 2001). The winter surveys were reinstated in 2007 and this time included additional 10–30 m strata in an attempt to index elephantfish and red gurnard which were included in the list of target species. Only 2007, 2012, and 2014 surveys provide full coverage of the 10–30 m depth range.

The 2014 barracouta biomass estimate was the highest recorded in the east coast South Island winter trawl survey time series core strata (30–400 m) (Table 6, Figure 2). Biomass has been steadily increasing and in 2014 was more than four-fold larger than the average biomass of the early 1990s. The additional biomass captured in the 10–30 m depth range accounted for 15% and 6% of the biomass in the core plus shallow strata (10–400 m) for 2007 and 2012 respectively, but was less than 1% in 2014, however shallow strata should continue to be monitored for this species (Table 5, Figure 2).

### 4.1.3 Length frequency distributions

The length distributions from the east coast South Island winter trawl survey show at least three clear pre-recruit modes at about 20 cm, 25 cm, and 50 cm (combined males, females, and unsexed) consistent with ages of 0+, 1+, and 2+. Length frequency distributions are consistent among the surveys, showing the presence of the pre-recruited cohorts, with indications that these could be tracked through time (modal progression)(Beentjes et al., 2015). The addition of the 10–30 m depth range does not change the shape of the length distributions.



**Figure 2: Barracouta total biomass and 95% confidence intervals for the all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012, and 2014**

**Table 6: Relative biomass indices (t) and coefficients of variation (CV) for barracouta for east coast South Island (ECSI) - winter, east coast North Island (ECNI), west coast South Island (WCSI) and Southland survey areas. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16 and 17). –, not measured; NA, not applicable.**

Region	Fishstock	Year	Trip number	Total Biomass estimate	CV (%)	Total Biomass estimate	CV (%)
ECSI (winter)	BAR 1			30–400 m		10–400 m	
		1991	KAH9105	8 361	29	-	-
		1992	KAH9205	11 672	23	-	-
		1993	KAH9306	18 197	22	-	-
		1994	KAH9406	6 965	34	-	-
		1996	KAH9608	16 848	19	-	-
		2007	KAH0705	21 132	17	24 939	19
		2008	KAH0806	25 544	16	-	-
		2009	KAH0905	33 360	16	-	-
		2012	KAH1207	34 325	17	36 526	16
		2014	KAH1402	46 563	19	46 903	19
ECNI	BAR 1	1993	KAH9304	2 673	15	-	-
		1994	KAH9402	8 433	33	-	-
		1995	KAH9502	2 103	29	-	-
		1996	KAH9602	2 495	23	-	-



**BARRACOUTA (BAR)**


**Table 6 [Continued]**

Region	Fishstock	Year	Trip number	Total Biomass	CV (%)	Total	CV (%)
WCSI	BAR 7	1992	KAH9203	2 478	14	-	-
		1994	KAH9404	5 298	16	-	-
		1995	KAH9504	4 480	13	-	-
		1997	KAH9701	2 993	19	-	-
		2000	KAH0004	1 787	11	-	-
		2003	KAH0304	4 485	20	-	-
		2005	KAH0503	2 763	13	-	-
		2013	KAH1305	3 423	16	-	-
Southland	BAR 5	1993	TAN9301	11 587	18	-	-
		1994	TAN9402	6 151	20	-	-
		1995	TAN9502	4 539	17	-	-
		1996	TAN9604	7 693	19	-	-

**4.2 BAR 5 Southland, Sub-Antarctic**

**4.2.1 CPUE indices**

McGregor (in prep.) used merged (stratified) and unmerged (tow level) data to fit CPUE indices for the southern Snares fishery. The WG considered that the trip level CPUE was an adequate time series of CPUE to monitor this stock. After being flat for a long period the CPUE shows a recent increase since 2008 possibly from a recent recruitment pulse. The current stock status is unknown.




**Figure 3 : South CPUE Model 1a (South) Arithmetic, geometric, and standardised CPUE indices for BAR 1990-2011.**

### 4.3 BAR 7 Challenger, Central (W) Auckland (W)


#### 4.3.1 CPUE indices

McGregor (in prep.) looked at the separate fisheries on the WCNI and WCSI. The three CPUE options for the WCNI all gave similar patterns to the inshore *Kaharoa* WCSI trawl survey. The WG considered that the tow level CPUE was the best data to use to monitor this stock. The CPUE shows an increasing trend from 2000 to 2004 and then generally flat (Figure 4).



**Figure 4: West Coast CPUE for Models 2b (tow level), 3 (JMA target) and 4 (no target) and Trawl Survey abundance index for calendar years 1990-2010. Model 3 (JMA target) is actually based on fishing years, months Nov-May, whereas the other models here are calendar year, Jun-Nov. Trawl survey is based on fishing year.**

The WCSI data series shows a similar increase from 2000 and then generally flat, for the tow level CPUE based on all target from June to October (Figure 5).



**Figure 5: West Coast South Island current and previous CPUE, West Coast North Island CPUE and trawl survey abundance index for calendar years 1990-2010. Trawl survey is based on fishing year.**

## BARRACOUTA (BAR)

### 4.4 Yield estimates and projections

It is not feasible to estimate *MCY* from commercial landings data for most Fishstocks (except for BAR 1), as the amount of effort has varied considerably since the beginning of the fishery in the late 1960s i.e., foreign licensed access has declined, effort was encouraged by subsidies in 1979 and 1981, an unknown amount of fish has been and may still be dumped, and effort is related to the availability of more preferred, higher value species. These, and other factors, also result in CPUE data being of limited use.

Estimates of current biomass are not available and *CAY* cannot be estimated.

#### 4.4.1 Auckland (East), Central (East) and South-East (Coast) (BAR 1)

*MCY* was estimated using the equation  $MCY = cY_{AV}$  (Method 4), where  $Y_{AV}$  average estimated catch from 1968–1975 and  $c = 0.7$ . The estimated average catch includes 2000 t which is assumed to have been caught and either dumped or not reported. Fishing activity is assumed to have been on the total stock, even though the entire area was not fished. Due to problems with QMA boundaries not corresponding to the fishing history boundaries, 500 t is subtracted and added to BAR 7.

$$MCY = 0.7 * (12\ 000\ t - 500\ t) = 8050\ t.$$

The level of risk to the stock by harvesting the population at the estimated *MCY* value cannot be determined. However, the risk is probably low given the sustainability of catches at about the *MCY* level since 1970.

*MCY* has not been determined for the other Fishstocks.

### 4.5 Other factors

The relationship of the southern area stock to the east and west coast South Island stocks is uncertain, so these areas have been treated separately as in the past. However, if fish from BAR 5 overlap significantly with other South Island stocks, then the *MCY*s for all Fishstocks on the South Island may all need adjusting downward.

Barracouta are part of the shelf (30–300 m) mixed fishery and are usually the dominant species in these depths around the South Island (except perhaps in good red cod years in Canterbury Bight). Any increase or decrease in barracouta quotas will have overflow effects onto bycatch species. The economics of targeting on barracouta is probably affected by its availability relative to other more preferred species and this will, in turn, affect fishing patterns.

An analysis of trends in biomass of the Southland fishery suggests that recruitment may have been relatively low in the years after 1989 and that biomass may have declined between surveys by the *Shinkai Maru* (1981 and 1986) and the *Tangaroa* (annually 1993 to 1996). The scale of decline appeared to be greater than could be explained by different catching efficiency of the two vessels.

## 5. STATUS OF THE STOCKS

Estimates of current and reference biomass are not available for any barracouta stocks and therefore it is not known if current TACCs and recent catches are sustainable or whether they are at levels which will allow the stocks to move towards a size that will support the maximum sustainable yield.

Yield estimates and reported landings are summarised in Table 7.

**Table 7: Summary of yields (t), TACCs (t), and reported landings (t) for barracouta for the most recent fishing year.**

Fishstock	FMAAs	<i>MCY</i>	2013–14 Actual TACC	2013–14 Reported landings	
BAR 1	Auckland (East), Central (East), South-East (Coast)	1, 2, & 3	8 050	11 000	11 309
BAR 4	South-East (Chatham)	4	-	3 019	1 482
BAR 5	Southland, Sub-Antarctic	5 & 6	-	7 470	6886

Table 7 [Continued]

		FMA	MCY	2013–14 Actual TACC	2013–14 Reported landings
Fishstock					
BAR 7	Challenger, Central (West), Auckland (West)	7, 8, & 9	-	11 173	6 637
BAR 10	Kermadec	10	-	10	0
Total				32 672	26314

## 6. FOR FURTHER INFORMATION

- Beentjes, M.P.; MacGibbon, D.J. (2013). Review of QMS species for inclusion in the east coast South Island winter trawl survey reports. *New Zealand Fisheries Assessment Report 2013/35*. 102 p.
- Beentjes, M.P.; MacGibbon, D.; Lyon, W.S. (2015). Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2014 (KAH1402). *New Zealand Fisheries Assessment Report 2015/14*. 136 p.
- Bagley, N W; Hurst, R J (1995) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1994 (TAN 9402). *New Zealand Fisheries Data Report 57*. 50 p.
- Bagley, N W; Hurst, R J (1996) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1995 (TAN 9502). *New Zealand Fisheries Data Report 73*. 47 p.
- Bagley, N W; Hurst, R J (1996) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1996 (TAN 9604). *New Zealand Fisheries Data Report 77*. 51 p.
- Beentjes, M P (1995) Inshore trawl survey of the Canterbury Bight and Pegasus Bay, May–June 1992 (KAH9205). *New Zealand Fisheries Data Report 55*. 58 p.
- Beentjes, M P (1995) Inshore trawl survey of the Canterbury Bight and Pegasus Bay, May–June 1993 (KAH9306). *New Zealand Fisheries Data Report 56*. 56 p.
- Beentjes, M P; Wass, R T (1994) Inshore trawl survey of the Canterbury Bight and Pegasus Bay, May–June 1991 (KAH9105). *New Zealand Fisheries Data Report 48*. 49 p.
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16 27 p. (Unpublished report held by NIWA library, Wellington.)
- Drummond, K L; Stevenson, M L (1995) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1992 (KAH9204). *New Zealand Fisheries Data Report 63*. 58 p.
- Drummond, K L; Stevenson, M L (1995) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1994 (KAH9404). *New Zealand Fisheries Data Report 64*. 55 p.
- Drummond, K L; Stevenson, M L (1996) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1994 (KAH9504). *New Zealand Fisheries Data Report 74*. 60 p.
- Grant, C J; Cowper, T R; Reid, D D (1978) Age and growth of snoek, *Leionura atun* (Euphrasen) in South-eastern Australian waters. *Australian Journal of Marine and Freshwater Research 29*: 435–444.
- Harley, S J; Horn, P L; Hurst, R J; Bagley, N W (1999) Analysis of commercial catch and effort data and age determination and catch-at-age of barracouta in BAR 5. New Zealand Fisheries Assessment Research Document 1999/39. 39 p. (Unpublished document held by NIWA library, Wellington.)
- Hatanaka, H; Uozumi, Y; Fukui, J; Aizawa, M; Hurst, R J (1989) Japan New Zealand trawl survey off southern New Zealand, October–November 1983. *New Zealand Fisheries Technical Report 9*. 52 p.
- Hurst, R J (1988a) The barracouta, *Thyrsites atun*, fishery around New Zealand: historical trends to 1984. *New Zealand Fisheries Technical Report 5*. 43 p.
- Hurst, R J (1988b) Barracouta. New Zealand Fisheries Assessment Research Document 1988/8. 29p. (Unpublished document held by NIWA library, Wellington.)
- Hurst, R J; Bagley, N W (1987) Results of a trawl survey of barracouta and associated finfish near the Chatham Islands, New Zealand, December 1984. *New Zealand Fisheries Technical Report 3*. 44 p.
- Hurst, R J; Bagley, N W (1989) Movements and possible stock relationships of the New Zealand barracouta, *Thyrsites atun*, from tag returns. *New Zealand Journal of Marine and Freshwater Research 23*: 105–111.
- Hurst, R J; Bagley, N W (1992) Results of a trawl survey of barracouta and associated finfish near the Chatham Islands, New Zealand, December 1985. *New Zealand Fisheries Technical Report 30*. 36 p.
- Hurst, R J; Bagley, N W (1997). Trends in Southland trawl surveys of inshore and middle depth species 1993–1996. *New Zealand Fisheries Technical Report No.50*. 66p.
- Hurst, R J; Bagley, N W (1994) Trawl survey of middle depth and inshore bottom species off Southland, February–March 1993 (TAN 9301). *New Zealand Fisheries Data Report 52*. 58 p.
- Hurst, R J; Bagley, N W; Uozumi, Y (1990) New Zealand - Japan trawl survey of shelf and upper slope species off southern New Zealand, June 1986. *New Zealand Fisheries Technical Report 18*. 50 p.
- Hurst, R J; Fenaughty, J M (1985) Report on biomass surveys 1980–84: summaries and additional information. Fisheries Research Division Internal Report 21: 53 p. (Unpublished report held by NIWA library, Wellington.)
- Langley, A D; Bentley, N (2002) An examination of the stock structure of barracouta (*Thyrsites atun*) around the South Island of New Zealand. *New Zealand Fisheries Assessment Report 2002/30*. 57 p.
- Langley, A D; Walker, N A (2002) Characterisation of the barracouta (*Thyrsites atun*) fishery in BAR 1. *New Zealand Fisheries Assessment Report 2002/44*. 37 p.
- Langley, A D; Walker, N (2002) CPUE analysis of the southeast South Island BAR 1 fishery, 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report 2002/11*. 28 p.
- McGregor, V. (in prep.). Fishery characterisation and standardised CPUE analyses for barracouta *Thyrsites atun*, (Euphrasen, 1791) (Gempylidae), 1989 to 2010–11 New Zealand Fisheries Assessment Report (draft)
- Stevenson, M L (1996) Bottom trawl survey of the inshore waters of the east coast North Island, February–March 1995 (KAH9502). *New Zealand Fisheries Data Report 78*. 57 p.
- Stevenson, M L (1998) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 1997 (KAH9701). *NIWA Technical Report 12*. 70 p.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)