


COCKLES (COC)

(*Austrovenus stutchburyi*)
Tuangi



1. INTRODUCTION

Cockles are important shellfish both commercially and for non-commercial fishers.

Commercial picking of cockles, *Austrovenus stutchburyi*, is carried out on Snake Bank, Whangarei Harbour (FMA 1), Papanui and Waitati Inlets, Otago (FMA 3) and Pakawau Beach, Ferry Point and Tapu Bay in Tasman and Golden Bays (FMA 7). Cockles have also been commercially harvested since August 2009 under a special permit from Otago Harbour. Cockles were introduced into the QMS on 1 October 2002. The fishing year runs from 1 October until September 30 and catches are measured in greenweight for all stocks. There is no minimum legal size for commercial or non-commercial fishers for cockles in any stock. Cockles are managed under Schedule 6 of the Fisheries Act for all stocks listed in Table 1, which allows cockles to be returned to where they were taken as soon as practicable after the cockle is taken as long as the cockle is likely to survive.

For assessment purposes, individual reports on the largest fisheries have been produced separately:

1. Snake Bank, Whangarei Harbour, in COC 1A.
2. Papanui Inlet, Waitati Inlet, and Otago Harbour, Otago Peninsula in COC 3.
3. Tasman and Golden Bays in COC 7A.

The landings, by stock, of these cockle fisheries are dominated by catch from COC 3 (Figure 1). Landings from COC 3 are relatively stable since 2002–03; by contrast landings from COC 1A and COC 7A have generally declined over that time period.

Information on cockles that applies to all stocks is included below rather than being repeated in the reports for each fishery.

New Zealand operates a mandatory shellfish quality assurance programme for all bivalve shellfish commercial growing or harvesting areas for human consumption. Shellfish caught outside this programme can only be sold for bait. This programme is based on international best practice and managed by MPI in cooperation with the District Health Board Public Health Units and the shellfish

industry¹ and is summarised below. Before any area can be used to grow or harvest bivalve shellfish, public health officials survey both the water catchment area to identify any potential pollution issues and microbiologically sampling water and shellfish over at least a 12-month period, so all seasonal influences are explored. This information is evaluated and, if suitable, the area classified and listed by MPI for harvest. There is then a requirement for regular monitoring of the water and shellfish flesh to verify levels of microbiological and chemical contaminants. Management measures stemming from this testing include closure after rainfall, to deal with microbiological contamination from runoff. Natural marine biotoxins can also cause health risks, therefore testing for these also occur at regular intervals. If toxins are detected above the permissible level the harvest areas are closed until the levels fall below the permissible level. Products are also traceable so that the source and time of harvest can always be identified in case of contamination.

Table 1: TACC, Recreational, customary allowances and TAC (t) for all cockle stocks.

Code	Description	TACC	Recreational allowance	Customary allowance	TAC
COC 1A	Whangarei Harbour	346	25	25	396
COC 1B	East Northland	0	22	22	44
COC 1C	Hauraki Gulf and Bay of Plenty	5	32	32	69
COC 2	Central	0	2	2	4
COC 3	Otago	1 470	10	10	1 490
COC 3B	Part South East Coast	1	27	27	55
COC 4	South East (Chatham Rise)	0	1	1	2
COC 5	Southland and Sub-Antarctic	2	2	2	6
COC 7A	Nelson Bays	1 390	85	25	1 500
COC 7B	Marlborough	0	5	5	10
COC 7C	Part Challenger	0	3	3	6
COC 8	Central (Egmont)	0	1	1	2
COC 9	Auckland (West)	0	6	6	12




Figure 1: Commercial landings and the sum total (solid line) of the three main commercial COC stocks throughout time. Note that this figure does not show data prior to entry into the QMS.

2. BIOLOGY

The cockle, *Austrovenus stutchburyi*, formerly known as *Chione stutchburyi*, is a shallow-burrowing suspension feeder of the family Veneridae. It is found in soft mud to fine sand on protected beaches and enclosed shores around the North and South Islands, Stewart Island, the Chatham Islands and the Auckland Islands (Morton & Miller 1973, Spencer et al 2002). Suspension feeders such as *A. stutchburyi* tend to be more abundant in sediments with a larger grain size. Cockles have been shown

¹For full details of this programme, refer to the Animal Products (Regulated Control Scheme-Bivalve Molluscan Shellfish) Regulations 2006 and the Animal Products (Specifications for Bivalve Molluscan Shellfish) Notice 2006 (both referred to as the BMSRCS), at: <http://www.nzfsa.govt.nz/industry/sectors/seafood/bms/page-01.htm>

COCKLES (COC)

to be most abundant in sediments of below 12 percent mud in two separate studies (Thrush et al 2003, Anderson 2008). They are also common in eelgrass (e.g., *Zostera* sp.), which often co-occurs with sand flats.

Cockles are found from the lowest high water neap tide mark to the lowest part of the shore. Larcombe (1971) suggested that the upper limit is found where submergence is only 3.5 hours per day. *A. stutchburyi* is often a dominant species and densities as high as 4500 per m² have been reported in some areas. In Pauatahanui Inlet the cockle biomass was estimated at 80% (5000 t) of the total intertidal biomass in 1976 (Richardson et al 1979). Calculations based on laboratory measurements of filtration rates suggested that cockles over 35 mm shell length were capable of filtering 1.1×10^6 m³ of water or enough to filter all the water in Papanui Inlet every two tidal cycles (Pawson 2004).

Sexes are separate and the sex ratio is usually close to 1:1. Size at maturity has been estimated at about 18 mm shell length (Larcombe 1971). Spawning extends over spring and summer, and fertilisation is followed by a planktonic larval stage lasting about three weeks. Significant depression of larval settlement has been recorded for areas of otherwise suitable substrate from which all live cockles have been removed. This suggests the presence of some conditioning factor.

Work on Snake Bank also showed moderate differences among years in the level of recruitment of juveniles to the population. The variability of recruitment was estimated as $\sigma_R = 0.41$ using all available data (1983–1996) but as $\sigma_R = 0.31$ using data only from those years since the fishery has been considered to be fully developed (1991–96). Given the variability of most shellfish populations and the shortness of the time series, this is probably an underestimate of the real variability of recruitment in the Snake Bank population.

Small cockles grow faster than large cockles, but overall, maximum growth occurs on the first of January, and a period of no growth occurs at the beginning of July (Tuck & Williams 2012). Growth is slower in the higher tidal ranges and in high density beds. Significant increases in growth rates have been observed for individuals remaining in areas that have been ‘thinned out’ by simulated harvesting. Tagging work at Pakawau beach also highlighted the variability in growth that can occur within a beach (Osborne 2010).

Growth parameters and length weight relationships are listed in Table 2 (Stewart 2008, Williams et al 2009, Osborne 2010). However, considerable variability in growth has been seen in all three QMAs over time. At Snake bank (1A) growth to 30 mm has been estimated as taking between 2 and 5 years in separate studies (Martin 1984, Cryer 1997). Additional tagging work on Snake Bank from 2001 to 2010 showed that on average, cockles reach maturity (18 mm; Larcombe 1971) in their second year of growth, and recruit to harvestable size (about 28 mm SL) in about 3 to 4 years, although these results showed great variability in growth rate (tabulated in Table 8, Tuck & Williams 2012). At Pakawau beach (7A) K has varied between 0.36 and 0.41 and L_∞ between 47 and 49mm (Osborne 1992, 1999). The work of Breen et al (1999) in Papanui and Waitati Inlets, Purakanui and Otago Harbour showed no significant growth after one year and modes in the length frequency distributions did not shift when measured over four sampling periods within a year. They concluded that it was unlikely that average growth is really as slow as the results indicated, but there may be high inter-annual variability in growth.

Quite extensive movements of juveniles have been documented, but individuals over 25 mm shell length remain largely sessile, moving only in response to disturbance.

Given that cockles recruit to the spawning biomass at about 18 mm shell length, but do not recruit to commercial or non-commercial fisheries until closer to 30 mm shell length, there is some protection for the stock against egg overfishing, especially as the Snake Bank and Papanui and Waitati Inlet stocks are probably not isolated as far as recruitment of juveniles is concerned. However, this generality should be treated with some caution, given that some population of adults seems to be required to stimulate settlement of spat.

Natural mortality arises from a number of sources. Birds are a major predator of cockles (up to about 23 mm shell length). Other predators include crabs and whelks. Cockles are also killed after being

smothered by sediments shifted during storms or strong tides. A mass mortality that killed an estimated 56–63% of all cockles and 80–84% of cockles over 30 mm in shell length (MPI unpublished data) has been reported from sites within the Whangateau harbour (north of Auckland). This mortality was attributed to a potential weakening of cockles due to heat stress then mortality from a coccidian parasite and a mycobacterium². Sediments, both suspended and deposited, both impact upon cockle fitness or survival, with terrestrial sediments having greater effects than marine sediments (Gibbs & Hewitt 2004). Increasing suspended sediment concentrations have induced increased physiological stress, decreased reproductive status and decreased juvenile growth rates (Nicholls et al 2003, Gibbs & Hewitt 2004). Sediment deposition has also been shown to negatively impact upon densities of cockles (Lohrer et al 2004). The sum of these effects is seen in the distribution of cockles which decline in abundance across a number of sites with increasing mud content in the sediments, either above zero or 11% mud content, depending upon the study (Thrush et al 2003, Anderson 2008).

Experimental work on Snake Bank led to estimates of absolute mortality of 17–30% per annum, instantaneous natural mortality (M) of 0.19–0.35, with a midpoint of $M = 0.28$. The estimated mortality rates for cockles of over 30 mm shell length were slightly greater at 19–37% per annum, (M of 0.21–0.46 with a midpoint of 0.33). This higher estimate was caused by relatively high mortality rates for cockles of over 35 mm shell length and, as these are now uncommon in the population, $M = 0.30$ (range 0.20–0.40) has been assumed for yield calculations across all three stocks (Table 2). Tagging (both notch and individual numbered tags) has been ongoing on Mair Bank from 2001 to 2009 and the last recoveries occurred in 2010 (Tuck & Williams 2012). Annualised mortality estimates (M) (averaged over 3, 6 and 9 month recoveries) were 0.356 and 0.465 from studies in 2008 and 2009.

Table 2: Biological parameters used for cockle assessments for different stocks. SL = shell length, within area 7A, P = Pakawau, FP = Ferry Point, TBR = Tapu Bay/Riwaka.

	1A	3	7A
1. Natural mortality (M)	0.3	0.3	0.3
2. Weight (grams)	= $a(\text{shell length})^b = a(\text{shell length}) + b = a(\text{shell length})^b$		
a	0.00014	0.7211	P = 0.000018, FP = 0.0002, TBR = 0.00015
b	3.29	11.55	P = 3.78, FP = 3.153, TBR = 3.249
3. von Bertalanffy growth parameters	Not used instead growth = $a(\ln(\text{age in years})) + b$		
K	0.26	0.311	a = 11.452
L_∞ (mm)	35	40.95	b = 16.425
SL at recruitment to the fishery (mm)	28	28	30

3. STOCKS AND AREAS

Little is known of the stock boundaries of cockles. Given the planktonic larval phase, many populations may receive spat fall from other nearby populations and may, in turn, provide spat for these other areas. In the absence of more detailed knowledge, each commercial fishery area is managed as a discrete population.

4. FOR FURTHER INFORMATION

- Anderson, M J (2008) Animal-sediment relationships re-visited: Characterising species' distributions along an environmental gradient using canonical analysis and quantile regression splines. *Journal of Experimental Marine Biology and Ecology* 366(1–2): 16–27.
- Breen, P A; Carbines, G C; Kendrick, T H (1999) Stock assessment of cockles in Papanui and Waitati Inlets, Otago Harbour, and Purakanui, Otago. Final Report for the Ministry of Fisheries research project COC9701 dated July 1999. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Cryer, M (1997) Assessment of cockles on Snake Bank, Whangarei Harbour, for 1996. New Zealand Fisheries Assessment Research Document 97/2. 29 p. (Unpublished report held in NIWA library, Wellington.)
- Ellis, J I (2003) Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology-Progress Series* 263: 101–112.
- Gibbs, M; Hewitt, J E (2004) Effects of sedimentation on macrofaunal communities: a synthesis of research studies for ARC. *Auckland Regional Council Technical Publication* 264.
- Irwin, C (2004) The impacts of harvesting and the sustainability of a New Zealand Littleneck Clam (*Austrovenus stutchburyi*) Fishery in Papanui and Waitati Inlets, New Zealand. Marine Science. Dunedin, University of Otago. PhD.

² <http://www.biosecurity.govt.nz/media/21-08-09/cockle-death-whangateau-estuary>

COCKLES (COC)

- Larcombe, M (1971) The ecology, population dynamics, and energetics of some soft shore molluscs. . Department of Zoology, University of Auckland. PhD. 250 p.
- Lohrer, A; Thrush, S; Hewitt, J E; Berkenbusch, K; Ahrens, M; Cummings, V J (2004) Terrestrially derived sediment: response of marine macrobenthic communities to thin terrigenous deposits. *Marine Ecology Progress Series* 273: 121–138.
- McKinnon, J (1996) Studies of the age, growth and shell increment patterns in the New Zealand cockle (*Austrovenus stutchburyi*). Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- Manly, B F J; Akroyd, J M; Walshe, K A R (2002) Two-phase stratified random surveys on multiple populations at multiple locations. *New Zealand Journal of Marine and Freshwater Research* 36: 581–591.
- Martin, N D (1984) *Chione stutchburyi* population responses to exploitation. Unpublished MSc Thesis, University of Auckland, Auckland.
- Morton, J; Miller, M (1973) *The New Zealand Sea Shore*. Collins, Auckland. 653 p.
- Nicholls, P; Hewitt, J; Halliday, J (2003) Effects of suspended sediment concentrations on suspension and deposit feeding marine macrofauna. *Auckland Regional Council Technical Publication* 211. 40 p.
- Osborne, T A (1992) Biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared for Westhaven Shellfish Co. Ltd. 27 p.
- Osborne, T A (1999) 1999 biomass survey and stock assessment of the New Zealand littleneck clam (*Chione stutchburyi*) on Pakawau Beach, Golden Bay. Unpublished report prepared by Osborne Research Co. Ltd. for Westhaven Shellfish Co. Ltd. 15 p.
- Osborne, T A (2010) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) in area COC 7A: Tapu Bay, Ferry Point, and Pakawau. *New Zealand Fisheries Assessment Report* 2010/44.
- Pawson, M (2004) The cockle *Austrovenus stutchburyi* and chlorophyll depletion in a southern New Zealand Inlet. Marine Science. Dunedin, University of Otago. MSc. 113 p.
- Richardson, J R; Alridge, A E; Main, W, deL (1979) Distribution of the New Zealand cockle, *Chione stutchburyi* at Pauatahanui Inlet. NZOI Oceanographic Field Report 14. 10 p.
- Spencer, H G; Willan, R C; Marshall, B A; Murray, T J (2002) Checklist of the Recent Mollusca described from the New Zealand Exclusive Economic Zone. <http://toroa.otago.ac.nz/pubs/spencer/Molluscs/index.html>
- Stewart, B (2008) Stock Assessment of Clams (*Austrovenus stutchburyi*) in Waitati Inlet, Otago, 2007. Report Prepared by Ryder Consulting Ltd. for Southern Clams Ltd.
- Williams, J R; Smith, M D; Mackay, G (2009) Biomass survey and stock assessment of cockles (*Austrovenus stutchburyi*) on Snake Bank, Whangarei Harbour, 2009. *New Zealand Fisheries Assessment Report* 2009/29. 22 p.
- Thrush, S F; Hewitt, J E; Norkko, A; Nicholls, P E; Funnell, G A; Ellis, J I (2003) Habitat change in estuaries: predicting broad-scale responses of intertidal macrofauna to sediment mud content. *Marine Ecology-Progress Series* 263: 101–112.
- Tuck, I; Williams, J (2012) Cockle growth at Snake Bank, Whangarei Harbour, 2001–10. Fisheries Research Report, Unpublished Report for Ministry for Primary Industries, 40p.