


**TUATUA (TUA)***(Paphies subtriangulata)*

Tuatua

**1. FISHERY SUMMARY**

Tuatua (*Paphies subtriangulata*) were introduced into the QMS on 1 October 2005. The fishing year runs from 1 October to 30 September, and commercial catches are measured in greenweight. In October of 2005 all TUA QMAs were allocated customary and recreational catch allowances. A breakdown of each QMA Total Allowable Catch (TAC) is listed in Table 1.

**1.1 Commercial fisheries**

QMA boundaries for tuatua were set the same as those established for FMAs, except for FMA 1 (the area between North Cape and Cape Runaway), which was divided into two QMAs, TUA 1A and TUA 1B, on either side of Te Arai Point (Pakiri Beach). The formerly specified historic commercial areas within TUA 1B (Papamoa domain to Maketu Beach, Bay of Plenty) and TUA 9 (i.e., Ninety Mile Beach, Hokianga Harbour to Maunganui Bluff, and specific areas between Maunganui Bluff to the North Head of the Kaipara Harbour) were revoked, and regulations were amended to remove the commercial daily catch limits for tuatua, which were no longer applicable. Commercial fishing was allowed to continue only in TUA 9 in the specified commercial area of the Kaipara Harbour entrance. A TACC of 43 t, which reflected the average of the reported landings taken from the Kaipara fishery between 1990–91 and 2003–04, was allocated to the TUA 9 stock in recognition that commercial tuatua fishing was constrained to the Kaipara Harbour entrance.

There is no minimum legal size (MLS) for tuatua, although fishers probably favor large individuals. Tuatua are available for harvest year-round, so there is no apparent seasonality in the fishery. Significant landings since 1989–90 have been reported from TUA 9 only (Table 2), and there have been no reported landings from TUA 5, TUA 6, and TUA 8. Landings from TUA 9 reached a peak of 192 t in 1997–98, and subsequently decreased, ranging from 4 to 76 t (average 32 t) between 1998–99 and 2003–04. This decline in commercial catches from the Kaipara bed is probably related to historic participants retiring from the fishery. The commercial effort had greatly reduced by 1992, post moratorium implementation, and catches have been influenced by the fact that commercial fishing is intermittent with only one or two fishers involved. No landings were reported from TUA 9 for 2004–05 to 2010–11. Landings of 4.881 t were recorded for TUA 9 in the 2011–12 fishing year.

## TUATUA (TUA)

**Table 1: Recreational, customary, and other mortality allowances (t); Total Allowable Commercial Catches (TACC, t) and Total Allowable Catch (TAC, t) declared for TUA in October 2005.**

Fishstock	Recreational Allowance	Customary non-commercial Allowance	Other Mortality	TACC	TAC
TUA 1A	40	40	4	0	84
TUA 1B	60	60	6	0	126
TUA 2	3	3	1	0	7
TUA 3	3	3	1	0	7
TUA 4	1	1	1	0	3
TUA 5	1	1	1	0	3
TUA 7	1	1	1	0	3
TUA 8	1	1	1	0	5
TUA 9	26	26	7	43	102

**Table 2: Reported landings (t) of tuatua (*Paphies subtriangulata*) by Fishstock from 1989–90 to the present day. Data up to 2003–04 taken from page 163 of MFish's Initial Position Paper (IPP), dated 31 March 2005, data since from CELR and CLR (early CELR and CLR data erroneously record commercial landings from FMA 9 as FMA 1 because permit holders were not filling in the forms correctly). There have been no reported landings of tuatua in TUA 5, TUA 6, and TUA 8. There were no landings reported from 2004–05 to 2010–11. Tuatua were introduced into the QMS on 1 October 2005; a TACC of 43 t was allocated (to TUA 9 only), and FMA 1 was divided into TUA 1A and TUA 1B.**

Year	TUA 1	TUA 2	TUA 3	TUA 4	TUA 7	TUA 9	Total	TACC
1989–90	0	0	0	0	0	69.015	69.015	-
1990–91	0	0	0	0	0.176	68.245	68.421	-
1991–92	0	0	0	0	1.667	82.002	83.669	-
1992–93	0	0	0	0	0.891	109.280	110.171	-
1993–94	0	0	0.042	0	0	177.165	177.207	-
1994–95	0	0	0	0	0	182.262	182.262	-
1995–96	0	0	0	0	0	100.016	100.016	-
1996–97	0	0	0.125	0	0.005	68.575	68.705	-
1997–98	0	0	0.184	0	0	192.262	192.446	-
1998–99	0	0	0	0	0	76.205	76.205	-
1999–00	0	0	0	0	0	44.450	44.450	-
2000–01	0	0	0	0	0	16.150	16.150	-
2001–02	0	0	0	0	0	4.900	4.900	-
2002–03	0	0	0	0	0	36.160	36.160	-
2003–04	0	0	0.054	0	0	34.336	34.390	-
2004–05	0	0	0	0	0	0	0	-
2005–06	0	0	0	0	0	0	0	43
2006–07	0	0	0	0	0	0	0	43
2007–08	0	0	0	0	0	0	0	43
2008–09	0	0	0	0	0	0	0	43
2009–10	0	0	0	0	0	0	0	43
2010–11	0	0	0	0	0	0	0	43
2011–12	0	0	0	0	0	4.881	4.881	43
2012–13	0	0	0	0	0	5.294	5.294	43
2013–14	0	0	0	0.02	0	0	0.02	43

### 1.2 Recreational fisheries

Tuatua support an extensive recreational fishery, with harvesting occurring in all stocks wherever there are accessible beds, particularly in the upper North Island. Tuatua are harvested entirely by hand gathering, and there is no MLS (although large tuatua are preferred).

There is a recreational daily catch limit of 150 tuatua per person, except in the Auckland - Coromandel region where the limit has been 50 per day per person since November 1999.

Currently, there are no reliable estimates of the recreational harvest of tuatua. Estimates of tuatua catch by recreational fishers have been made on three occasions (1996, 1999–2000, and 2000–01) as part of national recreational fishing (telephone and diary) surveys. These estimates indicate that the majority of recreational tuatua harvests were taken from FMA 1, moderate harvests were taken from QMA 9, and smaller quantities were taken from other areas. A review by the Marine Recreational Fisheries

Technical Working Group concluded that these estimates were not likely to be reliable. The current level of recreational harvest and its impact on the status of tuatua beds are unknown. There are concerns about the depletion of popular tuatua beds in some areas, whereas in other areas it appears they are in a healthy state.

### 1.3 Customary non-commercial fisheries

In common with many other intertidal shellfish, tuatua are an important customary species taken as kaimoana. Both oral tradition and the numerous middens of *P. triangulata* shells around the coastline clearly show that this fishery has been an important one to Maori for at least several hundred years. However, no quantitative information on the level of customary non-commercial take is available.

### 1.4 Illegal catch

The illegal catch of tuatua is probably significant in some areas, with some recreational fishers exceeding their bag limit, but no quantitative information on the level of illegal catch is available.

### 1.5 Other sources of fishing-related mortality

No quantitative information on the level of other sources of mortality is available. Tuatua are generally sedentary and beds are susceptible to localised depletion, not only by harvesting pressure, but also by habitat disturbance and degradation. Incidental mortality of tuatua is likely in the Kaipara Harbour dredge fishery if tuatua are damaged during encounters with the dredge. Changes in bank stability could arise from dredging operations and might cause additional incidental mortality. However, the level of dredge-related mortality is unknown. As suspension feeders, tuatua may also be adversely affected by high sedimentation loads in the water column. In some areas, such as Ninety Mile Beach, Dargaville and Muriwai, vehicles driven along the beach pass directly over tuatua beds, increasing mortality either directly by damaging tuatua or indirectly by adversely modifying surface sand conditions leading to desiccation of tuatua.

## 2. BIOLOGY

Tuatua (*Paphies subtriangulata*) belong to the family Mesodesmatidae, a group of moderate to large wedge-shaped surf clams that include toheroa (*Paphies ventricosum*), deepwater tuatua (*Paphies donacina*), and pipi (*Paphies australis*). *P. subtriangulata* is extensively distributed around New Zealand in localised abundant populations, but mainly occurs around the North Island, and at more scattered locations in the northern South Island, Stewart Island, and the Chatham Islands.

Tuatua are ecological markers of fine, clean, fluid sands on ocean beaches with moderate wave exposure. The densest beds are found in the zone from the low intertidal to the shallow subtidal (down to about 4 m depth). The tuatua is a suspension feeder with short siphons. It is usually wedged only a few centimetres into the sand, with the straight siphonal end often characteristically exposed and discoloured by a green or brown algal film. Individuals are often dragged about the surface and redistributed by swash and backwash before actively burrowing back into the sand.

Tuatua have separate sexes (1:1 sex ratio) and reproduce by broadcast spawning, synchronously releasing eggs and sperm into the water column for external fertilisation. In north-eastern New Zealand, two main spawning periods have been documented, one between September and November, the other between February and April. Spawning events have been observed *in situ* at high water on a number of occasions, with only a small proportion of the population participating in each event. These spawning events were synchronous with pipi spawning in the same area.

Planktonic larval development takes about two to three weeks, so larvae have the potential to disperse widely if conditions allow. Larval settlement is thought to occur high in the intertidal, but spat and juveniles are highly mobile, moving around with the tidal flow before reburying themselves rapidly. Tuatua appear to migrate down the beach to occupy the lower intertidal and shallow subtidal as they grow larger. Growth appears to be rapid but variable, with tuatua reaching 40–70 mm shell length in about 3 years. Maximal length is variable among areas, ranging from about 50 to 80 mm, and the maximum age is probably about 5 or more years. Highly variable recruitment has been observed on the northwest coast

## TUATUA (TUA)

of the North Island, and this is likely to occur in other areas. As in other surf clams, natural mortality is likely to be high.

A length-weight relationship has been estimated for tuatua sampled from East Auckland, and a southern population (probably Dunedin) where weight (in g) = a (length (in mm))<sup>b</sup>, where a = 0.2 x 10<sup>-3</sup> and b = 2.927. Data source: D. Allen unpublished data. Because the samples were from one northern and one southern population, the estimated relationship may not be representative of other populations.

### 3. STOCKS AND AREAS

Little is known of the stock structure of tuatua. There have been no biological studies directly relevant to the identification of separate stocks of *P. subtriangulata* around New Zealand, although “stocks” are likely to be linked by larval dispersal. For management purposes stock boundaries are based on FMAs, with the exception of TUA 1, which was divided into TUA 1A and TUA 1B on either side of Te Arai Point because there are likely to be significant differences in the state and use of the tuatua beds between the Northland and Hauraki Gulf / Bay of Plenty areas, and the respective alignment of recreational and customary fishing interests to those management areas. The circulation patterns that maintain the separation of the surf zone habitat to form a self contained ecosystem also retain planktonic larvae of surf clams probably isolating surf clams genetically as well as ecologically.

### 4. STOCK ASSESSMENT

#### 4.1 Estimates of fishery parameters and abundance

There are no estimates of fishery parameters or abundance for any tuatua fishstock.

#### 4.2 Biomass estimates

There is no time series of biomass surveys for tuatua either in the bed in the Kaipara Harbour entrance where commercial harvesting by dredge occurs now, or anywhere else that would indicate whether tuatua populations are changing in response to past and current levels of harvesting.

#### 4.3 Yield estimates and projections

MCY has not been estimated for *P. subtriangulata*.

CAY has not been estimated for *P. subtriangulata*.

### 5. STATUS OF THE STOCKS

There are no estimates of biomass or sustainable yields of tuatua for any tuatua stock and the status of all stocks is unknown. Because natural mortality is high and recruitment is variable, the biomass of tuatua is likely to be highly variable.

- TUA - *Paphies subtriangulata*

Stock Status	
Year of Most Recent Assessment	No formal assessment conducted for any of the stocks
Assessment Runs Presented	Recruited biomass (shells $\geq$ 50mm)
Reference Points <sup>4</sup>	Target: Undefined Soft Limit: 20% $B_0$ Hard Limit: 10% $B_0$
Status in relation to Target	Unknown

Status in relation to Limits	Unknown
<b>Fishery and Stock Trends</b>	
Recent Trend in Biomass or Proxy	Unknown
Recent Trend in Fishing Mortality or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Landings are less than a quarter of the TACC and have generally been declining since 2002–03.
<b>Projections and Prognosis</b>	
Stock Projections or Prognosis	-
Probability of Current Catch or TACC causing decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
<b>Assessment Methodology</b>	
Assessment Type	-
Assessment Method	-
Main data inputs	-
Period of Assessment	- Next assessment: Unknown
Changes to Model Structure and Assumptions	-
Major Sources of Uncertainty	-

**Qualifying Comments**

Landings are thought to have been declining in recent times because of economic rather than biological reasons.

**Fishery Interactions**

There are concerns about the potential impacts of dredge fishing on complex habitats.

**6. FOR FURTHER INFORMATION**

- Beu, A G; De Rooij-Schuiling, L A (1982) Subgeneric classification of New Zealand and Australian species of *Paphies lesson* (Bivalvia: Mesodesmatidae), and names for the two species of tuatua in New Zealand. *New Zealand Journal of Zoology* 9: 211–230.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Final Research Report for Ministry of Fisheries Project REC2000/03 (Unpublished report held by Ministry for Primary Industries, Wellington).
- Boyd, R O; Reilly, J L (2002) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for the Ministry of Fisheries Project 1998/03 28 p. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Bradford, E (1998) Harvest estimates from the 1996 national marine recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington).
- Cranfield, H J; Michael, K P; Stotter, D; Doonan, I J (1994) Distribution, biomass and yield estimates of surf clams off New Zealand beaches. New Zealand Fisheries Assessment Research Document 1994/1. 27 p. (Unpublished report held by Ministry for Primary Industries, Wellington).
- Grant, C M (1994) Demographics and reproduction of the tuatua *Paphies subtriangulata*. Unpublished MSc thesis. University of Auckland, Auckland, New Zealand. 120 p.
- Grant, C M; Creese, R G (1995) The reproductive cycle of the tuatua-*Paphies subtriangulata* (Wood, 1828), in New Zealand. *Journal of Shellfish Research* 14: 287–292.
- Grant, C M; Hooker, S H; Babcock, R C; Creese, R G (1998) Synchronous spawning and reproductive incompatibility of two bivalve species: *Paphies subtriangulata* and *Paphies australis*. *Veliger* 41: 148–156.
- Haddon, M; Wear, R (1987) Biology of feeding in the New Zealand paddle crab *Ovalipes catharus* (Crustacea, Portunidae). *New Zealand Journal of Marine and Freshwater Research* 14: 55–64.
- Ministry of Fisheries Science Group (Comps.) (2006) Report from the Fishery Assessment Plenary, May 2006: stock assessments and yield estimates. 875 p. (Unpublished report held in NIWA library, Wellington.)
- Morton, J E; Miller, M C (1968) *The New Zealand sea shore*. Collins, Auckland, New Zealand. 638 p.
- Powell, A W B (1979) *New Zealand Mollusca: marine, land and freshwater shells*. Collins, Auckland, New Zealand. 500 p.
- Redfearn, P (1987) Larval shell development of the northern tuatua, *Paphies subtriangulata* (Bivalvia, Mesodesmatidae). *New Zealand Journal of Marine and Freshwater Research* 21: 65–70.