



Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2012–13: Supplementary information

New Zealand Aquatic Environment and Biodiversity Report 162

Y. Richard
E. R. Abraham

ISSN 1179-6480 (online)
ISBN 978-1-77665-110-8 (online)

November 2015



Requests for further copies should be directed to:

Publications Logistics Officer
Ministry for Primary Industries
PO Box 2526
WELLINGTON 6140

Email: brand@mpi.govt.nz
Telephone: 0800 00 83 33
Facsimile: 04-894 0300

This publication is also available on the Ministry for Primary Industries websites at:
<http://www.mpi.govt.nz/news-resources/publications.aspx>
<http://fs.fish.govt.nz> go to Document library/Research reports

© Crown Copyright - Ministry for Primary Industries

TABLE OF CONTENTS

1 OVERVIEW

i

2 SPECIES DATA

S-1

S.1	Gibson's albatross (<i>Diomedea antipodensis gibsoni</i>)	S-1
S.2	Antipodean albatross (<i>Diomedea antipodensis antipodensis</i>)	S-2
S.3	Southern royal albatross (<i>Diomedea epomophora</i>)	S-3
S.4	Northern royal albatross (<i>Diomedea sanfordi</i>)	S-4
S.5	Campbell black-browed albatross (<i>Thalassarche impavida</i>)	S-5
S.6	New Zealand white-capped albatross (<i>Thalassarche steadi</i>)	S-6
S.7	Salvin's albatross (<i>Thalassarche salvini</i>)	S-7
S.8	Chatham Island albatross (<i>Thalassarche eremita</i>)	S-8
S.9	Grey-headed albatross (<i>Thalassarche chrysostoma</i>)	S-9
S.10	Southern Buller's albatross (<i>Thalassarche bulleri bulleri</i>)	S-10
S.11	Northern Buller's albatross (<i>Thalassarche bulleri platei</i>)	S-11
S.12	Light-mantled sooty albatross (<i>Phoebastria palpebrata</i>)	S-12
S.13	Northern giant petrel (<i>Macronectes halli</i>)	S-13
S.14	Grey petrel (<i>Procellaria cinerea</i>)	S-14
S.15	Black petrel (<i>Procellaria parkinsoni</i>)	S-15
S.16	Westland petrel (<i>Procellaria westlandica</i>)	S-16
S.17	White-chinned petrel (<i>Procellaria aequinoctialis</i>)	S-17
S.18	Flesh-footed shearwater (<i>Puffinus carneipes</i>)	S-18
S.19	Wedge-tailed shearwater (<i>Puffinus pacificus</i>)	S-19
S.20	Buller's shearwater (<i>Puffinus bulleri</i>)	S-20
S.21	Sooty shearwater (<i>Puffinus griseus</i>)	S-21
S.22	Fluttering shearwater (<i>Puffinus gavia</i>)	S-22
S.23	Hutton's shearwater (<i>Puffinus huttoni</i>)	S-23
S.24	Little shearwater (<i>Puffinus assimilis</i>)	S-24
S.25	Snares Cape petrel (<i>Daption capense australe</i>)	S-25
S.26	Fairy prion (<i>Pachyptila turtur</i>)	S-26
S.27	Antarctic prion (<i>Pachyptila desolata</i>)	S-27
S.28	Broad-billed prion (<i>Pachyptila vittata</i>)	S-28
S.29	Pycroft's petrel (<i>Pterodroma pycrofti</i>)	S-29
S.30	Cook's petrel (<i>Pterodroma cookii</i>)	S-30
S.31	Chatham petrel (<i>Pterodroma axillaris</i>)	S-31
S.32	Mottled petrel (<i>Pterodroma inexpectata</i>)	S-32
S.33	White-necked petrel (<i>Pterodroma cervicalis</i>)	S-33
S.34	Kermadec petrel (<i>Pterodroma neglecta</i>)	S-34
S.35	Grey-faced petrel (<i>Pterodroma macroptera</i>)	S-35
S.36	Chatham Island taiko (<i>Pterodroma magentae</i>)	S-36
S.37	White-headed petrel (<i>Pterodroma lessonii</i>)	S-37
S.38	Soft-plumaged petrel (<i>Pterodroma mollis</i>)	S-38
S.39	Common diving petrel (<i>Pelecanoides urinatrix</i>)	S-39
S.40	South Georgia diving petrel (<i>Pelecanoides georgicus</i>)	S-40
S.41	New Zealand white-faced storm petrel (<i>Pelagodroma marina</i>)	S-41
S.42	White-bellied storm petrel (<i>Fregetta grallaria</i>)	S-42
S.43	Black-bellied storm petrel (<i>Fregetta tropica</i>)	S-43
S.44	Kermadec white-faced storm petrel (<i>Pelagodroma marina albiclunis</i>)	S-44
S.45	New Zealand storm petrel (<i>Oceanites maorianus</i>)	S-45
S.46	Yellow-eyed penguin (<i>Megadyptes antipodes</i>)	S-46
S.47	Northern little penguin (<i>Eudyptula minor</i>)	S-47
S.48	White-flipped little penguin (<i>Eudyptula minor</i>)	S-48
S.49	Southern little penguin (<i>Eudyptula minor</i>)	S-49

S.50 Chatham Island little penguin (<i>Eudyptula minor</i>)	S-50
S.51 Eastern rockhopper penguin (<i>Eudyptes chrysocome filholi</i>)	S-51
S.52 Fiordland crested penguin (<i>Eudyptes pachyrhynchus</i>)	S-52
S.53 Snares crested penguin (<i>Eudyptes robustus</i>)	S-53
S.54 Erect-crested penguin (<i>Eudyptes sclateri</i>)	S-54
S.55 Australasian gannet (<i>Morus serrator</i>)	S-55
S.56 Masked booby (<i>Sula dactylatra</i>)	S-56
S.57 Pied shag (<i>Phalacrocorax varius varius</i>)	S-57
S.58 Little black shag (<i>Phalacrocorax sulcirostris</i>)	S-58
S.59 New Zealand king shag (<i>Phalacrocorax carunculatus</i>)	S-59
S.60 Stewart Island shag (<i>Phalacrocorax chalconotus</i>)	S-60
S.61 Chatham Island shag (<i>Phalacrocorax onslowi</i>)	S-61
S.62 Bounty Island shag (<i>Phalacrocorax ranfurlyi</i>)	S-62
S.63 Auckland Island shag (<i>Phalacrocorax colensoi</i>)	S-63
S.64 Campbell Island shag (<i>Phalacrocorax campbelli</i>)	S-64
S.65 Spotted shag (<i>Phalacrocorax punctatus</i>)	S-65
S.66 Pitt Island shag (<i>Phalacrocorax featherstoni</i>)	S-66
S.67 Subantarctic skua (<i>Catharacta antarctica lonnbergi</i>)	S-67
S.68 Black-backed gull (<i>Larus dominicanus</i>)	S-68
S.69 Caspian tern (<i>Sterna caspia</i>)	S-69
S.70 Common white tern (<i>Gygis alba</i>)	S-70

S SUPPLEMENTARY REFERENCES

S-71

1. OVERVIEW

This supplementary information presents a summary of population and distributional data for the 70 species included in the risk assessment of the impact of fishing-related mortalities on seabirds breeding in the New Zealand region (Richard & Abraham 2015). For each seabird species included in the risk assessment, the demographic parameters used were the New Zealand population size, the age at first reproduction, and the survival rate. For species for which no demographic estimates were available, values from proxy species were used, as indicated with a reference to the data source. Distributional data are presented as maps of the at-sea distribution of each species, with separate maps for the non-breeding and breeding distributions. The distribution of non-breeders was derived from existing maps published by NABIS (National Aquatic Biodiversity Information System) and Birdlife International. A single distribution map was generated when the breeding season extended throughout the year. Included in the distributional maps are data of any incidental captures in commercial trawl, longline and set-net fisheries between the 2006–2007 and 2012–2013 fishing years, recorded by fisheries observers.

A detailed description of the methods used to derive the data presented here is provided in Section 2 of the risk assessment (Richard & Abraham 2015).

2. SPECIES DATA

S.1 Gibson's albatross (*Diomedea antipodensis gibsoni*)

Population (NZ)	4792 pairs [2014]	Elliott & Walker (2014), Baker & Jensz (2014)
Age at first reproduction	10 to 12 years	de L. Brooke (2004)
Survival rate	93.8 to 98.5% [1996]	Croxall & Gales (1998)
	96 to 98% [1997]	Walker & Elliott (1999)
	95.9 ± 0.6% [2004]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)




Figure S-1: Relative density of Gibson's albatross (*Diomedea antipodensis gibsoni*). The breeding season runs throughout the year, hence a single distribution map was created. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl and surface-longline (SLL) fisheries.

S.2 Antipodean albatross (*Diomedea antipodensis antipodensis*)

Population (NZ)	3320 pairs [2014]	G. Elliot (pers. comm.)
Age at first reproduction	10 to 13 years [1997]	Walker & Elliott (2002)
Survival rate	$95.7 \pm 0.7\%$ [2004]	Walker & Elliott (1999)




Figure S-2: Relative density of Antipodean albatross (*Diomedea antipodensis antipodensis*). The breeding season runs throughout the year, hence a single distribution map was created. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in surface-longline (SLL) fisheries.

S.3 Southern royal albatross (*Diomedea epomophora*)

Population (NZ)	7886 pairs [2008]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
Age at first reproduction	8.5 to 10.6 years	Robertson (1993)
Survival rate	94.9 ± 0.8% [2001]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)




Figure S-3: Relative density of southern royal albatross (*Diomedea epomophora*). The breeding season runs throughout the year, hence a single distribution map was created. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.4 Northern royal albatross (*Diomedea sanfordi*)

Population (NZ)	5832 pairs [2003]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
Age at first reproduction	8.5 to 10.6 years	Robertson (1993)
Survival rate	95.2% [1993]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
	94.6 ± 1.5% [1993]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)





Figure S-4: Relative density of northern royal albatross (*Diomedea sanfordi*). The breeding season runs throughout the year, hence a single distribution map was created. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in surface-longline (SLL) fisheries.

S.5 Campbell black-browed albatross (*Thalassarche impavida*)

Population (NZ)	21 000 pairs [1998]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
Age at first reproduction	10 (6 – 13) years [1995]	Waugh et al. (1999)
Survival rate	94.5% [1996]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-5: Relative density of Campbell black-browed albatross (*Thalassarche impavida*). The breeding season runs from August to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.6 New Zealand white-capped albatross (*Thalassarche steadi*)

Population (NZ)	96 018 (95% c.i.: 85 882 – 106 260) pairs	Richard et al. (2015)
Age at first reproduction	12 years [2011]	Southern Buller's albatross as proxy, Francis & Sagar (2012)
Survival rate	96% [2011]	Francis (2012)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-6: Relative density of New Zealand white-capped albatross (*Thalassarche steadi*). The breeding season runs from November to August. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.7 Salvin's albatross (*Thalassarche salvini*)

Population (NZ)	33 000 to 41 000 pairs [2014]	Baker et al. (2014)
Age at first reproduction	12 years [2011]	Southern Buller's albatross as proxy, Francis & Sagar (2012)
Survival rate	96.7% [2011]	Sagar et al. (2011)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-7: Relative density of Salvin's albatross (*Thalassarche salvini*). The breeding season runs from September to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.8 Chatham Island albatross (*Thalassarche eremita*)

Population (NZ)	5247 pairs [2007]	Birdlife International (2009)
Age at first reproduction	12 years [2011]	Southern Buller's albatross as proxy, Francis & Sagar (2012)
Survival rate	96.7% [2011]	Salvin's albatross as proxy, Sagar et al. (2011)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-8: Relative density of Chatham Island albatross (*Thalassarche eremita*). The breeding season runs from August to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl and bottom-longline (BLL) fisheries.

S.9 Grey-headed albatross (*Thalassarche chrysostoma*)

Population (NZ)	6600 pairs [1997]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
Age at first reproduction	7 to 13 years	Schreiber & Burger (2001)
Survival rate	$95.3 \pm 0.9\%$ ($N = 225$) [1996]	Waugh et al. (1999)

(a) Breeding distribution

(b) Non-breeding distribution




Figure S-9: Relative density of grey-headed albatross (*Thalassarche chrysostoma*). The breeding season runs from September to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.10 Southern Buller's albatross (*Thalassarche bulleri bulleri*)

Population (NZ)	13 625 pairs [2002]	Sagar & Stahl (2005)
Age at first reproduction	12 years [2011]	Francis & Sagar (2012)
Survival rate	93 to 98% [2011]	Francis & Sagar (2012)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-10: Relative density of southern Buller's albatross (*Thalassarche bulleri bulleri*). The breeding season runs from January to September. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.11 Northern Buller's albatross (*Thalassarche bulleri platei*)

Population (NZ)	16 346 pairs [2008]	Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)
Age at first reproduction	12 years [2011]	Southern Buller's albatross as proxy, Francis & Sagar (2012)
Survival rate	93.5 (93 – 98)% [2011]	Southern Buller's albatross as proxy, Francis & Sagar (2012)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-11: Relative density of northern Buller's albatross (*Thalassarche bulleri platei*). The breeding season runs from October to June. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.12 Light-mantled sooty albatross (*Phoebastria palpebrata*)

Population (NZ)	6770 to 6900 pairs	Taylor (2000a)
Age at first reproduction	12 years	de L. Brooke (2004)
Survival rate	96 to 98% [1997]	Gibson's albatross as proxy, Walker & Elliott (1999)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-12: Relative density of light-mantled sooty albatross (*Phoebastria palpebrata*). The breeding season runs from September to June. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.13 Northern giant petrel (*Macronectes halli*)

Population (NZ) 2567 pairs [1993]

Age at first reproduction 6 to 10 years

Survival rate 92.3%
88 to 93% [1981]
88% [2003]

Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)


Trivelpiece & Trivelpiece (1998)

de L. Brooke (2004)

Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)

Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-13: Relative density of northern giant petrel (*Macronectes halli*). The breeding season runs from August to February. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.14 Grey petrel (*Procellaria cinerea*)

Population (NZ) 50 000 pairs [2010]

Walker et al. (2015)


Age at first reproduction 7 years

Barbraud et al. (2009)

Survival rate 90 to 97%

White-chinned petrel as proxy, Dillingham & Fletcher (2008)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-14: Relative density of grey petrel (*Procellaria cinerea*). The breeding season runs from February to November. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.


S.15 Black petrel (*Procellaria parkinsoni*)

Population (NZ)	4627 (95% c.i.: 1972 – 9777) pairs	Richard & Abraham (2015)
Age at first reproduction	6.6 ± 0.2 years [2010]	Bell et al. (2011)
Survival rate	95%	Walker et al. (2015)

(a) Pre-egg laying (Oct–Nov)



(b) Incubation (Dec–Jan)



(c) Chick rearing (Feb–May)




Figure S-15: Relative density of black petrel (*Procellaria parkinsoni*). The breeding season runs from October to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), and bottom-longline (BLL) fisheries.

S.16 Westland petrel (*Procellaria westlandica*)

Population (NZ) 4000 pairs [2008]

Agreement on the Conservation of Albatrosses and Petrels (ACAP) (2010)


Age at first reproduction 6.5 years [2002]

Waugh et al. (2006)

Survival rate 90 to 97%

White-chinned petrel as proxy, Dillingham & Fletcher (2008)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-16: Relative density of Westland petrel (*Procellaria westlandica*). The breeding season runs from March to December. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.17 White-chinned petrel (*Procellaria aequinoctialis*)

Population (NZ)	168 725 pairs	Birdlife International (2012)
Age at first reproduction	6.5 years	Schreiber & Burger (2001)
Survival rate	90 to 97%	Dillingham & Fletcher (2008)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-17: Relative density of white-chinned petrel (*Procellaria aequinoctialis*). The breeding season runs from October to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.18 Flesh-footed shearwater (*Puffinus carneipes*)

Population (NZ)	10 000 pairs [2013]	Walker et al. (2015)
Age at first reproduction	4 to 9 years [1973]	Bradley et al. (1999)
Survival rate	94%	Walker et al. (2015)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-18: Relative density of flesh-footed shearwater (*Puffinus carneipes*). The breeding season runs from October to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.19 Wedge-tailed shearwater (*Puffinus pacificus*)

Population (NZ)	52 500 to 60 000 pairs	Taylor (2000b)
Age at first reproduction	4 years	Schreiber & Burger (2001)
Survival rate	93.1 (88.9 – 95.8)% [1999]	Hutton's shearwater as proxy, Cuthbert & Davis (2002)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-19: Relative density of wedge-tailed shearwater (*Puffinus pacificus*). The breeding season runs from October to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.20 Buller's shearwater (*Puffinus bulleri*)

Population (NZ) 200 000 pairs
 Age at first reproduction 4 to 9 years [1973]
 Survival rate 92%

de L. Brooke (2004)
 Bradley et al. (1999)
 Short-tailed shearwater as proxy, de L. Brooke (2004)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-20: Relative density of Buller's shearwater (*Puffinus bulleri*). The breeding season runs from September to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.21 Sooty shearwater (*Puffinus griseus*)

Population (NZ)	5 000 000 pairs	Taylor (2000b)
Age at first reproduction	5 to 7 years	de L. Brooke (2004)
Survival rate	86 to 97.9% [2005]	Clucas et al. (2008)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-21: Relative density of sooty shearwater (*Puffinus griseus*). The breeding season runs from October to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.22 Fluttering shearwater (*Puffinus gavia*)

Population (NZ)	20 000 to 200 000 pairs	Taylor (2000b)
Age at first reproduction	4 to 6 years	Hutton's shearwater as proxy, Waugh et al. (1999)
Survival rate	93.1 (88.9 – 95.8)% [1999]	Hutton's shearwater as proxy, Cuthbert & Davis (2002)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-22: Relative density of fluttering shearwater (*Puffinus gavia*). The breeding season runs from July to February. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in bottom-longline (BLL) and set-net (SN) fisheries.

S.23 Hutton's shearwater (*Puffinus huttoni*)

Population (NZ)	94 000 pairs	Taylor (2000a)
Age at first reproduction	4 to 6 years	Waugh et al. (1999)
Survival rate	93.1 (88.9 – 95.8)% [1999]	Cuthbert & Davis (2002)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-23: Relative density of Hutton's shearwater (*Puffinus huttoni*). The breeding season runs from September to April. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.24 Little shearwater (*Puffinus assimilis*)

Population (NZ)	100 000 to 220 000 pairs	Taylor (2000a)
Age at first reproduction	4 to 6 years	Hutton's shearwater as proxy, Waugh et al. (1999)
Survival rate	93.1 (88.9 – 95.8)% [1999]	Hutton's shearwater as proxy, Cuthbert & Davis (2002)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-24: Relative density of little shearwater (*Puffinus assimilis*). The breeding season runs from April to November. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.25 Snares Cape petrel (*Daption capense australe*)

Population (NZ)	8420 pairs	de L. Brooke (2004)
Age at first reproduction	6 years 3 to 5 years [1968]	Schreiber & Burger (2001) Beck (1969)
Survival rate	77.1 to 93.9% [1987]	Sagar et al. (1996)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-25: Relative density of Snares Cape petrel (*Daption capense australe*). The breeding season runs from November to February. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.26 Fairy prion (*Pachyptila turtur*)

Population (NZ)	more than 1 000 000 pairs	Taylor (2000b)
Age at first reproduction	4 to 5 years	Schreiber & Burger (2001)
Survival rate	84%	de L. Brooke (2004)

(a) Breeding distribution

(b) Non-breeding distribution




Figure S-26: Relative density of fairy prion (*Pachyptila turtur*). The breeding season runs from March to January. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.27 Antarctic prion (*Pachyptila desolata*)

Population (NZ)	100 000 to 1 000 000 pairs	Taylor (2000b)
Age at first reproduction	5 to 6 years	de L. Brooke (2004)
Survival rate	84%	Fairy prion as proxy, de L. Brooke (2004)

(a) Breeding distribution

(b) Non-breeding distribution




Figure S-27: Relative density of Antarctic prion (*Pachyptila desolata*). The breeding season runs from November to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.28 Broad-billed prion (*Pachyptila vittata*)

Population (NZ) 1 000 000 pairs

de L. Brooke (2004)

Age at first reproduction 4 to 5 years

Fairy prion as proxy, Schreiber & Burger (2001)

Survival rate 84%

Fairy prion as proxy, de L. Brooke (2004)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-28: Relative density of broad-billed prion (*Pachyptila vittata*). The breeding season runs from February to January. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.29 Pycroft's petrel (*Pterodroma pycrofti*)

Population (NZ)	2000 to 3000 pairs [1998]	Taylor (2000a)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-29: Relative density of Pycroft's petrel (*Pterodroma pycrofti*). The breeding season runs from October to April. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.30 Cook's petrel (*Pterodroma cookii*)

Population (NZ) 50 000 to 60 000 pairs

Taylor (2000a)


Age at first reproduction 6 to 7 years

Grey-faced petrel as proxy, Schreiber & Burger (2001)

Survival rate 94%

Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-30: Relative density of Cook's petrel (*Pterodroma cookii*). The breeding season runs from September to April. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.31 Chatham petrel (*Pterodroma axillaris*)

Population (NZ) 250 pairs [2009]

Age at first reproduction 6 to 7 years


Survival rate 94%

Birdlife International (2012)

Grey-faced petrel as proxy, Schreiber & Burger (2001)

Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-31: Relative density of Chatham petrel (*Pterodroma axillaris*). The breeding season runs from November to June. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.32 Mottled petrel (*Pterodroma inexpectata*)

Population (NZ)	300 000 to 400 000 pairs [1999]	Taylor (2000b)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-32: Relative density of mottled petrel (*Pterodroma inexpectata*). The breeding season runs from October to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.33 White-necked petrel (*Pterodroma cervicalis*)

Population (NZ)	50 000 pairs [1988]	Taylor (2000a)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-33: Relative density of white-necked petrel (*Pterodroma cervicalis*). The breeding season runs from October to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.34 Kermadec petrel (*Pterodroma neglecta*)

Population (NZ)	5000 to 7000 pairs	Taylor (2000b)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)





Figure S-34: Relative density of Kermadec petrel (*Pterodroma neglecta*). No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.35 Grey-faced petrel (*Pterodroma macroptera*)

Population (NZ)	200 000 to 300 000 pairs	Taylor (2000b)
Age at first reproduction	6 to 7 years	Schreiber & Burger (2001)
Survival rate	94%	Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-35: Relative density of grey-faced petrel (*Pterodroma macroptera*). The breeding season runs from March to January. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in surface-longline (SLL) and bottom-longline (BLL) fisheries.

S.36 Chatham Island taiko (*Pterodroma magentae*)

Population (NZ)	17 pairs [2010]	Birdlife International (2012)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-36: Relative density of Chatham Island taiko (*Pterodroma magentae*). The breeding season runs from September to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.37 White-headed petrel (*Pterodroma lessonii*)

Population (NZ) 200 000 pairs

de L. Brooke (2004)


Age at first reproduction 5.5 years

Schreiber & Burger (2001)

Survival rate 94%

Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-37: Relative density of white-headed petrel (*Pterodroma lessonii*). The breeding season runs from November to June. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.38 Soft-plumaged petrel (*Pterodroma mollis*)

Population (NZ)	1000 to 9999 pairs	Taylor (2000b)
Age at first reproduction	6 to 7 years	Grey-faced petrel as proxy, Schreiber & Burger (2001)
Survival rate	94%	Grey-faced petrel as proxy, Marchant & Higgins (1990)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-38: Relative density of soft-plumaged petrel (*Pterodroma mollis*). The breeding season runs from August to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.39 Common diving petrel (*Pelecanoides urinatrix*)

Population (NZ)	300 000 to 2 150 000 pairs	Taylor (2000b)
Age at first reproduction	2 to 3 years	de L. Brooke (2004)
Survival rate	75 to 87%	Schreiber & Burger (2001)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-39: Relative density of common diving petrel (*Pelecanoides urinatrix*). The breeding season runs from September to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.40 South Georgia diving petrel (*Pelecanoides georgicus*)

Population (NZ)	64 pairs [1998]	Taylor (2000b)
Age at first reproduction	2 to 3 years	Common diving petrel as proxy, de L. Brooke (2004)
Survival rate	75 to 87%	Common diving petrel as proxy, Schreiber & Burger (2001)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-40: Relative density of South Georgia diving petrel (*Pelecanoides georgicus*). The breeding season runs from September to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.41 New Zealand white-faced storm petrel (*Pelagodroma marina*)

Population (NZ)	more than 1 000 000 pairs	Taylor (2000b)
Age at first reproduction	4 to 5 years more than 3 years	Several species as proxy, Croxall (1987) de L. Brooke (2004)
Survival rate	90%	Several species as proxy, Croxall (1987)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-41: Relative density of New Zealand white-faced storm petrel (*Pelagodroma marina*). The breeding season runs from September to April. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.42 White-bellied storm petrel (*Fregetta grallaria*)

Population (NZ)	1000 pairs	Taylor (2000a)
Age at first reproduction	4 to 5 years	Several species as proxy, Croxall (1987)
Survival rate	90%	Several species as proxy, Croxall (1987)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-42: Relative density of white-bellied storm petrel (*Fregetta grallaria*). The breeding season runs from April to August. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.43 Black-bellied storm petrel (*Fregetta tropica*)

Population (NZ)	50 000 to 100 000 pairs	Taylor (2000b)
Age at first reproduction	4 to 5 years	Several species as proxy, Croxall (1987)
Survival rate	90%	Several species as proxy, Croxall (1987)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-43: Relative density of black-bellied storm petrel (*Fregetta tropica*). The breeding season runs from October to May. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl fisheries.

S.44 Kermadec white-faced storm petrel (*Pelagodroma marina albiclunis*)

Population (NZ)	fewer than 100 pairs	Taylor (2000a)
Age at first reproduction	4 to 5 years more than 3 years	Several species as proxy, Croxall (1987) de L. Brooke (2004)
Survival rate	90%	Several species as proxy, Croxall (1987)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-44: Relative density of Kermadec white-faced storm petrel (*Pelagodroma marina albiclunis*). The breeding season runs from June to December. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.45 New Zealand storm petrel (*Oceanites maorianus*)

Population (NZ)	20 to 1000 pairs	
Age at first reproduction	4 to 5 years	Several species as proxy, Croxall (1987)
Survival rate	90%	Several species as proxy, Croxall (1987)




Figure S-45: Relative density of New Zealand storm petrel (*Oceanites maorianus*). No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BL), and set-net (SN) fisheries.

S.46 Yellow-eyed penguin (*Megadyptes antipodes*)

Population (NZ)	1700 to 2420 pairs	Taylor (2000a)
Age at first reproduction	2 to 3 years	Schreiber & Burger (2001)
Survival rate	87%	Schreiber & Burger (2001)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-46: Relative density of yellow-eyed penguin (*Megadyptes antipodes*). The breeding season runs from August to April. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in set-net (SN) fisheries.

S.47 Northern little penguin (*Eudyptula minor*)

Population (NZ)	5000 to 10 000 pairs [1984]	Taylor (2000b)
Age at first reproduction	2 to 3 years	Schreiber & Burger (2001)
Survival rate	83%	Sidhu et al. (2007)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-47: Relative density of northern little penguin (*Eudyptula minor*). The breeding season runs from July to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.48 White-flipped little penguin (*Eudyptula minor*)

Population (NZ)	2200 pairs [1998]	Taylor (2000a)
Age at first reproduction	2 to 3 years	Schreiber & Burger (2001)
Survival rate	83%	Sidhu et al. (2007)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-48: Relative density of white-flipped little penguin (*Eudyptula minor*). The breeding season runs from July to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.49 Southern little penguin (*Eudyptula minor*)

Population (NZ)	5000 to 10 000 pairs [1984]	Taylor (2000b)
Age at first reproduction	2 to 3 years	Schreiber & Burger (2001)
Survival rate	83%	Sidhu et al. (2007)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-49: Relative density of southern little penguin (*Eudyptula minor*). The breeding season runs from July to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.50 Chatham Island little penguin (*Eudyptula minor*)

Population (NZ)	5000 to 10 000 pairs [1984]	Taylor (2000b)
Age at first reproduction	2 to 3 years	Schreiber & Burger (2001)
Survival rate	83%	Sidhu et al. (2007)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-50: Relative density of Chatham Island little penguin (*Eudyptula minor*). The breeding season runs from July to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.51 Eastern rockhopper penguin (*Eudyptes chrysocome filholi*)

Population (NZ)	38 961 to 58 500 pairs	Taylor (2000a)
Age at first reproduction	4.7 years	Moseley's rockhopper penguin as proxy, Guinard et al. (1998)
Survival rate	$84 \pm 1.1\%$ [1995]	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-51: Relative density of eastern rockhopper penguin (*Eudyptes chrysocome filholi*). The breeding season runs from October to May. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.52 Fiordland crested penguin (*Eudyptes pachyrhynchus*)

Population (NZ)	3000 pairs	Roots (2006)
Age at first reproduction	3 to 4 years 5 to 6 years	Schreiber & Burger (2001) Marchant & Higgins (1990)
Survival rate	84 ± 1.1% [1995]	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-52: Relative density of Fiordland crested penguin (*Eudyptes pachyrhynchus*). The breeding season runs from July to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in set-net (SN) fisheries.

S.53 Snares crested penguin (*Eudyptes robustus*)

Population (NZ)	30 000 pairs	Roots (2006)
Age at first reproduction	5 to 6 years	Roots (2006)
Survival rate	$84 \pm 1.1\%$ [1995]	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-53: Relative density of Snares crested penguin (*Eudyptes robustus*). The breeding season runs from September to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.54 Erect-crested penguin (*Eudyptes sclateri*)

Population (NZ)	81 000 (77 000 – 85 000) pairs	Taylor (2000a)
Age at first reproduction	5 to 6 years	Fiordland crested penguin as proxy, Roots (2006)
Survival rate	84 ± 1.1% [1995]	Northern rockhopper penguin as proxy, Guinard et al. (1998)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-54: Relative density of erect-crested penguin (*Eudyptes sclateri*). The breeding season runs from September to March. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.55 Australasian gannet (*Morus serrator*)

Population (NZ) 46 004 pairs [1981]

Age at first reproduction 3 to 7 years


Survival rate 94%

Wodzicki et al. (1984)

Schreiber & Burger (2001)

Northern gannet as proxy, Schreiber & Burger (2001)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-55: Relative density of Australasian gannet (*Morus serrator*). The breeding season runs from August to March. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.56 Masked booby (*Sula dactylatra*)

Population (NZ)	240 pairs	Veitch et al. (2004)
Age at first reproduction	2 to 4 years	Schreiber & Burger (2001)
Survival rate	85% [1979]	Harris (1979)




Figure S-56: Relative density of masked booby (*Sula dactylatra*). No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.57 Pied shag (*Phalacrocorax varius varius*)

Population (NZ)	6400 pairs [2013]	Bell (2013)
Age at first reproduction	more than 2 years	Schreiber & Burger (2001)
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)




Figure S-57: Relative density of pied shag (*Phalacrocorax varius varius*). Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in set-net (SN) fisheries.

S.58 Little black shag (*Phalacrocorax sulcirostris*)

Population (NZ)	1500 pairs	Walker et al. (2015)
Age at first reproduction	2 years	Pied shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-58: Relative density of little black shag (*Phalacrocorax sulcirostris*). The breeding season runs from October to December. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.59 New Zealand king shag (*Phalacrocorax carunculatus*)

Population (NZ)	102 to 126 pairs [2002]	Birdlife International (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-59: Relative density of New Zealand king shag (*Phalacrocorax carunculatus*). The breeding season runs from March to October. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.60 Stewart Island shag (*Phalacrocorax chalconotus*)

Population (NZ)	2075 to 2482 pairs [2011]	Lalas & Perriman (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-60: Relative density of Stewart Island shag (*Phalacrocorax chalconotus*). The breeding season runs from August to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in set-net (SN) fisheries.

S.61 Chatham Island shag (*Phalacrocorax onslowi*)

Population (NZ)	357 pairs [2011]	Birdlife International (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-61: Relative density of Chatham Island shag (*Phalacrocorax onslowi*). The breeding season runs from September to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.62 Bounty Island shag (*Phalacrocorax ranfurlyi*)

Population (NZ)	120 pairs [2005]	Birdlife International (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-62: Relative density of Bounty Island shag (*Phalacrocorax ranfurlyi*). The breeding season runs from October to December. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.63 Auckland Island shag (*Phalacrocorax colensoi*)

Population (NZ)	more than 1366 pairs [2011]	Birdlife International (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-63: Relative density of Auckland Island shag (*Phalacrocorax colensoi*). The breeding season runs from November to March. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.64 Campbell Island shag (*Phalacrocorax campbelli*)

Population (NZ)	2000 pairs [1975]	Birdlife International (2012)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-64: Relative density of Campbell Island shag (*Phalacrocorax campbelli*). The breeding season runs from November to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.65 Spotted shag (*Phalacrocorax punctatus*)

Population (NZ)	10 000 to 30 000 pairs	Taylor (2000b)
Age at first reproduction	2 years	Schreiber & Burger (2001)
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)





Figure S-65: Relative density of spotted shag (*Phalacrocorax punctatus*). Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl and set-net (SN) fisheries.

S.66 Pitt Island shag (*Phalacrocorax featherstoni*)

Population (NZ)	669 pairs [1997]	Taylor (2000a)
Age at first reproduction	more than 3 years	Black shag as proxy
Survival rate	87.8 (85.9 – 89.7)%	European shag as proxy, Harris et al. (1994)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-66: Relative density of Pitt Island shag (*Phalacrocorax featherstoni*). The breeding season runs from September to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.67 Subantarctic skua (*Catharacta antarctica lonnbergi*)

Population (NZ)	450 to 470 pairs	Wilson (2006)
Age at first reproduction	8.03 ± 0.21 years ($N = 96$) [1996]	Young (1998)
Survival rate	93.8 (91 – 97)% [1965]	Wood (1971)

(a) Breeding distribution

(b) Non-breeding distribution





Figure S-67: Relative density of subantarctic skua (*Catharacta antarctica lonnbergi*). The breeding season runs from September to February. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.68 Black-backed gull (*Larus dominicanus*)

Population (NZ)	more than 1 000 000 pairs	Taylor (2000b)
Age at first reproduction	4 years	Schreiber & Burger (2001)
Survival rate	81%	Schreiber & Burger (2001)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-68: Relative density of black-backed gull (*Larus dominicanus*). The breeding season runs from September to March. Also shown are incidental captures recorded by observers between the 2006–07 and 2012–13 fishing years in trawl and bottom-longline (BLL) fisheries.

S.69 Caspian tern (*Sterna caspia*)

Population (NZ)	1000 pairs [1992]
Age at first reproduction	2 to 4 years
Survival rate	87 to 91% 89% [1980]

Taylor (2000b)
Schreiber & Burger (2001)
Schreiber & Burger (2001)
Gill & Mewaldt (1983)

(a) Breeding distribution



(b) Non-breeding distribution





Figure S-69: Relative density of Caspian tern (*Sterna caspia*). The breeding season runs from September to January. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S.70 Common white tern (*Gygis alba*)

Population (NZ)	60 to 100 pairs	Taylor (2000b)
Age at first reproduction	3 to 5 years	Schreiber & Burger (2001)
Survival rate	78 to 83%	Bridled tern as proxy, Schreiber & Burger (2001)

(a) Breeding distribution



(b) Non-breeding distribution




Figure S-70: Relative density of common white tern (*Gygis alba*). The breeding season runs from September to April. No incidental capture was recorded by observers between the 2006–07 and 2012–13 fishing years in trawl, surface-longline (SLL), bottom-longline (BLL), and set-net (SN) fisheries.

S. SUPPLEMENTARY REFERENCES

- Agreement on the Conservation of Albatrosses and Petrels (ACAP). (2010). ACAP species assessment. Retrieved 7 May 2010, from <http://www.acap.aq>
- Baker, G.B.; Jensz, K. (2014). Gibson's albatross at Disappointment island—analysis of aerial photographs. Unpublished report prepared for the Department of Conservation, Wellington, New Zealand. Retrieved from <http://www.doc.govt.nz/Documents/conservation/marine-and-coastal/marine-conservation-services/reports/gibsons-albatross-disappointment-island-2014-draft-report.pdf>
- Baker, G.B.; Jensz, K.; Sagar, P. (2014). 2013 Aerial survey of Salvin's albatross at the Bounty Islands. Unpublished report prepared for the Department of Conservation, Wellington, New Zealand. Retrieved from <http://www.doc.govt.nz/Documents/conservation/marine-and-coastal/marine-conservation-services/meetings/pop2012-06-salvins-albatross-aerial-population-estimate.pdf>
- Barbraud, C.; Delord, K.; Marteau, C.; Weimerskirch, H. (2009). Estimates of population size of white-chinned petrels and grey petrels at Kerguelen Islands and sensitivity to fisheries. *Animal Conservation* 12: 258–265.
- Beck, J.R. (1969). Food, moult and age of first breeding in the Cape pigeon, *Daption capensis* (Linnaeus). *British Antarctic Survey Bulletin* 21: 33–44.
- Bell, E.A.; Sim, J.L.; Scofield, P.; Francis, R.I.C.C. (2011). Population parameters of the black petrels (*Procellaria parkinsoni*) on Great Barrier Island (Aotea Island), 2009/10. Unpublished report prepared for the Department of Conservation. Retrieved 15 January 2012, from <http://www.doc.govt.nz/publications/conservation/marine-and-coastal/conservation-services-programme/csp-reports/population-parameters-of-black-petrels-on-great-barrier-aotea-island-2009-10/>
- Bell, M. (2013). Pied shag: A national population review. Final Research Report for the Department of Conservation. Project POP2011-07, (Unpublished report held by the Department of Conservation, Wellington.) Retrieved from <http://www.doc.govt.nz/Documents/conservation/marine-and-coastal/marine-conservation-services/reports/pied-shag-population-review-final-report.pdf>
- Birdlife International. (2009). Birdlife International. Species factsheets. <http://www.birdlife.org>. Retrieved 21 May 2010, from <http://www.birdlife.org>
- Birdlife International. (2012). Birdlife International. Species factsheets. <http://www.birdlife.org>. Retrieved 10 May 2012, from <http://www.birdlife.org>
- Bradley, J.S.; Gunn, B.M.; Skira, I.J.; Meathrel, C.E.; Wooller, R.D. (1999). Age-dependent prospecting and recruitment to a breeding colony of short-tailed shearwaters *Puffinus tenuirostris*. *Ibis* 141: 277–285.
- Clucas, R.J.; Fletcher, D.J.; Moller, H. (2008). Estimates of adult survival rate for three colonies of sooty shearwater (*Puffinus griseus*) in New Zealand. *Emu* 108: 237–250.
- Croxall, J.P. (1987). The status and conservation of Antarctic seals and seabirds: a review. *Environment International* 13: 55–70.
- Croxall, J.P.; Gales, R. (1998). An assessment of the conservation status of albatrosses. In: Robertson, G.; Gales, R. (Eds.), *The albatross: biology and conservation*, Surrey Beatty & Sons, Chipping Norton, Australia, pp. 46–65.
- Cuthbert, R.; Davis, L.S. (2002). Adult survival and productivity of Hutton's shearwaters. *Ibis* 144(3): 423–432.
- de L. Brooke, M. (2004). *Albatrosses and petrels across the world*. Oxford University Press. 499 p.
- Dillingham, P.W.; Fletcher, D. (2008). Estimating the ability of birds to sustain additional human-caused mortalities using a simple decision rule and allometric relationships. *Biological Conservation* 141: 1783–1792.
- Elliott, G.; Walker, K. (2014). Gibson's wandering albatross research, Adams Island 2014. Unpublished report prepared for the Department of Conservation, Wellington, New Zealand. Retrieved from <http://www.doc.govt.nz/Documents/conservation/marine-and-coastal/marine-conservation-services/meetings/pop2013-03-gibsons-albatross-population-study-draft-final-report.pdf>
- Francis, R.I.C.C. (2012). Fisheries risks to the population viability of white-capped albatross (*Thalasarche steadi*). *New Zealand Aquatic Environment and Biodiversity Report No. 104*. 24 p.

- Francis, R.I.C.C.; Sagar, P.M. (2012). Modelling the effect of fishing on southern Buller's albatross using a 60-year dataset. *New Zealand Journal of Zoology* 39(1): 3–17.
- Gill, R.E.; Mewaldt, L.R. (1983). Pacific coast caspian terns: dynamics of an expanding population. *The Auk* 100(2): 369–381.
- Guinard, E.; Weimerskirch, H.; Jouventin, P. (1998). Population changes and demography of the northern rockhopper penguin on Amsterdam and Saint Paul islands. *Colonial waterbirds* 21(2): 222–228.
- Harris, M.P. (1979). Survival and ages of first breeding of Galapagos seabirds. *Bird-Banding* 50(1): 56–61. doi:10.2307/4512409
- Harris, M.P.; Buckland, S.T.; Russell, S.M.; Wanless, S. (1994). Year-and age-related variation in the survival of adult European shags over a 24-year period. *Condor* 96(3): 600–605.
- Lalas, C.; Perriman, L. (2012). Distribution and abundance of Stewart Island shags (*Leucocarbo chalconotus*). (Unpublished report held by the Department of Conservation, Wellington.)
- Marchant, S.; Higgins, P.J. (1990). Handbook of Australian, New Zealand and Antarctic birds. Volume 1, part A. Oxford University Press, Melbourne, Australia. 735 p.
- Richard, Y.; Abraham, E.R. (2015). Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 162*. 89 p.
- Richard, Y.; Perriman, L.; Lalas, C.; Abraham, E.R. (2015). Demographic rates of northern royal albatross at Taiaroa Head, New Zealand. *PeerJ* 3: e906. doi:10.7717/peerj.906
- Robertson, C.J.R. (1993). Survival and longevity of the northern royal albatross *Diomedea epomophora sanfordi* at Taiaroa Head 1937–93. *Emu* 93(4): 269–276.
- Roots, C. (2006). Flightless birds. Greenwood Press, Connecticut. 248 p.
- Sagar, P.M.; Stahl, J.C. (2005). Increases in the numbers of breeding pairs in two populations of Buller's albatross (*Thalassarche bulleri bulleri*). *Emu* 105(1): 49–55.
- Sagar, P.; Carroll, J.; Charteris, M.; Thompson, D.; Scofield, P. (2011). Population assessment of Salvin's albatrosses at the Snares Western Chain, 25 September – 14 October 2010. Final Research Report for research project PRO200601-E (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Sagar, P.; Tennyson, A.; Miskelly, C. (1996). Breeding and survival of Snares Cape pigeons *Daption capense australe* at The Snares, New Zealand. *Notornis* 43: 197–207.
- Schreiber, E.A.; Burger, J. (2001). Biology of marine birds. CRC Press, Boca Raton.
- Sidhu, L.A.; Catchpole, E.A.; Dann, P.; Shaffer, T.L. (2007). Mark-recapture-recovery modeling and age-related survival in little penguins (*Eudyptula minor*). *The Auk* 124(3): 815–827.
- Taylor, G.A. (2000a). Action plan for seabird conservation in New Zealand. Part A: Threatened seabirds. *Threatened Species Occasional Publication No. 16*. 234 p.
- Taylor, G.A. (2000b). Action plan for seabird conservation in New Zealand. Part B: Non-threatened seabirds. *Threatened Species Occasional Publication No. 17*. 201 p.
- Trivelpiece, S.G.; Trivelpiece, W.Z. (1998). Post-fledging dispersal of southern giant petrels *Macronectes giganteus* banded at Admiralty Bay, King George Island, Antarctica. *Marine Ornithology* 26: 63–68.
- Veitch, C.R.; Miskelly, C.M.; Harper, G.A.; Taylor, G.A.; Tennyson, A.J.D. (2004). Birds of the Kermadec Islands, south-west Pacific. *Notornis* 51(2): 61–90.
- Walker, K.; Elliott, G. (1999). Population changes and biology of the wandering albatross *Diomedea exulans gibsoni* at the Auckland Islands. *Emu* 99: 239–247.
- Walker, K.; Elliott, G. (2002). Monitoring Antipodean and Gibson's wandering albatross, 1996/97. *Department of Conservation Science Internal Series* 75. 14 p.
- Walker, N.; Smith, N.; Sharp, B.; Cryer, M. (2015). A qualitative review of New Zealand's 2013 level two risk assessment for seabirds. *New Zealand Fisheries Science Review* 2015/1: 53 p. Retrieved 10 November 2015, from <https://fs.fish.govt.nz/Page.aspx?pk=113&dk=23943>.
- Waugh, S.M.; Doherty, P.F.; Freeman, A.N.D.; Adams, L.; Woods, G.C.; Bartle, J.A.; Hedley, G.K. (2006). Demography of Westland petrels (*procellaria westlandica*), 1995–2003. *Emu* 106: 219–226.

- Waugh, S.M.; Weimerskirch, H.; Moore, P.J.; Sagar, P.M. (1999). Population dynamics of black-browed and grey-headed albatrosses *Diomedea melanophrys* and *D. chrysostoma* at Campbell Island, New Zealand, 1942–96. *Ibis* 141: 216–225.
- Wilson, K.-J. (2006). The state of New Zealand's birds. Special report seabirds. Ornithological Society of New Zealand, Nelson.
- Wodzicki, K.; Robertson, C.; Thompson, H.; Alderton, C. (1984). The distribution and numbers of gannets in New Zealand. *Notornis* 31: 232–261.
- Wood, R.C. (1971). Population dynamics of breeding south polar skuas of unknown age. *The Auk* 88(4): 805–814.
- Young, E.C. (1998). Dispersal from natal territories and the origin of cooperatively polyandrous breeding groups in the brown skua. *Condor* 100: 335–342.