


## BLUE MACKEREL (EMA)

*(Scomber australasicus)*

Tawatawa



## 1. FISHERY SUMMARY

Blue mackerel were introduced into the QMS on 1 October 2002. Since then allowances, TACCs and TACs (Table 1) have not changed.

**Table 1: Recreational and Customary non-commercial allowances, TACCs and TACs for blue mackerel by Fishstock.**

Fishstock	Recreational Allowance	Customary Non-Commercial Allowance	TACC	TAC
EMA 1	40	20	7 630	7 690
EMA 2	5	2	180	187
EMA 3	1	1	390	392
EMA 7	1	1	3 350	3 352
EMA 10	0	0	0	0
Total	47	24	11 550	11 621

### 1.1 Commercial fisheries

Blue mackerel are taken by a variety of methods but for most of these methods the catches are very low. The largest and most consistent catches have been from the target purse seine fishery in EMA 1, 2 and 7, and as non-target catch in the jack mackerel mid-water trawl fishery in EMA 7. Most catch is taken north of latitude 43° S (Kaikoura). Historical estimated and recent reported blue mackerel landings and TACCs are shown in Tables 2 and 3, while Figure 1 shows the historical landings and TACC values for these three main stocks. Since 1983–84 the catch of blue mackerel in New Zealand waters has grown substantially (Table 3), primarily in the purse seine fishery in EMA 1, and have averaged about 10 000 t annually since 1990–91.

Most blue mackerel purse seine catch comes from the Bay of Plenty (BoP) and East Northland, where it is primarily taken between July and December. Purse seine fishing effort on blue mackerel has been strongly influenced by the availability and market value of other pelagic species, particularly skipjack tuna and kahawai, with effort increasing as limits have been placed on the purse seine catch of kahawai. The purse seine fishery has accounted for more than 97% of annual EMA 1 landings since at least 1990, and about 90% of this was targeted (Ballara 2016).

Total blue mackerel catches peaked in 1991–92 at more than 15 000 t, of which 60–70% was taken by purse seine. More recently, commercial landings of over 12 500 t were taken in 1998–99 (13 500 t), 2000–01 (13 100 t) and 2004–05 (12 750 t), with the highest landings recorded in EMA 1 and EMA 7. EMA 1 landings exceeded the TACC in 2004–05, 2006–07, 2009–10, 2011–12 and 2014–15. The 2004–05, 2005–06, and 2008–09 EMA 7 landings also exceeded the TACC. The EMA 7 landings in 2014–15, however, were the lowest since the mid-1980s. Landings from EMA 2 and EMA 3 have been well

(EMA)

below the TACCs since the early to mid-1990s; they are primarily a bycatch of purse seine (EMA 2) and trawl (EMA 3) fisheries.

The blue mackerel catch from EMA 7 is now principally non-target catch from the jack mackerel mid-water trawl fishery. Highest catches are taken during June, July and October in areas 034 and 035 on the WCSI and areas 041 and 801 further north (WCNI). Fishing has shifted from south to north in the last decade. Since the late 1990s, a fleet of Ukrainian vessels has taken most of the catch in the JMA 7 target fishery and these vessels have taken the EMA as bycatch. Since 2004, 0–11% of the EMA 7 catch has been taken annually by purse seine, down from an average of about 25% between 1991 and 2003.

A number of factors have been identified that can influence landing volumes in the blue mackerel fisheries. In the purse seine fishery, blue mackerel has become the second most preferred species because of decreased TACCs on kahawai. Skipjack tuna is the preferred species and blue mackerel will not be targeted once the skipjack season has begun in late-spring, early summer. Thus, early arrival of skipjack can result in reduced volumes of blue mackerel being landed.

Management of company quota is complicated by the relative timing of the fishing season and the fishing year and this, along with the timing of the main market, may influence whether the blue mackerel TACC can all be taken in a particular year. The fishing season usually begins in about July–August, runs through to the end-beginning of subsequent fishing years, and finishes in about November. The main market for purse seined blue mackerel takes up to 80% of the catch and requires premium fish to be available from early spring. To meet the demands of this market and to minimise the costs of storing fish from the previous season, fishing companies must carry over some proportion of their quota for a given year until fish become available the following season. If availability is delayed until after October 1, only 10% of the total quota can then be carried over into the new fishing year.

Because blue mackerel is taken principally as bycatch in the jack mackerel TCEPR target fishery in JMA 7, factors influencing the targeting of jack mackerel also affect blue mackerel landings. Other bycatch species taken in this fishery include barracouta, gurnard, John dory, kingfish, and snapper, and, although non-availability of ACE is unlikely to be constraining in the first three of these, the same is not true of kingfish and snapper. Fishing company spokespersons have stated that known hotspots of snapper are avoided. Other factors in this fishery include strategies to avoid the catch of marine mammals, and a code of practice operates in which gear is not deployed between 2 a.m. and 4 a.m. It is unknown whether this affects total landing volumes.

**Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.**

Year	EMA 1	EMA 2	EMA 3	EMA 7	Year	EMA 1	EMA 2	EMA 3	EMA 7
1931-32	0	0	0	0	1957	0	0	0	0
1932-33	0	0	0	0	1958	0	0	0	0
1933-34	0	0	0	0	1959	0	0	0	0
1934-35	0	0	0	0	1960	0	0	0	0
1935-36	0	0	0	0	1961	0	0	0	0
1936-37	0	0	0	0	1962	0	0	0	0
1937-38	0	0	0	0	1963	0	0	0	0
1938-39	0	0	0	0	1964	0	0	0	0
1939-40	0	0	0	0	1965	0	0	0	0
1940-41	0	0	0	0	1966	0	0	0	0
1941-42	0	0	0	0	1967	0	0	0	0
1942-43	0	0	0	0	1968	0	0	0	0
1943-44	0	0	0	0	1969	0	0	0	0
1944	0	0	0	0	1970	0	0	0	0
1945	0	0	0	0	1971	0	0	0	0
1946	0	0	0	0	1972	0	0	0	0
1947	0	0	0	0	1973	0	0	0	0
1948	0	0	0	0	1974	38	8	0	6
1949	0	0	0	0	1975	10	0	0	2
1950	0	0	0	0	1976	50	49	0	0
1951	0	0	0	0	1977	34	135	0	0
1952	0	0	0	0	1978	14	55	0	128
1953	0	0	0	0	1979	185	31	0	317
1954	0	0	0	0	1980	752	32	0	407
1955	0	0	0	0	1981	459	49	0	1363
1956	0	0	0	0	1982	305	0	0	791

**Notes**

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.

2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.


3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

## BLUE MACKEREL (EMA)

**Table 3: Reported landings (t) of blue mackerel by QMA, and where area was unspecified (Unsp.), from 1983–84 to 2015–16. CELR data from 1986–87 to 2000–01. MHR data from 2001–02 to present.**

Fishing year	QMA					Unsp	Total
	1	2	3	7	10#		
1983–84*	480	259	44	245	0	1	1 028
1984–85*	565	222	18	865	0	73	1 743
1985–86*	618	30	190	408	0	51	1 296
1986–87	1 431	7	424	489	0	49	2 399
1987–88	2 641	168	864	1 896	0	58	5 625
1988–89	1 580	< 1	1 141	1 021	0	469	4 211
1989–90	2 158	76	518	1 492	0	< 1	4 245
1990–91	5 783	94	478	3 004	0	0	9 358
1991–92	10 926	530	65	3 607	0	0	15 128
1992–93	10 684	309	133	1 880	0	0	13 006
1993–94	4 178	218	223	1 402	5	0	6 025
1994–95	6 734	94	154	1 804	10	149	8 944
1995–96	4 170	119	173	1 218	0	1	5 680
1996–97	6 754	78	340	2 537	0	< 1	9 708
1997–98	4 595	122	78	2 310	0	< 1	7 104
1998–99	4 505	186	62	8 756	0	4	13 519
1999–00	3 602	73	3	3 169	0	0	6 847
2000–01	9 738	113	6	3 278	0	< 1	13 134
2001–02	6 368	177	49	5 101	0	0	11 694
2002–03	7 609	115	88	3 563	0	0	11 375
2003–04	6 523	149	1	2 701	0	0	9 373
2004–05	7 920	9	< 1	4 817	0	0	12 746
2005–06	6 713	13	133	3 784	0	0	10 643
2006–07	7 815	133	42	2 698	0	0	10 688
2007–08	5 926	6	122	2 929	0	0	8 982
2008–09	3 147	2	88	3 503	0	0	6 740
2009–10	8 539	3	14	3 260	0	0	11 816
2010–11	6 630	2	9	1 996	0	0	8 638
2011–12	8 080	2	28	2 707	0	0	10 817
2012–13	7 213	3	100	2 401	0	0	9 716
2013–14	6 860	4	29	1 200	0	0	8 092
2014–15	8 134	16	87	892	0	0	9 129
2015–16	7 226	18	27	761	0	0	8 033

\* FSU data, # Landings reported from QMA 10 are probably attributable to Statistical Area 010 in the Bay of Plenty (i.e., QMA 1).



**Figure 1: Reported commercial landings and TACC for the three main EMA stocks. EMA 1 (Auckland East)**  
(continued on next page).

(EMA)




Figure 1: Reported commercial landings and TACC for the three main EMA stocks. From top: EMA 2 (Central East), and EMA 7 (Challenger to Auckland West).

## 1.2 Recreational fisheries

Blue mackerel does not rate highly as a recreational target species although it is popular as bait.

There is some uncertainty with all recreational harvest estimates for blue mackerel and there is some confusion between blue and jack mackerels in the recreational data. The harvest estimates from the diary surveys should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and, c) the 2000 and 2001 estimates are implausibly high for many important fisheries.

Recreational catch in the northern region (EMA 1) was estimated at 114 000 fish by a diary survey in 1993–94 (Bradford 1996), 47 000 fish in a national recreational survey in 1996 (Bradford 1998), 84 000 fish (CV 42%) in the 2000 survey (Boyd & Reilly 2005) and 58 000 fish (CV 27%) in the 2001 survey (Boyd et al 2004). The surveys suggest a harvest of 35–90 t per year for EMA 1, insignificant in the context of the commercial catch. Estimates from other areas are very low (between 500 and 3000 fish) and are likely to be insignificant in the context of the commercial catch.

## 1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial catch is not available.

#### 1.4 Illegal catch

There is no known illegal catch of blue mackerel.

#### 1.5 Other sources of mortality

There is no information on other sources of mortality.

## 2. BIOLOGY

The geographical distribution and habitat of blue mackerel vary with life history stage. Juvenile and immature blue mackerel are northerly in their distribution, having been recorded from commercial and research catches around the North Island and into Golden and Tasman Bay at the top of the South Island.

By contrast, adults have been recorded around both the North and South Islands to Stewart Island and across the Chatham Rise almost to the Chatham Islands. Sporadic catches of small numbers of yearling blue mackerel have been made by bottom trawl in shallow waters.

The distribution of blue mackerel at the surface is seasonal and differs from its known geographical range. During summer, surface schools are found in Northland, BoP, South Taranaki Bight, and Kaikoura, but they disappear during winter, when only occasional individuals are found in Northland and the BoP. A possible corollary to this winter disappearance comes from the peak in bycatch of blue mackerel in the winter jack mackerel mid-water trawl fishery in EMA 7. This suggests an increased partitioning of the population in deeper water at this time of the year, reflecting an observed behavioural characteristic of the related Atlantic species, *Scomber scombrus*. Summaries from aerial sightings data show that blue mackerel can be found in mixed schools with jack mackerel (*Trachurus* spp.), kahawai (*Arripis trutta*), skipjack tuna (*Katsuwonus pelamis*) and trevally (*Pseudocaranx dentex*), and that its appearance in mixed schools varies seasonally.

Blue mackerel are serial spawners, releasing eggs in batches over several months. Based on gonad condition, sexual maturity for both sexes of blue mackerel taken in the Great Australian Bight between January 1979 and December 1980 was estimated to be about 28 cm FL, which translates to an age of about 2 years. Eggs are pelagic and development rate is dependent on temperature. In plankton surveys, blue mackerel eggs have been found from North Cape to East Cape, with highest concentrations from Northland, the Hauraki Gulf, and the Western BoP. Eggs have been described throughout the Hauraki Gulf from November to the end of January, at surface temperatures in the range 15–23°C. Individuals in spent or spawning condition have been taken in a few tows off Tasman Bay and Taranaki, in EMA 7 and in the BoP in EMA 1.

Age and growth studies suggest a difference in the age structures of catches taken in the BoP (New Zealand, EMA 1) and New South Wales (Australia). For fish from the New South Wales study, a peak was found at 1 year that accounted for more than 55% of the fish sampled, with a maximum age of 7 yr. The BoP results show a much broader distribution, with a maximum age of 24 yr, and a mode in the data around 8 to 10 yr. Growth parameters estimated in the BoP study are given in Table 4. Following a quantitative test of competing growth models in the BoP study, no evidence was found of statistically significant differences in growth between the sexes in BoP blue mackerel.

Australian studies may underestimate the ages of larger, older blue mackerel in their catch. The Australian method for estimating blue mackerel ages is based on reading otoliths whole in oil, whereas the New Zealand method is based on otolith thin-sections (Marriott & Manning 2011). Results from the New South Wales study referred to above, suggest that blue mackerel 25–40 cm fork length may be 3–7 years old. Using the New Zealand method, fish in this length range could be as old as 16 years. Australian scientists, reading whole otoliths, may be missing opaque zones near the margin, which are visible in sectioned otoliths.

(EMA)

**Table 4: Von Bertalanffy growth parameters for Bay of Plenty (EMA 1) blue mackerel (Manning et al. 2006).**

	Males	Females	Both sexes
$L_{\infty}$	52.49	53.10	52.79
$K$	0.15	0.15	0.15
$t_0$	-3.29	-3.18	-3.19
Age range	1.8–21.9	1.8–21.9	1.8–21.9
$N$	240	269	509

Although Australian scientists have validated the timing of the first opaque zone in blue mackerel otoliths, their results do not cover the complete life history defined using either the Australian or New Zealand method. A study attempting to validate the New Zealand age estimation method using lead-radium dating indicated that blue mackerel in New Zealand are a relatively long lived small pelagic species, living to at least 17 to 49 years, with the real age most likely nearer the lower value (Marriott et al. 2010). While this range of age estimates is less than desirable for the validation of the growth zone counting method for this species, the findings are consistent with the New Zealand method where otolith ageing studies from commercial catches have blue mackerel living to at least 24 years.

Instantaneous natural mortality ( $M$ ) for male and female fish was estimated using Hoenig's method (Morrison et al. 2001a). Based on age estimates from otoliths collected during the mid 1980s when fishing pressure was presumably light, natural mortality estimates of 0.22 for males and 0.20 for females were derived.

In New Zealand, the diet of blue mackerel has been described as zooplankton, which consists mainly of copepods, but also includes larval crustaceans and molluscs, fish eggs and fish larvae. Feeding involves both filtering of the water and active pursuit of prey, with blue mackerel able to take much smaller animals than, for example, kahawai can.

### 3. STOCKS AND AREAS

Sampling of eggs, larvae, and spawning blue mackerel indicate at least three spawning centres for this species: Northland-Hauraki Gulf; Western BoP; and South Taranaki Bight. Nothing is known of migratory patterns or the fidelity of fish to a particular spawning area. Examination of mitochondrial DNA shows no geographical structuring between New Zealand and Australian fish. Meristic characters show significant regional differentiation within New Zealand fisheries waters and, combined with parasite marker information, Smith et al. (2005) sub-divided blue mackerel into at least three stocks in New Zealand fisheries waters: EMA 1, EMA 2, and EMA 7. No information is currently available on the stock affinity of fish in EMA 3.

### 4. STOCK ASSESSMENT

#### 4.1 EMA 1

##### 4.1.1 Estimates of fishery parameters and abundance

Analysis of aerial sightings data for east Northland (part of EMA 1) from 1985–86 to 2002–03 found no apparent trends in abundance, apart from a peak off east Northland in 1991–92 for both the number of schools and the estimated tonnage, and a further strong signal for the number of schools and the estimated tonnage from 2000–01 to 2002–03.

Using market and catch sampling data collected from 2002 to 2005, estimated numbers-at-length and numbers-at-age were calculated based on all available groomed length and length-at-age data (Manning et al. 2007). These were done separately by sex and scaled to estimates of the total catch from the purse seine fishery. Results showed that the EMA 1 purse seine fishery was composed of fish between 2–21 years of age, although most were between 5 and 15 years.

## 4.2. EMA 7

### 4.2.1 Estimates of fishery parameters and abundance

A standardised CPUE analysis for EMA 7 was carried out using TCEPR tow by tow data from the mid-water trawl jack mackerel target fishery up to 2013–14 in which blue mackerel form a significant and important bycatch (Ballara 2016). The initial dataset comprised tows that targeted jack mackerel with blue mackerel caught as bycatch. Tows that targeted blue mackerel were not considered as they constituted a small amount of catch and effort (about 30 tows each year for the last 10 years by all vessels) and they were confined to a few areas in the fishery and were directed at large sub-surface schools of blue mackerel. Tows that targeted jack mackerel but did not report any blue mackerel catch were also excluded. The data used for the CPUE analyses consisted of catch and effort by core vessels that targeted jack mackerel; core vessels were those participating in the fishery for five or more years, and reporting at least 20 tows per vessel-year. Estimates of relative year effects were obtained using a forward stepwise multiple regression method, where the data were fitted using binomial-lognormal model structure.

Separate standardisations were carried out to two subgroups of core vessels corresponding to an early and late period of the data series respectively. CPUE indices were developed for the early time series from 1989–90 to 1997–98 using catch and effort by 12 core vessels and the late time series from 1996–97 to 2013–14 using catch and effort by 7 core vessels (Table 5). The residual deviance explained was 33% for the early time series and 35% for the late time series. For both data series, the main terms selected by the models are statistical area, vessel, and month.

The early time series increased from 1990 to 1992, and was then relatively constant to 1998. The late time series declined steadily from 1997 to about 2005, and has been relatively constant since then (although the three most recent years produced the lowest indices from this series). Similar trends were also apparent for the later series analysed separately by WCSI and WCNI areas (Figure 1). The series from 2000 onwards shows a decline of more than 50%.

The WG concluded that standardised CPUE series based on the blue mackerel bycatch in the WCNI and WCSI jack mackerel trawl fishery appears to provide reliable indices of abundance.

Using market and catch sampling data collected from 2002 to 2005, estimated numbers-at-length and numbers-at-age were calculated based on all available groomed length and length-at-age data (Manning et al. 2007). These were done separately by sex and scaled to estimates of the total catch from the purse seine and the trawl fisheries. Results showed that the EMA 7 purse seine fishery was composed of fish between 2–24 years of age, although most were between 5 and 15 years. Catch-at-age in the EMA 7 mid-water trawl TCEPR bycatch (jack mackerel target) fishery also showed a wide range, with fish between 2–24 years represented, and small peaks evident between 10 and 11 years in both sexes. These results were generally consistent with those from previous years, although relatively low numbers of small fish in the sampled fisheries were noted.

**Table 5: Standardised lognormal CPUE catch/hr indices for the core West coast TCEPR tow-by-tow target JMA data indices for fishing years 1990–2014. The Standardised CPUE indices for the early series is from 1990 to 1998 (from Fu & Taylor 2011) and for the late series from 1997 to 2014 (Ballara 2016).**

Year	Indices	CV	Indices	CV
1990	0.67	0.20	–	–
1991	0.87	0.10	–	–
1992	1.24	0.11	–	–
1993	1.01	0.13	–	–
1994	0.99	0.09	–	–
1995	1.05	0.07	–	–
1996	0.87	0.11	–	–
1997	1.34	0.08	2.27	0.09
1998	1.13	0.08	1.99	0.07
1999	–	–	2.22	0.05
2000	–	–	2.03	0.05
2001	–	–	1.66	0.05

(EMA)

2002	—	—	1.73	0.04
2003	—	—	1.17	0.05
2004	—	—	0.80	0.04
2005	—	—	0.70	0.04
2006	—	—	0.86	0.04
2007	—	—	0.60	0.04
2008	—	—	0.69	0.04
2009	—	—	0.84	0.04
2010	—	—	0.71	0.04
2011	—	—	0.75	0.04
2012	—	—	0.55	0.05
2013	—	—	0.57	0.05
2014	—	—	0.52	0.06




Figure 2: Blue mackerel CPUE for 1997-2014 for West coast (WC); WCSI, and WCNI. Indices have been standardised to have a mean of one.

#### 4.3 Biomass estimates

No estimates of biomass are available for any blue mackerel stocks.

#### 4.4 Other factors

Catch sampling in the period from 2002 to 2005 indicated that catch-at-length and catch-at-age is relatively stable between years in EMA 1. Although total mortality in EMA 1 is poorly understood, the relatively stable age-length composition between years and the number of year-classes that compose the catch-at-age within fishing years, suggested that blue mackerel may be capable of sustaining the catch levels at that time in EMA 1.

### 5. STATUS OF THE STOCKS

Based on studies of stock structure within New Zealand waters blue mackerel may be sub-divided blue mackerel into at least three stocks: EMA 1, EMA 2, and EMA 7. No information is currently available on the stock affinity of fish in EMA 3.

Little is known about the status of blue mackerel stocks and no estimates of current and reference biomass, or yield, are available for any blue mackerel area.


#### • EMA 1

For EMA 1, the stability of the age composition data and the large number of age classes that comprise the catches suggests that blue mackerel may be capable of sustaining current commercial fishing mortality, at least in the short-term.



## BLUE MACKEREL (EMA)

### • EMA 7

Stock Status	
Year of Most Recent Assessment	2016
Assessment Runs Presented	Standardised trawl CPUE
Reference Points	Target: 40% $B_0$ Soft Limit: 20% $B_0$ Hard Limit: 10% $B_0$ Overfishing threshold: $F_{40\%B_0}$
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	Unknown
Historical Stock Status Trajectory and Current Status	
 <p>West coast blue mackerel CPUE: Comparison of indices for the TCEPR tow-by-tow datasets for fishing years 1990–2014, current analysis with CPUE indices from Fu (2013) and Fu &amp; Taylor (2011). Indices have been standardised to have a mean of one.</p>	

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE has continued to decline since 2009
Recent Trend in Fishing Intensity or Proxy	
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	Broad age structure of the trawl catch (2004-05) did not support a large decrease in biomass from 1999 to 2005 as suggested by the CPUE series

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

(EMA)

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE from the jack mackerel target fishery WCSI and WCNI	
Assessment Dates	Latest assessment: 2016	Next assessment: 2019
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Standardised CPUE - Proportions at age data from the commercial trawl fishery	1 – High Quality 1 – High Quality
Data not used (rank)		
Changes to Model Structure and Assumptions	-	
Major sources of Uncertainty	-	

### Qualifying Comments

The decline in CPUE from 1999 to 2005 was not consistent with the broad range of ages in the trawl catch. However, no recent age data are available.

### Fishery Interactions

There is a small target fishery for blue mackerel on the WCNI but the bulk of the catch is taken as bycatch in the jack mackerel mid-water trawl fishery on the WCSI and WCNI, which has issues with bycatch of kingfish and snapper. Incidental interactions and associated mortality of common dolphins occur in the jack mackerel fishery.

**Table 6: Summary of reported landings (t) and TACCs by QMA for the most recent fishing year.**

		2015–16	2015–16
Fishstock	FMA	TACC	Reported Landings
EMA 1	1	7 630	7 226
EMA 2	2	180	18
EMA 3	3–6	390	27
EMA 7	7–9	3 350	761
EMA 10	10	0	0
TOTAL		11 550	8 033

## 6. FOR FURTHER INFORMATION

- Ballara, S (2016). Characterisation analyses for blue mackerel (*Scomber australasicus*) in EMA 1, 2, 3, and 7, 1989–90 to 2013–14. *New Zealand Fisheries Assessment Report 2016/04*. 108 p.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. Draft New Zealand Fisheries Assessment Report (Unpublished report held by Ministry for Primary Industries, Wellington.). 93 p.
- Boyd, R O; Reilly, J L (2005) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries Research Project REC9803. (Unpublished report held by Ministry for Primary Industries, Wellington.).
- Bradford, E (1996) A comparison of the 1993–94 diary and boatramp surveys of recreational fishing in the Ministry of Fisheries North region. New Zealand Fisheries Assessment Research Document 1996/5. 21 p. (Unpublished report held by NIWA library, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national marine recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Bradford, E; Taylor, P R (1995) Trends in pelagic fish abundance from aerial sightings data. New Zealand Fisheries Assessment Research Document 1995/8. 60 p. (Unpublished report held by NIWA library, Wellington.)
- Fu, D (2013). Characterisation analyses for blue mackerel (*Scomber australasicus*) in EMA 1, 2, 3, and 7, 1989–90 to 2009–10. *New Zealand Fisheries Assessment Report 2013/16*. 54 p.
- Fu, D; Taylor, P R (2007) Standardised CPUE analysis for blue mackerel (*Scomber australasicus*) in EMA 7, 1989–90 to 2004–05. *New Zealand Fisheries Assessment Report 2007/33*. 33 p.
- Fu, D; Taylor, P R (2011) Characterisation and standardised CPUE analyses for blue mackerel (*Scomber australasicus*) in EMA 7, 1989–90 to 2008–09. *New Zealand Fisheries Assessment Report 2011/56*.
- Langley, A; Anderson, F (1998) Sampling the length and age composition of EMA 1 blue mackerel catches from the 1997–98 target purse seine fishery. Unpublished research report, held by Sanfords Tauranga Ltd.
- Marriott, P M; Manning, M J (2011) Reviewing and refining the method for estimating blue mackerel (*Scomber australasicus*) ages. *New Zealand Fisheries Assessment Report 2011/11*.
- Marriott, P M; Manning, M J; Andrews, A H (2010) Investigating blue mackerel (*Scomber australasicus*) age-estimation error. Final Research Report for Ministry of Fisheries Research Project EMA2005-02, Objectives 2–3. 42 p. (Unpublished document held by Ministry for Primary Industries, Wellington.)

## BLUE MACKEREL (EMA)

- Manning, M J; Devine, J A; Marriott, P M; Taylor, P R (2007) The length and age composition of the commercial catch of blue mackerel (*Scomber australasicus*) in EMA 1 and EMA 7 during the 2004–05 fishing year. *New Zealand Fisheries Assessment Report 2007/35*. 36 p.
- Manning, M J; Marriot, P M; Taylor, P (2006) The length and age composition of the commercial catch of blue mackerel (*Scomber australasicus*) in EMA 1 during the 2002–03 fishing year, including a comparison with data collected during the 1997–98 fishing year, and some remarks on estimating blue mackerel ages from otoliths. *New Zealand Fisheries Assessment Report 2006/42*. 42 p.
- Manning, M J; Marriot, P M; Taylor, P R (2007) Length and age composition of the commercial catch of blue mackerel (*Scomber australis*) in EMA 1 and 7 during the 2003–04 fishing year. *New Zealand Fisheries Assessment Report 2007/13*. 41 p.
- Morrison, M; Taylor, P; Marriott, P; Sutton, C (2001) An assessment of information on blue mackerel (*Scomber australasicus*) stocks. *New Zealand Fisheries Assessment Report 2001/44*. 26 p.
- Parrish, R H; MacCall, A D (1978) Climatic variation and exploitation in the Pacific mackerel fishery. *California Department of Fish and Game Bulletin 167*. 110 p.
- Robertson, D A (1978) Blue Mackerel, pilchard, anchovy, sprat, saury, and lantern fish. *New Zealand Fisheries Research Division Occasional Publication 15*. 85–89.
- Rohde, K (1987) Different populations of *Scomber australasicus* in New Zealand and south-eastern Australia, demonstrated by a simple method using monogenean sclerites. *Journal of Fish Biology* 30(6): 651–657.
- Sette, O E (1950) Biology of the Atlantic mackerel (*Scomber scombrus*) of North America. Part II. Migrations and habits. *Fishery Bulletin of the United States Fish and Wildlife Service* 51(49): 251–358.
- Shaun-ror, W (1970) Age and growth of Taiwan spotted mackerel, *S. australasicus*. MSc Thesis, Department of Zoology, National Taiwan University. 38 p. [In Chinese].
- Smith, P J; Diggles, B; Kim, S (2005) Stock structure of blue mackerel, *Scomber australasicus*. *New Zealand Fisheries Assessment Report 2005/43*. 38 p.
- Stevens, J D; Hausfeld, H F; Davenport, S R (1984) Observations on the biology, distribution and abundance of *Trachurus declivis*, *Sardinops neopilchardus* and *Scomber australasicus* in the Great Australian Bight. *Report (CSIRO Marine Laboratories) No. 164*. 29 p.
- Stewart, J; Ferrell, D J; Andrew, N L (1998) Ageing yellowtail (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) in New South Wales. *New South Wales Fisheries Final Report Series, No. 3*. 59 p.
- Stewart, J; Ferrell, D J; Andrew, N L (1999) Validation of the formation and appearance of annual marks in the otoliths of yellowtail (*Trachurus novaezelandiae*) and blue mackerel (*Scomber australasicus*) in New South Wales. *Australian Journal of Marine and Freshwater Research* 50: 389–395.
- Taylor, P R (2002) A summary of information on blue mackerel, (*Scomber australasicus*), characterisation of its fishery in QMAs 7, 8, and 9, and recommendations on appropriate methods to monitor the status of its stock. *New Zealand Fisheries Assessment Report 2002/50*. 68 p.
- Taylor, P R (2008) Factors affecting fish size and landed volumes in the purse-seine and TCEPR charter-boat fisheries in 2004–05 and 2005–06. *New Zealand Fisheries Assessment Report 2008/32*. 17 p.