


RED GURNARD (GUR)

RED GURNARD (GUR)

(Chelidonichthys kumu)

Kumukumu



1. FISHERY SUMMARY

1.1 Commercial fisheries

Red gurnard are a major bycatch of inshore trawl fisheries in most areas of New Zealand, including fisheries for red cod in the southern regions and flatfish on the west coast of the South Island (WCSI) and in Tasman Bay. They are also directly targeted in some areas e.g. GUR 2. Some minor target fisheries for red gurnard are known in Pegasus Bay, off Mahia and off the west coast South Island. Red gurnard is also a minor bycatch in the jack mackerel trawl fishery in the South Taranaki Bight. Up to 15% of the total red gurnard catch is taken by bottom longline and setnet.

Red gurnard was introduced into the Quota Management System (QMS) in 1986. The 1986 TACCs were based on 1984 landings for Southland and 1983 landings for other regions. TACCs for GUR 3 and 7 were increased by 76 t (14%) and 137 t (20%) respectively for the 1991–92 fishing year under the Adaptive Management Programme (AMP), to 600 t in GUR 3 and to 815 t in GUR 7. The GUR 7 TACC was reduced to 678 t, in 1997–98. All AMP programmes ended on 30 September 2009. For the 2009–10 fishing season, the TACC in GUR 7 was increased from 681 t to 715 t, including an allocation of 10 t for customary, 20 t for recreational use, and 14 t allocation for other sources of mortality. The GUR 7 TACC was further increased to 785 t in October 2012 and 845 t in October 2015. The TACC for GUR 3 was increased, by 300 t (50%) to 900 t, for the 1996–97 fishing year under the AMP, but decreased to 800 t in 2002–03. For the 2009–10 fishing season, the TACC for GUR 3 was increased from 800 t to 900 t, with allocations of 3 t, 5 t, and 45 t for customary, recreational, and other sources of mortality respectively. The GUR 3 TACC was increased to 1100 t in October 2012 and to 1220 t in October 2015. This TACC can be seen in Table 1 along with all current allowances, TACCs and TACs.

Table 1: TACs, TACCs and allowances (t) for Red Gurnard by Fishstock.

| Fishstock | TAC | TACC | Customary allowance | Recreational allowance | Other mortality |
|-----------|-------|-------|---------------------|------------------------|-----------------|
| GUR 1 | | 2 287 | | | |
| GUR 2 | | 725 | | | |
| GUR 3 | 1 290 | 1 220 | 3 | 6 | 61 |
| GUR 7 | 919 | 845 | 10 | 22 | 42 |
| GUR 8 | | 543.2 | | | |
| GUR 10 | | 10 | | | |

Reported landings since 1931 are shown in Tables 2 and 3, while an historical record of landings and TACC values for the five main GUR stocks are depicted in Figure 1.

Annual landings of GUR 1 have been relatively stable since 1986–87, generally ranging between 900 and 1300 t; substantially lower than the 2287 t TACC. About 60% of the GUR 1 total is taken from FMA 1, as a bycatch of a number of fisheries including inshore trawl fisheries for snapper, John Dory and tarakihi. The remaining 40% is taken from FMA 9, mainly as a bycatch of the snapper and trevally inshore trawl fisheries.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

| Year | GUR 1 | GUR 2 | GUR 3 | GUR 7 | Year | GUR 1 | GUR 2 | GUR 3 | GUR 7 |
|---------|-------|-------|-------|-------|------|-------|-------|-------|-------|
| 1931–32 | 67 | 0 | 1 | 16 | 1957 | 494 | 402 | 737 | 409 |
| 1932–33 | 42 | 0 | 0 | 13 | 1958 | 430 | 394 | 745 | 400 |
| 1933–34 | 67 | 84 | 1 | 20 | 1959 | 460 | 320 | 806 | 212 |
| 1934–35 | 50 | 179 | 0 | 2 | 1960 | 489 | 417 | 1008 | 421 |
| 1935–36 | 75 | 147 | 18 | 2 | 1961 | 559 | 419 | 1180 | 419 |
| 1936–37 | 114 | 215 | 37 | 25 | 1962 | 505 | 592 | 1244 | 322 |
| 1937–38 | 205 | 193 | 83 | 21 | 1963 | 576 | 562 | 1364 | 367 |
| 1938–39 | 109 | 118 | 151 | 31 | 1964 | 977 | 814 | 1708 | 397 |
| 1939–40 | 121 | 149 | 147 | 25 | 1965 | 1020 | 668 | 1459 | 400 |
| 1940–41 | 124 | 222 | 215 | 38 | 1966 | 1157 | 754 | 1178 | 436 |
| 1941–42 | 107 | 200 | 267 | 38 | 1967 | 1051 | 836 | 745 | 522 |
| 1942–43 | 124 | 332 | 287 | 58 | 1968 | 1137 | 583 | 510 | 368 |
| 1943–44 | 128 | 244 | 294 | 53 | 1969 | 1345 | 632 | 487 | 256 |
| 1944 | 238 | 292 | 291 | 60 | 1970 | 1493 | 823 | 841 | 381 |
| 1945 | 360 | 338 | 222 | 94 | 1971 | 1225 | 570 | 940 | 379 |
| 1946 | 426 | 387 | 290 | 119 | 1972 | 770 | 347 | 662 | 333 |
| 1947 | 376 | 297 | 243 | 162 | 1973 | 1278 | 406 | 1393 | 491 |
| 1948 | 385 | 243 | 267 | 226 | 1974 | 881 | 299 | 1083 | 586 |
| 1949 | 371 | 264 | 316 | 323 | 1975 | 691 | 199 | 655 | 365 |
| 1950 | 306 | 186 | 486 | 332 | 1976 | 1055 | 217 | 960 | 545 |
| 1951 | 221 | 231 | 750 | 202 | 1977 | 1288 | 381 | 975 | 579 |
| 1952 | 394 | 378 | 658 | 211 | 1978 | 1571 | 519 | 1106 | 487 |
| 1953 | 490 | 494 | 614 | 334 | 1979 | 1936 | 382 | 690 | 349 |
| 1954 | 496 | 462 | 660 | 382 | 1980 | 1845 | 438 | 672 | 253 |
| 1955 | 495 | 283 | 652 | 490 | 1981 | 2349 | 603 | 438 | 318 |
| 1956 | 434 | 312 | 782 | 435 | 1982 | 2084 | 454 | 379 | 368 |

| Year | GUR 8 | Year | GUR 8 |
|---------|-------|------|-------|
| 1931–32 | 0 | 1957 | 46 |
| 1932–33 | 0 | 1958 | 51 |
| 1933–34 | 0 | 1959 | 44 |
| 1934–35 | 0 | 1960 | 27 |
| 1935–36 | 0 | 1961 | 27 |
| 1936–37 | 1 | 1962 | 14 |
| 1937–38 | 0 | 1963 | 8 |
| 1938–39 | 2 | 1964 | 16 |
| 1939–40 | 1 | 1965 | 34 |
| 1940–41 | 1 | 1966 | 27 |
| 1941–42 | 0 | 1967 | 45 |
| 1942–43 | 0 | 1968 | 52 |
| 1943–44 | 0 | 1969 | 33 |
| 1944 | 0 | 1970 | 53 |
| 1945 | 3 | 1971 | 37 |
| 1946 | 4 | 1972 | 15 |
| 1947 | 10 | 1973 | 21 |
| 1948 | 9 | 1974 | 41 |
| 1949 | 13 | 1975 | 28 |
| 1950 | 13 | 1976 | 52 |
| 1951 | 10 | 1977 | 45 |
| 1952 | 5 | 1978 | 26 |
| 1953 | 3 | 1979 | 18 |
| 1954 | 7 | 1980 | 34 |
| 1955 | 25 | 1981 | 16 |
| 1956 | 29 | 1982 | 34 |

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

RED GURNARD (GUR)

Table 3: Reported landings (t) of red gurnard by Fishstock from 1983–84 to 2015–16 and actual TACCs (t) from 1986–87 to 2015–16. The QMS data is from 1986–present.

| Fishstock QMA (s) | GUR 1 1 & 9 | | GUR 2 2 | | GUR 3 3, 4, 5 & 6 | | GUR 7 7 | |
|----------------------|----------------|-------|------------|------|----------------------|-------|------------|------|
| | Landings | TACC | Landings | TACC | Landings | TACC | Landings | TACC |
| 1983–84* | 2 099 | - | 782 | - | 366 | - | 468 | - |
| 1984–85* | 1 531 | - | 665 | - | 272 | - | 332 | - |
| 1985–86* | 1 760 | - | 495 | - | 272 | - | 239 | - |
| 1986–87 | 1 021 | 2 010 | 592 | 610 | 210 | 480 | 421 | 610 |
| 1987–88 | 1 139 | 2 081 | 596 | 657 | 386 | 486 | 806 | 629 |
| 1988–89 | 1 039 | 2 198 | 536 | 698 | 528 | 489 | 479 | 669 |
| 1989–90 | 916 | 2 283 | 451 | 720 | 694 | 501 | 511 | 678 |
| 1990–91 | 1 123 | 2 284 | 490 | 723 | 661 | 524 | 442 | 678 |
| 1991–92 | 1 294 | 2 284 | 663 | 723 | 539 | 600 | 704 | 815 |
| 1992–93 | 1 629 | 2 284 | 618 | 725 | 484 | 601 | 761 | 815 |
| 1993–94 | 1 153 | 2 284 | 635 | 725 | 711 | 601 | 469 | 815 |
| 1994–95 | 1 054 | 2 287 | 559 | 725 | 685 | 601 | 455 | 815 |
| 1995–96 | 1 163 | 2 287 | 567 | 725 | 633 | 601 | 382 | 815 |
| 1996–97 | 1 055 | 2 287 | 503 | 725 | 641 | 900 | 378 | 815 |
| 1997–98 | 1 015 | 2 287 | 482 | 725 | 477 | 900 | 309 | 678 |
| 1998–99 | 927 | 2 287 | 469 | 725 | 395 | 900 | 323 | 678 |
| 1999–00 | 944 | 2 287 | 521 | 725 | 411 | 900 | 331 | 678 |
| 2000–01 | 1 294 | 2 287 | 623 | 725 | 569 | 900 | 571 | 678 |
| 2001–02 | 1 109 | 2 287 | 619 | 725 | 717 | 900 | 686 | 681 |
| 2002–03 | 1 256 | 2 287 | 552 | 725 | 888 | 800 | 793 | 681 |
| 2003–04 | 1 225 | 2 287 | 512 | 725 | 725 | 800 | 717 | 681 |
| 2004–05 | 1 354 | 2 287 | 708 | 725 | 854 | 800 | 688 | 681 |
| 2005–06 | 1 113 | 2 287 | 542 | 725 | 957 | 800 | 604 | 681 |
| 2006–07 | 1 180 | 2 287 | 575 | 725 | 1 004 | 800 | 714 | 681 |
| 2007–08 | 1 198 | 2 287 | 517 | 725 | 842 | 800 | 563 | 681 |
| 2008–09 | 1 060 | 2 287 | 621 | 725 | 939 | 800 | 595 | 681 |
| 2009–10 | 1 075 | 2 287 | 853 | 725 | 1 018 | 900 | 603 | 715 |
| 2010–11 | 1 046 | 2 288 | 587 | 725 | 929 | 900 | 545 | 715 |
| 2011–12 | 981 | 2 288 | 558 | 725 | 915 | 900 | 684 | 715 |
| 2012–13 | 1 103 | 2 288 | 603 | 725 | 1 168 | 1 100 | 763 | 785 |
| 2013–14 | 1 005 | 2 288 | 555 | 725 | 1 223 | 1 100 | 837 | 785 |
| 2014–15 | 1 020 | 2 288 | 695 | 725 | 1 150 | 1 100 | 852 | 785 |
| 2015–16 | 860 | 2 288 | 748 | 725 | 1 348 | 1 220 | 852 | 845 |

| Fishstock QMA (s) | GUR 8 | | GUR 10 | | Landings | Total TACC |
|----------------------|----------|------|----------|------|----------|---------------|
| | Landings | TACC | Landings | TACC | | |
| 1983–84* | 251 | - | 0 | - | 3 966 | - |
| 1984–85* | 247 | - | 0 | - | 3 047 | - |
| 1985–86* | 163 | - | 0 | - | 2 929 | - |
| 1986–87 | 159 | 510 | 0 | 10 | 2 403 | 4 230 |
| 1987–88 | 194 | 518 | 0 | 10 | 3 121 | 4 381 |
| 1988–89 | 167 | 532 | 0 | 10 | 2 749 | 4 596 |
| 1989–90 | 173 | 538 | 0 | 10 | 2 745 | 4 730 |
| 1990–91 | 150 | 543 | 0 | 10 | 2 866 | 4 762 |
| 1991–92 | 189 | 543 | 0 | 10 | 3 390 | 4 975 |
| 1992–93 | 208 | 543 | 0 | 10 | 3 700 | 4 978 |
| 1993–94 | 174 | 543 | 0 | 10 | 3 142 | 4 978 |
| 1994–95 | 217 | 543 | 0 | 10 | 2 969 | 4 982 |
| 1995–96 | 182 | 543 | 0 | 10 | 2 927 | 4 982 |
| 1996–97 | 219 | 543 | 0 | 10 | 2 796 | 5 281 |
| 1997–98 | 249 | 543 | 0 | 10 | 2 532 | 5 143 |
| 1998–99 | 170 | 543 | 0 | 10 | 2 284 | 5 143 |
| 1999–00 | 222 | 543 | 0 | 10 | 2 429 | 5 143 |
| 2000–01 | 291 | 543 | 0 | 10 | 3 348 | 5 143 |
| 2001–02 | 302 | 543 | 0 | 10 | 3 429 | 5 143 |
| 2002–03 | 342 | 543 | 0 | 10 | 3 831 | 4 993 |
| 2003–04 | 329 | 543 | 0 | 10 | 3 508 | 4 993 |
| 2004–05 | 370 | 543 | 0 | 10 | 3 974 | 4 993 |
| 2005–06 | 373 | 543 | 0 | 10 | 3 589 | 4 993 |
| 2006–07 | 349 | 543 | 0 | 10 | 3 822 | 4 993 |
| 2007–08 | 223 | 543 | 0 | 10 | 3 344 | 4 993 |
| 2008–09 | 274 | 543 | 0 | 10 | 3 489 | 4 993 |
| 2009–10 | 239 | 543 | 0 | 10 | 3 789 | 5 181 |
| 2010–11 | 182 | 543 | 0 | 10 | 3 289 | 5 181 |
| 2011–12 | 213 | 543 | 0 | 10 | 3 351 | 5 181 |
| 2012–13 | 170 | 543 | 0 | 10 | 3 807 | 5 451 |
| 2013–14 | 151 | 543 | 0 | 10 | 3 769 | 5 451 |
| 2014–15 | 193 | 543 | 0 | 10 | 3 910 | 5 451 |
| 2015–16 | 145 | 543 | 0 | 10 | 3 953 | 5 631 |

*FSU data.

RED GURNARD (GUR)

GUR 2 landings have fluctuated within the range of 400–853 t since 1991–92, typically well below the TACC. In addition to the target fishery, red gurnard are taken as a bycatch of the tarakihi, trevally and snapper inshore trawl fisheries.

GUR 3 landings regularly exceeded the TACC between 1988–89 and 1995–96. Ageing of fish collected during the east coast South Island trawl (ECSI) surveys suggests that there were 1 or 2 relatively strong year classes moving through the fishery, which may help explain the overcatches. GUR 3 has been consistently overcaught since 2004.




Figure 1: Reported commercial landings and TACCs for the five main GUR stocks. From top to bottom: GUR 1 (Auckland East) and GUR 2 (Central East). [Continued on next page].

RED GURNARD (GUR)




Figure 1 [Continued]: Reported commercial landings and TACCs for the five main GUR stocks. From top to bottom: GUR 3 (South East Coast), GUR 7 (Challenger) and GUR 8 (Central Egmont).

GUR 7 landings declined steadily from 761 t in 1992–93, to 309 t in 1997–98, but then increased to a peak of 793 t in 2002–03. They then generally declined to 2010–11, followed by an increase to 2012–13. Landings in GUR 8 have remained well below the levels of the TACC since 1986–87.

1.2 Recreational fisheries

Red gurnard is, by virtue of its wide distribution in harbours and shallow coastal waters, an important recreational species. It is often taken by fishers targeting snapper and tarakihi, particularly around the North Island. The allowances within the TAC for each Fishstock are shown in Table 1.

1.2.1 Management controls

The main methods used to manage recreational harvests of red gurnard are minimum legal size limits (MLS), method restrictions and daily bag limits. Fishers can take up to 20 GUR as part of their combined daily bag limit and the MLS is 25 cm.

1.2.2 Estimates of recreational harvest

Recreational catch estimates are given in Table 4. There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and, offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for red gurnard were calculated using an offsite approach, the offsite regional telephone and diary survey approach. Estimates for 1996 came from a national telephone and diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2002) and a rolling replacement of diarists in 2001 (Boyd & Reilly 2004) allowed estimates for a further year (population scaling ratios and mean weights were not re-estimated in 2001).

The harvest estimates provided by these telephone diary surveys are no longer considered reliable for various reasons. With the early telephone/diary method, fishers were recruited to fill in diaries by way of a telephone survey that also estimates the proportion of the population that is eligible (likely to fish). A “soft refusal” bias in the eligibility proportion arises if interviewees who do not wish to cooperate falsely state that they never fish. The proportion of eligible fishers in the population (and, hence, the harvest) is thereby under-estimated. Pilot studies for the 2000 telephone/diary survey suggested that this effect could occur when recreational fishing was established as the subject of the interview at the outset. Another equally serious cause of bias in telephone/diary surveys was that diarists who did not immediately record their day’s catch after a trip sometimes overstated their catch or the number of trips made. There is some indirect evidence that this may have occurred in all the telephone/diary surveys (Wright et al 2004).

Table 4: Recreational harvest estimates for red gurnard stocks. The telephone/diary surveys and earlier aerial-access surveys ran from December to November but are denoted by the January calendar year. The surveys since 2010 have run through the October to September fishing year but are denoted by the January calendar year. Mean fish weights were obtained from boat ramp surveys (for the telephone/diary and panel survey harvest estimates).

| Stock | Year | Method | Number of fish | Total weight (t) | CV |
|--------------|------|-----------------|----------------|------------------|------|
| <u>GUR 1</u> | 1996 | Telephone/diary | 262 000 | 108 | 0.07 |
| | 2000 | Telephone/diary | 465000 | 223 | 0.16 |
| FMA 1 only | 2005 | Aerial-access | - | 127 | 0.14 |
| FMA 1 only | 2012 | Aerial-access | | | |
| | 2012 | Panel survey | 241 957 | 103 | 0.15 |
| <u>GUR 2</u> | 1996 | Telephone/diary | 38 000 | 16 | 0.18 |
| | 2000 | Telephone/diary | 209 000 | 127 | 0.37 |
| | 2012 | Panel survey | 66 661 | 38 | 0.20 |
| <u>GUR 3</u> | 1996 | Telephone/diary | 1 000 | - | - |
| | 2000 | Telephone/diary | 11 000 | 5 | 0.70 |
| | 2012 | Panel survey | 4 605 | 2 | 0.62 |
| <u>GUR 7</u> | 1996 | Telephone/diary | 26 000 | 12 | 0.15 |
| | 2000 | Telephone/diary | 36 000 | 11 | 0.23 |
| | 2012 | Panel survey | 23 653 | 12 | 0.24 |
| <u>GUR 8</u> | 1996 | Telephone/diary | 67 000 | 28 | 0.15 |
| | 2000 | Telephone/diary | 99 000 | 40 | 0.36 |
| | 2012 | Panel survey | 93 656 | 47 | 0.23 |

RED GURNARD (GUR)

The recreational harvest estimates provided by the 2000 and 2001 telephone diary surveys are thought to be implausibly high for many species, which led to the development of an alternative maximum count aerial-access onsite method that provides a more direct means of estimating recreational harvests for suitable fisheries. The maximum count aerial-access approach combines data collected concurrently from two sources: a creel survey of recreational fishers returning to a subsample of ramps throughout the day; and an aerial survey count of vessels observed to be fishing at the approximate time of peak fishing effort on the same day. The ratio of the aerial count in a particular area to the number of interviewed parties who claimed to have fished in that area at the time of the overflight was used to scale up harvests observed at surveyed ramps, to estimate harvest taken by all fishers returning to all ramps. The methodology is further described by Hartill et al (2007).

This aerial-access method was first employed and optimised to estimate snapper harvests in the Hauraki Gulf in 2003–04. It was then extended to survey the wider SNA 1 fishery in 2004–05 and to provide estimates for other species, including red gurnard (FMA 1 only for GUR). In response to the cost and scale challenges associated with onsite methods, in particular the difficulties in sampling other than trailer boat fisheries, offsite approaches to estimating recreational fisheries harvest have been revisited. This led to the development and implementation of a national panel survey for the 2011–12 fishing year (Wynne-Jones et al 2014). The panel survey used face-to-face interviews of a random sample of New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and catch information collected in standardised phone interviews. Note that the national panel survey estimate does not include recreational harvest taken under s111 general approvals, or catches made by non-residents.

1.3 Customary non-commercial fisheries

Red gurnard is an important species for customary non-commercial fishing interests, by virtue of its wide distribution in shallow coastal waters. However, no quantitative estimates of customary non-commercial catch are currently available.

1.4 Illegal catch

No quantitative information is available.

1.5 Other sources of mortality

No quantitative information is available.

2. BIOLOGY

Gurnard growth rate varies with location, and females grow faster and are usually larger at age than males. Maximum age (A_{MAX}) is about 16 years and maximum size is 55+ cm. Red gurnard reach sexual maturity at an age of 2–3 years and a fork length (FL) of about 23 cm, after which the growth rate slows. An analysis of the age and growth of red gurnard in FMA 7 revealed that young fish 1–4 years old tend to be most common in Tasman and Golden Bays. Three to six year old fish are found on the inshore areas of the West coast South Island and the older fish are predominantly found further offshore (Lyon & Horn 2011).

M was estimated using the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Samples from the ECSI suggested an A_{MAX} of about 16 years for males and 13 years for females, giving estimates for M of 0.29 and 0.35 respectively. Samples from the WCSI indicate an A_{MAX} of about 15 years for both sexes, giving an estimate of 0.31 for M . These samples were not from virgin populations, so M may be overestimated.

Red gurnard have a long spawning period which extends through spring and summer with a peak in early summer. In the Hauraki Gulf, ripe adults can be found throughout the year. Spawning grounds appear to be widespread, although perhaps localised over the inner and central shelf. Egg and larval development takes place in surface waters, and there is a period of at least eight days before feeding starts. Small juveniles (under 15 cm FL) are often caught in shallow harbours, but rarely in commercial trawls.

Biological parameters relevant to the stock assessment are shown in Table 5.

Table 5: Estimates of biological parameters for red gurnard.

| Fishstock | | | Estimate | Source | | |
|---|------------|-------|------------|------------|-------|--------|
| <u>1. Natural mortality (M)</u> | | | | | | |
| | | | Female | Males | | |
| GUR 1W & 1E | | | 0.30 | 0.35 | | |
| GUR 3 | | | 0.29 | 0.35 | | |
| GUR 7 | | | 0.31 | 0.31 | | |
| <u>2. Weight = $a(\text{length})^b$ (Weight in g, length in cm fork length).</u> | | | | | | |
| | | | Both Sexes | | | |
| | | | a | b | | |
| GUR 1 | | | 0.00998 | 2.99 | | |
| GUR 1W & 1E | | | 0.026 | 2.775 | | |
| GUR 2 | | | 0.0053 | 3.19 | | |
| <u>3. von Bertalanffy growth parameters</u> | | | | | | |
| | Females | | | Males | | |
| | L_∞ | k | t_0 | L_∞ | k | t_0 |
| GUR 1 | 36.4 | 0.641 | 0.189 | 28.8 | 0.569 | -0.552 |
| GUR 1W | 45.3 | 0.25 | -0.88 | 36.5 | 0.45 | -0.30 |
| GUR 1E | 44.5 | 0.28 | -0.76 | 35.2 | 0.49 | -0.24 |
| GUR 3 | 48.2 | 0.44 | 0.1 | 42.2 | 0.49 | -0.26 |
| GUR 7 | 45.7 | 0.40 | -0.36 | 40.3 | 0.37 | -0.96 |

3. STOCKS AND AREAS

There are no data that would alter the current stock boundaries. No information is available on stock separation of red gurnard. For GUR 3 the Working Group noted that spatial information from the CPUE analyses indicated that separate stocks or sub-stocks may exist between the East and South coasts of the South Island.

4. STOCK ASSESSMENT

4.1 Biomass estimates

Relative abundance indices have been obtained from trawl surveys of the Bay of Plenty, west coast North Island and Hauraki Gulf within the GUR 1 Fishstock, west coast South Island and Tasman/Golden Bays combined (GUR 7), and east coast South Island (GUR 3) (Table 6). The west coast South Island (WCSI) and east coast South Island (ECSI) surveys are the only ongoing surveys, currently conducted on a biennial basis.

ECSI

The ECSI winter surveys from 1991 to 1996 in 30–400 m were replaced by summer trawl surveys (1996–97 to 2000–01) which also included the 10–30 m depth range, but these were discontinued after the fifth in the annual time series because of the extreme fluctuations in catchability between surveys (Francis et al 2001). The winter surveys were reinstated in 2007 and this time included additional 10–30 m strata in an attempt to index elephant fish and red gurnard which were included in the list of target species. Only the 2007, 2012, 2014 and 2016 surveys provide full coverage of the 10–30 m depth range.

In the 1990s, red gurnard biomass averaged 422 t in the core strata, increasing more than three-fold to 1453 t in 2007. From 2007 to 2014 biomass had an upward trend followed by a substantial decline in 2016 when biomass more than halved. (Table 6, Figure 2). Biomass for the four core plus shallow strata followed the same trend as that for the core strata. The proportion of pre-recruit biomass in the core strata varied greatly among surveys, but was generally low, 2–20%, and in 2016 it was 7%. Similarly, the proportion of juvenile biomass (based on the length-at-50% maturity) within the core strata was close to zero for all surveys (Beentjes et al 2016).

The additional red gurnard biomass captured in the 10–30 m depth range accounted for 29%, 52%, 36% and 61% of the biomass in the core plus shallow strata (10–400 m) for 2007, 2012, 2014 and 2016 respectively, indicating the importance of shallow strata for red gurnard biomass. These observations indicate that the core strata survey (30–400 m) may not be shallow enough to provide an index for sub-mature gurnard.

RED GURNARD (GUR)

Table 6: Relative biomass indices (t) and coefficients of variation (CV) for gurnard for east coast South Island (ECSI) - summer and winter, west coast South Island (WCSI) and the Stewart-Snares Island survey areas*. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16 and 17). The sum of pre-recruit and recruited biomass values do not always match the total biomass for the earlier surveys because at several stations length frequencies were not measured, affecting the biomass calculations for length intervals. – , not measured; NA, not applicable. Recruited is defined as the size-at-recruitment to the fishery (30 cm). [Continued on next page].

| Region | Fishstock | Year | Trip number | Total Biomass estimate | CV (%) | Total Biomass estimate | CV (%) | Pre- recruit | CV (%) | Pre- recruit | CV (%) | Recruited | CV (%) | Recruited | CV (%) |
|----------------------------|-----------|------|-------------|------------------------------|--------|------------------------------|--------|-----------------|--------|-----------------|--------|-----------|--------|-----------|--------|
| Bay of Plenty | | 1983 | KAH8303 | 380 | 23 | - | - | - | - | - | - | - | - | - | - |
| | | 1985 | KAH8506 | 57 | 17 | - | - | - | - | - | - | - | - | - | - |
| | | 1987 | KAH8711 | 410 | 28 | - | - | - | - | - | - | - | - | - | - |
| | | 1990 | KAH9004 | 432 | 12 | - | - | - | - | - | - | - | - | - | - |
| | | 1992 | KAH9202 | 290 | 9 | - | - | - | - | - | - | - | - | - | - |
| | | 1996 | KAH9601 | 332 | 14 | - | - | - | - | - | - | - | - | - | - |
| | | 1999 | KAH9902 | 364 | 14 | - | - | - | - | - | - | - | - | - | - |
| North Island west coast | GUR 9 | 1986 | KAH8612 | 1 763 | 16 | - | - | - | - | - | - | - | - | - | - |
| | | 1987 | KAH8715 | 2 022 | 24 | - | - | - | - | - | - | - | - | - | - |
| | | 1989 | KAH8918 | 1 013 | 12 | - | - | - | - | - | - | - | - | - | - |
| | | 1991 | KAH9111 | 1 846 | 23 | - | - | - | - | - | - | - | - | - | - |
| | | 1994 | KAH9410 | 2 498 | 30 | - | - | - | - | - | - | - | - | - | - |
| | | 1996 | KAH9615 | 1 820 | 14 | - | - | - | - | - | - | - | - | - | - |
| North Island west coast | GUR 8 | 1989 | KAH8918 | 628 | 15 | - | - | - | - | - | - | - | - | - | - |
| | | 1991 | KAH9111 | 817 | 9 | - | - | - | - | - | - | - | - | - | - |
| | | 1994 | KAH9410 | 685 | 22 | - | - | - | - | - | - | - | - | - | - |
| | | 1996 | KAH9615 | 370 | 37 | - | - | - | - | - | - | - | - | - | - |
| | | 1999 | KAH9915 | 2 099* | 13 | - | - | - | - | - | - | - | - | - | - |
| Hauraki Gulf | | 1984 | KAH8421 | 595 | 15 | - | - | - | - | - | - | - | - | - | - |
| | | 1985 | KAH8517 | 49 | 44 | - | - | - | - | - | - | - | - | - | - |
| | | 1986 | KAH8613 | 426 | 36 | - | - | - | - | - | - | - | - | - | - |
| | | 1987 | KAH8716 | 255 | 15 | - | - | - | - | - | - | - | - | - | - |
| | | 1988 | KAH8810 | 749 | 19 | - | - | - | - | - | - | - | - | - | - |
| | | 1989 | KAH8917 | 105 | 29 | - | - | - | - | - | - | - | - | - | - |
| | | 1990 | KAH9016 | 141 | 16 | - | - | - | - | - | - | - | - | - | - |
| | | 1992 | KAH9212 | 330 | 9 | - | - | - | - | - | - | - | - | - | - |
| | | 1993 | KAH9311 | 177 | 17 | - | - | - | - | - | - | - | - | - | - |
| | | 1994 | KAH9411 | 247 | 19 | - | - | - | - | - | - | - | - | - | - |
| | | 1997 | KAH9720 | 242 | 14 | - | - | - | - | - | - | - | - | - | - |
| | | 2000 | KAH0012 | 24 | 46 | - | - | - | - | - | - | - | - | - | - |

*Assuming areal availability, vertical availability and vulnerability equal 1.0. Biomass is only estimated outside 10 m depth except for COM9901 and CMP0001. Note: because trawl survey biomass estimates are indices, comparisons between different seasons (e.g., summer and winter ECSI) are not strictly valid
FMAs 8 and 9 combined

Table 6 [Continued]: Relative biomass indices (t) and coefficients of variation (CV) for gurnard for east coast South Island (ECSI) - summer and winter, west coast South Island (WCSI) and the Stewart-Snares Island survey areas*. Biomass estimates for ECSI in 1991 have been adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16 and 17). The sum of pre-recruit and recruited biomass values do not always match the total biomass for the earlier surveys because at several stations length frequencies were not measured, affecting the biomass calculations for length intervals. -, not measured; NA, not applicable. Recruited is defined as the size-at-recruitment to the fishery (30 cm). Biomass estimates from current surveys with extreme catchability are denoted with an #.

| Region | Fishstock | Year | Trip number | Total | | Total | | Pre-recruit | CV (%) | Pre-recruit | CV (%) | Recruited | CV (%) | Recruited | CV (%) | |
|-----------------------------------|-----------|---------|-------------|------------------|--------|------------------|--------|-----------------|--------|-----------------|--------|-----------------|--------|-----------------|--------|---|
| | | | | Biomass estimate | CV (%) | Biomass estimate | CV (%) | | | | | | | | | |
| South Island | | 1992 | KAH9204 | 572 | 15 | - | - | - | - | - | - | - | - | - | - | - |
| west coast and Tasman/Golden Bays | | 1994 | KAH9404 | 559 | 15 | - | - | - | - | - | - | - | - | - | - | - |
| | | 1995 | KAH9504 | 584 | 19 | - | - | - | - | - | - | - | - | - | - | - |
| | | 1997 | KAH9704 | 471 | 13 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2000 | KAH0004 | 625 | 15 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2003 | KAH0304 | #270 | 20 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2005 | KAH0503 | 442 | 17 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2007 | KAH0704 | 553 | 17 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2009 | KAH0904 | 651 | 18 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2011 | KAH1104 | 1 070 | 17 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2013 | KAH1305 | 754 | 12 | - | - | - | - | - | - | - | - | - | - | - |
| | | 2015 | KAH1503 | 1 774 | 16 | - | - | - | - | - | - | - | - | - | - | - |
| North Island east coast | | 1993 | KAH9304 | 439 | 44 | - | - | - | - | - | - | - | - | - | - | - |
| | | 1994 | KAH9402 | 871 | 16 | - | - | - | - | - | - | - | - | - | - | - |
| | | 1995 | KAH9502 | 178 | 26 | - | - | - | - | - | - | - | - | - | - | - |
| | | 1996 | KAH9605 | 708 | 29 | - | - | - | - | - | - | - | - | - | - | - |
| ECSI (winter) | GUR 3 | | | <u>30-400 m</u> | | <u>10-400 m</u> | | <u>30-400 m</u> | | <u>10-400 m</u> | | <u>30-400 m</u> | | <u>10-400 m</u> | | |
| | | 1991 | KAH9105 | 763 | 33 | - | - | NA | NA | - | - | NA | NA | - | - | |
| | | 1992 | KAH9205 | 142 | 30 | - | - | 21 | 58 | - | - | 121 | 30 | - | - | |
| | | 1993 | KAH9306 | 576 | 31 | - | - | 26 | 45 | - | - | 551 | 31 | - | - | |
| | | 1994 | KAH9406 | 123 | 34 | - | - | 2 | 42 | - | - | 121 | 34 | - | - | |
| | | 1996 | KAH9606 | 505 | 27 | - | - | 8 | 44 | - | - | 496 | 26 | - | - | |
| | | 2007 | KAH0705 | 1 453 | 35 | 2 048 | 27 | 298 | 40 | 494 | 32 | 1 155 | 35 | 1 554 | 27 | |
| | | 2008 | KAH0806 | 1 309 | 34 | - | - | 100 | 59 | - | - | 1 210 | 33 | - | - | |
| | | 2009 | KAH0905 | 1 725 | 30 | - | - | 62 | 34 | - | - | 1 663 | 30 | - | - | |
| | | 2012 | KAH1207 | 1 680 | 28 | 3 515 | 17 | 193 | 40 | 742 | 31 | 1 487 | 27 | 2 773 | 16 | |
| | | 2014 | KAH1402 | 2 063 | 25 | 3 215 | 17 | 409 | 45 | 585 | 32 | 1 654 | 23 | 2 630 | 16 | |
| | | 2016 | KAH1605 | 941 | 30 | 2 420 | 15 | 63 | 41 | 306 | 19 | 877 | 30 | 2 114 | 15 | |
| ECSI (summer) | GUR 3 | 1996-97 | KAH9618 | 765 | 13 | - | - | - | - | - | - | - | - | - | - | |
| | | 1997-98 | KAH9704 | 317 | 16 | - | - | - | - | - | - | - | - | - | - | |
| | | 1998-99 | KAH9809 | 493 | 13 | - | - | - | - | - | - | - | - | - | - | |
| | | 1999-00 | KAH9917 | 202 | 20 | - | - | - | - | - | - | - | - | - | - | |
| | | 2000-01 | KAH0014 | 146 | 34 | - | - | - | - | - | - | - | - | - | - | |

*Assuming areal availability, vertical availability and vulnerability equal 1.0. Biomass is only estimated outside 10 m depth except for COM9901 and CMP0001. Note: because trawl survey biomass estimates are indices, comparisons between different seasons (e.g., summer and winter ECSI) are not strictly valid

RED GURNARD (GUR)

The addition of the 10–30 m depth range had no significant effect on the length frequency distributions in 2007 and 2014, but in 2012 there was a strong 1+ cohort in 10–30 m, which was poorly represented in the core strata (Beentjes et al 2015). Based on the four surveys that included the 10–30 m strata, there are generally more pre-recruit fish in the shallow strata, suggesting that the core plus shallow strata (10 to 400 m) survey is probably indexing red gurnard abundance, including juveniles. The distribution of red gurnard hot spots varies, but overall this species is consistently well represented over the entire survey area from 10 to 100 m, but is most abundant in the shallow 10 to 30 m strata.

WCSI

There was a steady increase over the last five surveys and the estimate for 2015 (1776 t) was the highest in the time series, 66% higher than the previous high in 2011 (1070 t) (Figure 3). Seventy five percent of the total biomass in 2015 was recruited fish (30 cm and over). A significant proportion of the biomass has always occurred in the Tasman and Golden Bay region, although for the last three surveys markedly more was from the west coast South Island, with 71% of the recruited biomass having come from the latter. The increase in total biomass in 2015 comes from both regions. The trends in pre-recruit biomass for the entire survey area has largely followed that of the recruited (>30 cm) fish (Figure 4).

Scaled length frequencies are similar between surveys. Larger numbers of smaller fish are found in Tasman Bay and Golden Bay which is thought to be a nursery area, and larger number of large fish are found on the west coast, although a wide size range occurs in both areas (Stevenson & MacGibbon 2015).




Figure 2: Red gurnard total biomass for all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012, 2014 and 2016. Error bars are \pm two standard deviations.




Figure 3: Red gurnard biomass trends from the west coast South Island inshore trawl survey time series. Error bars are \pm two standard deviations. The red symbol denotes biomass estimated from a survey conducted when catchability was extremely low.




Figure 4: Red gurnard pre-recruit (<30 cm) and recruited biomass trends from the west coast South Island inshore trawl survey time series. Error bars are \pm two standard deviations. The red symbols denote biomass estimated from a survey conducted when catchability was extremely low.

RED GURNARD (GUR)

4.3 CPUE Analyses

GUR 1

In 2012, Kendrick & Bentley (in prep. a) updated CPUE analyses for GUR 1W, GUR 1E, and GUR 1BP (Figures 5 and 6). For each substock, positive catches from single bottom trawl targeted at gurnard, snapper, trevally, tarakihi or John dory were standardised using data from selected core vessels.

The analyses were based on tow based CPUE reported on TCEPR and TCER forms because adequate time series are available in the northern inshore trawl fisheries from 1995–96. Stratum based analyses were also done for each substock which included CELR forms and aggregated data to a common vessel-date-target-area stratum (Table 7). This produced longer time series (from 1989–90) that give an historical perspective to the recent trends.

For each CPUE analysis the suitability of alternative assumptions about the distributions of GLM errors were examined. The distribution which produced the lowest AIC when fitted using a simple, preliminary model was chosen.




Figure 5: Comparison of indices for GUR 1W (upper) and GUR 1E (lower) for bottom trawl based on TCEPR/ TCE format data (tow) with a longer time series (stratum) that includes CELR data, and also with the previous analysis (Kendrick & Bentley 2011) Error bars are ± 1 s.e.




Figure 6: Comparison of indices for GUR 1 BoP for bottom trawl based on TCEPR/ TCE format data (tow) with a longer time series (stratum) which includes CELR data, and also with the previous analysis (Kendrick & Bentley 2011). Error bars are ± 1 s.e.

Table 7: Details of CPUE analyses for each substock of red gurnard in GUR 1.

| | Core vessels | | | Error distribution |
|---------|-------------------------|--------|-----------|--------------------|
| | Criteria (trips, years) | Number | Catch (%) | |
| | West coast | | | |
| Tow | 3, 3 | 34 | 93 | Gamma |
| Stratum | 3, 3 | 46 | 97 | Weibull |
| | East coast | | | |
| Tow | 3, 3 | 41 | 98 | log-logistic |
| Stratum | 3, 3 | 64 | 96 | log-logistic |
| | Bay of Plenty | | | |
| Tow | 3, 3 | 44 | 98 | log-logistic |
| Stratum | 3, 3 | 61 | 97 | weibull |

All three series show strong cyclical fluctuations with a strong recovery from low levels reached between 1995 and 1999 to a peak in the early 2000s followed by a subsequent decline but with bigger magnitude changes evident in the east coast substock than in the other two. The series also differ with respect to the specific years for the nadir and the peak, as well as the nature of the trajectory after the peak in the early 2000s; each is currently near the mean for the series, but the west coast is increasing, while East coast and Bay of Plenty series are in a downward phase.

The Working Group accepted the tow-based series for ongoing monitoring of each substock.

GUR 2

GUR 2 is monitored using standardised CPUE from the bottom trawl fishery targeting gurnard, snapper or trevally.

In 2017, Schofield et al (in prep) updated CPUE analyses for GUR 2 (Figure 7). Landings were allocated to daily aggregated effort using methods described by Langley (2014) to improve the comparability between the data collected from two different statutory reporting forms (CELR and TCER). A core fleet of 49 vessels that had completed at least five trips per year in at least seven years was modelled using a Weibull distribution. The model adjusted for the recent positive influences of shifts in duration, vessel, an area \times month interaction term, and target species, and accounted for 47%

RED GURNARD (GUR)

of the variance in catch. A shorter time series based on TCEPR and TCER format data available since 2007–08, and analysed at tow by tow resolution closely resembles the mixed form series for the years in common (Figure 7).

The NINS WG noted that most of the records in the aggregated data had catches of gurnard and that a binomial index was flat. As a result the positive catch index was retained as the key monitoring series.

In the longer CPUE series using aggregated data there are indications of cyclical variations in abundance with a 4 to 5 year period (Figure 7). There was an overall decreasing trend in CPUE from 1990 to 2007, after which CPUE stabilised and then trended upwards with an especially rapid increase from 2014 to 2016. Spatial residuals indicated that gurnard distribution within Hawke Bay has altered over the period since high resolution data became available in 2008, with evidence of a shift to deeper areas in the centre of the Bay (Figure 8).




Figure 7: Comparison of standardised catch per unit effort (CPUE) indices for GUR 2 from bottom trawling targeting gurnard, snapper and trevally (BT_MIX(day)) combined over all form types, and more recently from data based on TCEPR/ TCER (tow) format data only (Schofield et al., in prep). Both series are scaled relative to the geometric mean of the years they have in common. Fishing years are labelled according to the second calendar year e.g. 1990 = 1989–90. In both standardisation models a Weibull error distribution was assumed.




Figure 8: The mean residuals from the abundance model for GUR 2 BT_MIX(tow). Residuals are aggregated to 0.1° bins and a threshold of 30 tows is applied before a bin was plotted. A: 2008 to 2010, B: 2011 to 2013, C: 2014 to 2016.

Chapman and Robson estimates of total mortality (Z) for GUR 2, based on the age composition of bottom trawl landings in 2009–10, were 0.518 (SE = 0.0159, CV=3.1%) and 0.632 (0.0196, 3.1), depending on whether the age of full recruitment was 2 or 3 years (Parker & Fu 2012). Assuming an instantaneous rate of natural mortality of 0.307, fishing mortality was estimated to be 0.189 or 0.303.

Although it was not possible to produce reliable estimates of spawner biomass per recruit based targets of F (due to unreliable estimates of growth rate and size at maturity), estimates of F from this study were either lower or approximately equal to the estimate of natural mortality (depending on the age at full recruitment assumed). Assuming that the fishery is sampling the age structure of the population, and given that catches and standardised CPUE have been reasonably constant over the last decade, these results suggest that GUR 2 was not over-exploited in 2010, and that the stock is likely to be at or above B_{MSY} .

Establishing B_{MSY} compatible reference points

In 2014, the Working Group adopted mean CPUE from the (BT(MIX)) model for the period 1990–91 to 2009–10 as an B_{MSY} -compatible proxy for GUR 2. The Working Group adopted the default Harvest Strategy Standard definitions for the Soft and Hard Limits of one half and one quarter the target, respectively.

RED GURNARD (GUR)

GUR 3

In 2012, the Working Group accepted two standardised CPUE series for GUR 3 with both series based on the bycatch of red gurnard in bottom trawl fisheries defined by different target species combinations from fishing within the inshore statistical areas of GUR 3 (018, 020, 022,024, 026, 025, 030). The BT(MIX) index included fishing effort targeting RCO, STA, BAR, TAR, GUR while the BT(FLA) index was comprised of FLA target trawls only (Starr & Kendrick 2013).

In 2014, the two CPUE analyses were updated with data from 1989–90 to 2012–13 (Langley 2014). The analysis also included several refinements to improve the comparability between the data collected from two statutory reporting forms (CELR and TCER) which collect data at different levels of detail (daily and by tow), including the approach used to apportion red gurnard landed catches from individual fishing trips to the associated fishing effort records and the daily aggregation of fishing effort. These refinements in data processing resulted in no appreciable change in the resulting CPUE indices for the corresponding period. The 2014 CPUE analyses used the equivalent model formulations to the previous analyses (dependent and explanatory variables and Weibull error structure following Starr & Kendrick 2013).

The two sets of indices were updated in 2015 to include data from 2013–14. The time-series of CPUE indices from the two fisheries are very similar. The indices were at a relatively low level in 1997–98 to 1999–2000 and increased steadily to a peak during 2007–08 to 2010–11 (Figure 9). Both sets of indices were lower than the peak level in 2011–12 to 2013–14, although the indices remained well above the longer term average level from the entire time-series (Figure 9).

The longer term trends in the CPUE indices are similar to the increase in estimates of recruited biomass (defined as fish at least 30 cm T.L.) from the time series of winter ECSI inshore trawl surveys (Figure 9), although the magnitude of the overall increase in the trawl survey biomass is greater than the overall increase in the CPUE indices. Since 2007, the trawl survey biomass estimates have increased and there is no indication of the recent reduction in the CPUE indices from 2011–12 to 2013–14.

Establishing B_{MSY} compatible reference points

In 2012, BT(MIX+FLA), the mean of the BT(MIX) and BT(FLA) series in each year, was accepted by the Working Group as the series for monitoring GUR 3. These fisheries cover different aspects of gurnard distribution, both by depth and spatially, but still have very similar trajectories, providing some confidence that these series are likely to be tracking abundance. The mean from 1997–98 to 1999–00 of BT(MIX+FLA) was selected as the Soft Limit because it was a well-defined low point in the series, along with the observations that both catch and CPUE increased simultaneously from that point. The Working Group accepted the default Harvest Strategy Standard definitions that the target “ B_{MSY} -compatible proxy” for GUR 3 would be twice the Soft Limit and the Hard Limit was one-half the Soft Limit.




Figure 9: Standardised CPUE indices for two east coast South Island bottom trawl fisheries [BT(MIX) and BT(FLA)] compared to trawl survey estimates of recruited (≥ 30 cm T.L.) biomass for red gurnard from the winter ECSI inshore trawl survey for two survey areas (30–400 m and 10–400 m). Error bars show $\pm 95\%$ confidence intervals.

GUR 7

In both 2014 and 2017, only two standardised CPUE analyses based on the catch of gurnard in bottom trawl fisheries operating off of the west coast of the South Island for monitoring GUR 7 were accepted. These fisheries are defined as follows:

- WCSI(FLA): bottom trawl effort targeted at FLA (or any of the species that make up this complex) and fishing in Statistical Areas 033, 034, 035 or 036;
- WCSI(MIX): bottom trawl effort targeted at GUR, RCO, TAR, BAR, STA, WAR and fishing in Statistical Areas 033, 034, 035 or 036;

The data for these analyses were prepared using the “daily effort” procedure documented in Langley (2014). The Plenary agreed in 2017 to use the combined model (lognormal model of positive catches and binomial model of probability of capture) using the delta-lognormal method (Vignaux 1994) for stock evaluations. This was done because the Inshore WGs have adopted the standard of combining positive catch and fishing success models when there is a trend in the proportion zero catch. In addition, simulation work has indicated that calculating a combined index may reduce bias when reporting small catch amounts (Langley 2015).

These fishery definitions build on the work of Kendrick et al (2011) and Langley (2014), who defined four fisheries for monitoring GUR 7, two on the WCSI and two in western Cook Strait/Tasman-Golden Bays, some with slightly different target species definitions than indicated above. These four GUR 7 BT fisheries were reviewed in 2014, comparing the CPUE series with the red gurnard biomass indices obtained from the west coast South Island trawl survey (Table 6). The Plenary rejected the two series based on catch-effort data from Tasman/Golden Bays, partly because those series did not match the biomass survey indices very well, and because there was a marked shift in the spatial distribution of fishing effort in the western Cook Strait fishery, with a reduction in the proportion of fishing effort within the areas of higher red gurnard catch rates and a shift towards trawling in deeper waters (Langley 2014). On the other hand, the two sets of CPUE indices from the west coast South Island fisheries showed similar cyclical trends with relatively high CPUE indices during 1990–91 to 1991–92 and 2001–02 to 2003–04 and also relatively low CPUE indices in 1993–94 to 1999–2000 and 2006–07 to 2010–11 (Figure 10). These CPUE indices have since steadily increased from 2009–10 to a high level in 2015–16.

RED GURNARD (GUR)

A composite series (WCSI(MIX+FLA)), which averaged the WCSI(MIX) and WCSI(FLA) series in each year, was accepted in 2014 by the Plenary as the best CPUE series for monitoring GUR 7.

The biomass estimates of recruited (≥ 30 cm T.L.) red gurnard from the WCSI trawl survey do not show the same strong abundance signal in the early to mid-2000s as do the CPUE indices. However, with the omission of the 2003 survey on the basis of an apparently large (negative) change in catchability (see Appendix 6, Stevenson & MacGibbon 2015), the trends are not incompatible. Also, recent survey biomass estimates in 2011, 2013 and 2015 are consistent with the high levels of CPUE observed in the two WCSI BT series (Figure 10).

Establishing B_{MSY} compatible reference points

The Plenary reviewed the WCSI trawl survey biomass estimates in 2017 and concluded that there was no need to separate the Tasman/Golden Bay strata from the WCSI strata, given the strong similarity in the biomass signals from the two survey components in 9 of the 11 survey years. Consequently, it was agreed that the recruited biomass from the total survey should be used as the main tool for monitoring GUR 7.

The Plenary concluded that the trawl survey time series is a better index of trends in abundance than the CPUE time series, primarily because it is more consistent through time and is not affected by changes in fishing behaviour. The mean of the WCSI trawl survey series from 1992–2013, but excluding 2003 because of a large negative change in catchability, was chosen as a “ B_{MSY} compatible proxy” for GUR 7 on the basis that this was a period of relative stability in the series. The Plenary then adopted the default Harvest Strategy Standard definitions that the Soft and Hard Limits would be one half and one quarter the target, respectively.

The averaged WCSI(MIX+FLA) series was retained for corroboration purposes only, with no associated reference points being derived from it.




Figure 10: Comparison of the combined indices from two independent CPUE series for GUR 7 from the inshore WCSI bottom trawl fisheries (Statistical Areas 033, 034, 035, and 036); a) WCSI(FLA): target FLA; b) WCSI(MIX): target, GUR, BAR, TAR, WAR, STA, RCO. Trawl survey biomass estimates of recruited (≥ 30 cm T.L.) red gurnard from the WCSI inshore trawl survey are also presented (the 2017 index [in red] is preliminary) and the excluded 2003 survey estimate is plotted with a hollow marker. The vertical bars represent the associated 95% confidence intervals.

4.4 Other factors

Red gurnard is a major bycatch of target fisheries for several different species, such as snapper and flatfish. The target species may differ between areas and seasons. The recorded landings are influenced directly by changes in the fishing patterns of fisheries for these target species and indirectly by the abundance of these target species. Some target fishing for gurnard also occurs.

4.5 Future research needs

- Investigate the potential benefits of undertaking a full stock assessment for GUR 7, which would entail conducting more ageing of otoliths.
- Further investigation of the relationship between pre-recruits and subsequent recruitment may be useful.

5. STATUS OF THE STOCKS

Stock Structure Assumptions


For the purpose of this summary GUR 1 is considered to be a single stock with three sub-stocks.

- **GUR 1W**


| Stock Status | |
|-----------------------------------|---|
| Year of Most Recent Assessment | 2013 |
| Assessment Runs Presented | Standardised CPUE |
| Reference Points | Target: B_{MSY} -compatible proxy based on the mean CPUE from 1994–95 to 2011–12 of the bottom trawl GUR 1 west (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period: 1994–95 to 2011–12 |
| Status in relation to Target | About as Likely as Not (40–60%) to be at or above B_{MSY} |
| Status in relation to Limits | Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%) |
| Status in relation to Overfishing | Overfishing is Unlikely (< 40%) to be occurring |
| | |

RED GURNARD (GUR)

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for red gurnard in GUR 1W from models of catch rate in successful bottom trawl trips done for tow by tow data from 1995–96 (± 2 s.e.) and at stratum level including CELR data from 1989–90 (Kendrick & Bentley in prep. a). Also shown is the trajectory of total landed GUR 1 from the sub-stock area. The two CPUE series have been scaled to the mean of each series for the years in common. Horizontal lines represent the target and soft and hard limits.



Annual relative exploitation rate for red gurnard in the GUR 1 west coast sub-stock.

Fishery and Stock Trends

| | |
|--|--|
| Recent Trend in Biomass or Proxy | The CPUE index cycles over a 4–8 year period consistent with the dynamics of a short lived species with variable recruitment. CPUE suggests that stock size has fluctuated around the long-term average since 1995–96, recovering from lows in 1998–99 and 2008–09. The CPUE has increased since 2008–09 and in 2011–12 was slightly above the long-term mean. |
| Recent Trend in Fishing Intensity or Proxy | Relative exploitation rate has fluctuated without trend since 1991–92. |
| Other Abundance Indices | The GUR 1West (stratum) series is slightly longer than the GUR 1 West (tow) series, but has a similar trend for the overlapping period. |
| Trends in Other Relevant | |

| | |
|-------------------------|---|
| Indicators or Variables | - |
|-------------------------|---|

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Unlikely if the catch remains at current levels Hard Limit: Unlikely if the catch remains at current levels Unknown whether catch at the level of the TACC would cause decline below both the soft and hard Limits |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | Probability of TACC causing overfishing to occur or commence: Unlikely if the catch remains at current levels Unknown whether catch at the level of the TACC would cause overfishing |

| Assessment Methodology and Evaluation | |
|--|---|
| Assessment Type | Level 2 - Partial quantitative stock assessment |
| Assessment Method | Standardised CPUE based on positive catches from bottom trawl |
| Assessment Dates | Latest assessment: 2013 Next assessment: 2018 |
| Overall assessment quality rank | 1 – High Quality |
| Main data inputs (rank) | Catch and effort data 1 – High Quality |
| Data not used (rank) | N/A |
| Changes to Model Structure and Assumptions | The accepted CPUE index is now a tow based index, rather than trip-stratum based. |
| Major Sources of Uncertainty | |

| Qualifying Comments |
|---|
| <p>As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation, and in two sub-stocks trends are currently downward. This makes it difficult to predict future trends without recruitment information. Given that the catch levels observed from 1986–87 to 2011–12 has been relatively consistent (averaging 1129 t for all of GUR 1) and that red gurnard are mainly taken as bycatch, current catch levels are unlikely to compromise the long-term viability of this stock.</p> <p>As the TACC is substantially higher than the current catch, it is not possible to evaluate potential impacts if catches increased to the level of the TACC.</p> |

| Fishery Interactions |
|--|
| Red gurnard is taken on the west coast by bottom trawl targeted at snapper and trevally. Incidental captures of seabirds occur and there is a risk of incidental capture of Maui's dolphins. |


- **GUR 1E**

| Stock Status | |
|--------------------------------|--|
| Year of Most Recent Assessment | 2013 |
| Assessment Runs Presented | Standardised CPUE |
| Reference Points | Target: B_{MSY} -compatible proxy based on the mean CPUE |


RED GURNARD (GUR)

| | |
|-----------------------------------|---|
| | from 1995–96 to 2011–12 for the bottom trawl GUR 1 East (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period: 1995–96 to 2011–12 |
| Status in relation to Target | About as Likely as Not (40–60%) to be at or above B_{MSY} |
| Status in relation to Limits | Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%) |
| Status in relation to Overfishing | Unknown whether Overfishing is occurring |

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for red gurnard in GUR 1E from models of catch rate in successful bottom trawl trips done for tow by tow data from 1995–96 (± 2 s.e.) and at stratum level including CELR data from 1989–90 (Kendrick & Bentley in prep. a). Also shown is the trajectory of total landed GUR 1 from the substock area. The two CPUE series have been scaled to the mean of each series for the years in common. Horizontal lines represent the target and the soft and hard limits.



Annual relative exploitation rate for red gurnard in the GUR 1 east coast sub-stock.

| Fishery and Stock Trends | |
|--|---|
| Recent Trend in Biomass or Proxy | The CPUE index fluctuates in a way that is consistent with the dynamics of a short lived species with variable recruitment, although the period is longer than that for other gurnard stocks. An increase from the lowest levels in 1995–96 was sustained over eight consecutive years, peaked in 2004–05 and has since declined to slightly below the target in 2011–12. |
| Recent Trend in Fishing Intensity or Proxy | Relative exploitation rate increased from 1989–90 to 1996–97, declined to 1998–99 and has since then fluctuated without trend below the long-term average. |
| Other Abundance Indices | The GUR 1 East (stratum) series is slightly longer than the GUR 1 East (tow) series, but has a similar trend for the overlapping period. |
| Trends in Other Relevant Indicators or Variables | - |

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Unknown Hard Limit: Unknown |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | Unknown if the catch remains at current levels Unknown whether catch at the level of the TACC would cause overfishing. |

| Assessment Methodology and Evaluation | |
|--|---|
| Assessment Type | Level 2 - Partial quantitative stock assessment |
| Assessment Method | Standardised CPUE based on positive catches from bottom trawl |
| Assessment Dates | Latest assessment: 2013 Next assessment: 2018 |
| Overall assessment quality rank | 1 – High Quality |
| Main data inputs (rank) | - Catch and effort data 1 – High Quality |
| Data not used (rank) | N/A |
| Changes to Model Structure and Assumptions | The accepted CPUE index is now a tow based index, rather than trip-stratum based. |
| Major Sources of Uncertainty | - |

| Qualifying Comments |
|---|
| <p>As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation, and in two sub-stocks trends are currently downward. This makes it difficult to predict future trends without recruitment information. Given that the catch levels observed from 1986–87 to 2011–12 has been relatively consistent (averaging 1129 t for all of GUR 1) and that red gurnard are mainly taken as bycatch, current catch levels are unlikely to compromise the long-term viability of this stock.</p> <p>As the TACC is substantially higher than the current catch, it is not possible to evaluate potential impacts if catches increased to the level of the TACC.</p> |


| Fishery Interactions |
|---|
| Red gurnard is taken as a bycatch on the east coast mainly by bottom longline targeted at snapper, with the balance taken almost equally by bottom trawl and Danish seine targeting snapper and John dory. Incidental captures of seabirds occur. |

RED GURNARD (GUR)


- GUR 1 Bay of Plenty

| Stock Status | |
|-----------------------------------|---|
| Year of Most Recent Assessment | 2013 |
| Assessment Runs Presented | Standardised CPUE |
| Reference Points | Target: B_{MSY} -compatible proxy based on the mean CPUE from 1994–95 to 2011–12 for the bottom trawl GUR 1 BoP (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period: 1994–95 to 2011–12 |
| Status in relation to Target | About as Likely as Not (40–60%) to be at or above B_{MSY} |
| Status in relation to Limits | Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%) |
| Status in relation to Overfishing | Unknown whether Overfishing is occurring |

Historical Stock Status Trajectory and Current Status



Comparison of standardised CPUE for red gurnard in GUR 1BoP from models of catch rate in successful bottom trawl trips done for tow by tow data from 1995–96 (± 2 s.e.) and at stratum level including CELR data from 1989–90 (Kendrick & Bentley in prep. a). Also shown is the trajectory of total landed GUR 1 from the substock area. The two CPUE series have been scaled to the mean of each series for the years in common. Horizontal lines represent the target and the soft and hard limits.



Annual relative exploitation rate for red gurnard in the Bay of Plenty.

| Fishery and Stock Trends | |
|--|---|
| Recent Trend in Biomass or Proxy | The CPUE index fluctuates in a way that is consistent with the dynamics of a short lived species with variable recruitment. An increase from the lowest levels in 1995–96 to a peak in 2000–01, and has since declined to slightly below the target in 2011–12. |
| Recent Trend in Fishing Intensity or Proxy | Relative exploitation rate has fluctuated without trend around the long-term mean since 1991–92 |
| Other Abundance Indices | The GUR 1 BoP (stratum) series is slightly longer than the GUR 1 BoP (tow) series, but has a similar trend for the overlapping period. |
| Trends in Other Relevant Indicators or Variables | - |

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Unknown Hard Limit: Unknown |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | Unknown if the catch remains at current levels Unknown whether catch at the level of the TACC would cause overfishing. |

| Assessment Methodology and Evaluation | | |
|--|---|-----------------------|
| Assessment Type | Level 2 - Partial quantitative stock assessment | |
| Assessment Method | Standardised CPUE based on positive catches from bottom trawl | |
| Assessment Dates | Latest assessment: 2013 | Next assessment: 2018 |
| Overall assessment quality rank | 1 – High Quality | |
| Main data inputs (rank) | - Catch and effort data | 1 – High Quality |
| Data not used (rank) | - | |
| Changes to Model Structure and Assumptions | The accepted CPUE index is now a tow based index, rather than trip-stratum based. | |
| Major Sources of Uncertainty | - | |

| Qualifying Comments |
|---|
| <p>As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation, and in two sub-stocks trends are currently downward. This makes it difficult to predict future trends without recruitment information. Given that the catch levels observed from 1986–87 to 2011–12 has been relatively consistent (averaging 1129 t for all of GUR 1) and that red gurnard are mainly taken as bycatch, current catch levels are unlikely to compromise the long-term viability of this stock.</p> <p>As the TACC is substantially higher than the current catch, it is not possible to evaluate potential impacts if catches increased to the level of the TACC.</p> |

| Fishery Interactions |
|--|
| Red gurnard is taken as a bycatch in the Bay of Plenty mainly by bottom longline targeted at snapper, with the balance taken almost equally by bottom trawl and Danish seine targeting snapper and John dory. Incidental captures of seabirds occur. |

RED GURNARD (GUR)


• **GUR 2**

Stock Structure Assumptions


For the purpose of this summary GUR 2 is considered to be a single stock.

| Stock Status | |
|-----------------------------------|--|
| Year of Most Recent Assessment | 2017 |
| Assessment Runs Presented | Standardised CPUE for BT.MIX |
| Reference Points | Target: B_{MSY} -compatible proxy based on the mean CPUE (BT(MIX)) for period 1990–91 to 2009–10 Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period 1990–91 to 2009–10 |
| Status in relation to Target | Very Likely (> 90%) to be above the target |
| Status in relation to Limits | Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%) |
| Status in relation to Overfishing | Overfishing is Unlikely (< 40%) to be occurring |

Historical Stock Status Trajectory and Current Status



Annual landings and standardised catch per unit effort (CPUE) index for GUR 2 from bottom trawling targeting gurnard, snapper and trevally (BT_MIX(day)) that combines all form types at a daily aggregation (Schofield et al in prep.). Scaling is relative to the years in common. A Weibull error distribution was assumed. Horizontal lines are the target and the soft/hard limits.



Annual relative exploitation rate (catch/CPUE) for red gurnard in GUR 2.

| Fishery and Stock Trends | |
|--|--|
| Recent Trend in Biomass or Proxy | CPUE indices generally trended downwards between 1990 and 2007, then flattened to 2012, with a strong increase to 2016. Standardised CPUE in 2015–16 is well above the target. |
| Recent Trend in Fishing Intensity or Proxy | Relative exploitation rate increased gradually from 1989–90 to 2009–10 and then dropped to below the long-term average from 2013–14. |
| Other Abundance Indices | Tow based analysis of 2007–08 to 2015–16 data closely resembles the mixed form type analysis. |
| Trends in Other Relevant Indicators or Variables | Catch curve analysis indicated that fishing mortality was at or below M in 2010 (depending on the age at full recruitment). |

| Projections and Prognosis | |
|---|--|
| Stock Projections or Prognosis | Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%) Unknown whether catch at the level of the TACC would cause decline below both the soft and hard Limits. |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | Unlikely (< 40%) if the catch remains at the average of 2000–2013 levels Unknown whether catch at the level of the TACC would cause overfishing |

| Assessment Methodology and Evaluation | |
|--|--|
| Assessment Type | Level 2 - Partial quantitative stock assessment |
| Assessment Method | Standardised CPUE |
| Assessment Dates | Latest assessment: 2017 Next assessment: 2018 |
| Overall assessment quality rank | 1 – High Quality |
| Main data inputs (rank) | BT.Mix CPUE series 1 – High Quality |
| Data not used (rank) | N/A |
| Changes to Model Structure and Assumptions | - Weibull instead of gamma error structure for CPUE analysis |

RED GURNARD (GUR)

| | |
|------------------------------|---|
| Major Sources of Uncertainty | - |
|------------------------------|---|

Qualifying Comments

Most of the GUR2 commercial catch is made in Hawke Bay, and the index of abundance is naturally weighted to abundance of GUR in this area.

Fishery Interactions


Red gurnard is taken in FMA 2 by the bottom trawl fishery targeting snapper, gurnard and trevally and as a bycatch in bottom trawl fisheries targeting flatfish and tarakihi. Incidental captures of seabirds occur and there is a risk of incidental capture of Hector's dolphins at the southern end of the QMA.

• GUR 3

Stock Structure Assumptions

No information is available on the stock separation of red gurnard. The Fishstock GUR 3 is treated in this summary as a unit stock.

| Stock Status | |
|--|---|
| Year of Most Recent Assessment | 2015 |
| Assessment runs presented | The CPUE series BT(MIX+FLA), which is the mean of two standardised bottom trawl CPUE series: one based on bottom trawls targeting mixed species (RCO, STA, BAR, TAR, GUR) and the other based on flatfish targeting. |
| Reference Points | Target: B_{MSY} -compatible proxy based on CPUE is twice the soft limit Soft Limit: Mean from 1997–98 to 1999–00 of BT(MIX+FLA) series, as defined in Starr & Kendrick (2012) Hard Limit: 50% of soft limit Overfishing threshold: F_{MSY} |
| Status in relation to Target | Likely (> 60%) to be above the target |
| Status in relation to Limits | Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below |
| Status in relation to Overfishing | About as Likely as Not (40–60%) to be overfishing |
| Historical Stock Status Trajectory and Current Status | |
| East coast South Island winter trawl survey, CPUE, Catch and TACC Trajectories | |



Comparison of east coast South Island winter trawl survey recruited biomass and CPUE indices (average FLA and MIX) and the trajectories of catch and TACCs from 1989–90 to 2013–14. The horizontal grey line represents the MSY proxy relative to the CPUE series. The black dotted and solid lines represent the soft and hard limits, respectively.

| Fishery and Stock Trends | |
|--|--|
| Recent Trend in Biomass or Proxy | Two bottom trawl CPUE series (one targeted at flatfish and the other at RCO, STA, BAR, TAR, GUR), which are considered to be an index of stock abundance, increased steadily from the late 1990s to 2009–10, and then declined, remaining above the target level. |
| Recent Trend in Fishing Intensity or Proxy | <p>The graph shows 'Standardised Effort = F proxy' on the y-axis (0.0 to 1.2) against 'Fishing year' on the x-axis (89/90 to 13/14). The data points, connected by a dotted line, fluctuate around a mean of approximately 1.0, with a notable peak in 2013-14 reaching above 1.2.</p> <p>Fishing mortality proxy is Standardised Fishing Effort = Total catch/CPUE (normalised). Fishing mortality proxy increased sharply from 2010–11 to 2013–14 to above the series mean in 2011–12 and 2013–14.</p> |
| Other Abundance Indices | ECSI winter survey (30–400 m) shows a substantial increase since the early 1990s, but declining in 2016. The expanded survey (10–400 m) shows a marked increase from 2007–2014, but declining in 2016 (n = 4). |
| Trends in Other Relevant Indicators or Variables | - |

RED GURNARD (GUR)

| Projections and Prognosis | |
|---|---|
| Stock Projections or Prognosis | Quantitative stock projections are unavailable. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Very Unlikely (< 40%) Hard Limit: Very Unlikely (< 10%) Current abundance is at historically high levels and is unlikely to decline below limits in 3–5 years. |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | GUR is mostly taken as a bycatch (about 10% targeted). The correspondence between relative abundance and catch suggests a constant exploitation rate. The current catch is therefore Unlikely (< 40%) to cause overfishing. |

| Assessment Methodology and Evaluation | | |
|--|---|------------------------------------|
| Assessment Type | Level 2: Partial Quantitative Stock Assessment | |
| Assessment Method | Agreed standardised CPUE series and trawl survey biomass indices | |
| Assessment Dates | Latest assessment: 2015 | Next assessment: 2018 |
| Overall assessment quality rank | 1 – High Quality | |
| Main data inputs (rank) | -Trawl survey biomass indices and associated length frequencies - Catch and effort data | 1– High Quality 1– High Quality |
| Data not used (rank) | N/A | |
| Changes to Model Structure and Assumptions | - | |
| Major Sources of Uncertainty | Prior to 2007 the ECSI trawl survey did not cover the entire depth range for red gurnard. Variable proportion of the population in the previously unsurveyed 10–30 m depth range suggests that survey catchability varies between years in the core survey area (30–400 m). | |

| Qualifying Comments |
|--|
| <p>Red gurnard are relatively short-lived and reasonably productive. They exhibit cyclic fluctuations and were at low levels in the mid-1990s. Stock size has increased substantially since then and commercial fishers indicate that they find it difficult to stay within the TACC despite the low level of targeting on this species.</p> <p>Two independent CPUE series and the winter trawl survey corroborate that stock size for GUR 3 has increased since the late 1990s.</p> <p>There are potentially sufficient data to undertake a quantitative stock assessment for GUR 3. This would allow the estimation of B_{MSY} and other reference points.</p> |

| Fishery Interactions |
|--|
| <p>Red gurnard in GUR 3 are taken almost entirely by bottom trawl in fisheries targeted at red cod, barracouta and flatfish. Some gurnard are also taken in the target tarakihi and stargazer bottom trawl fisheries. The level of targeting on this species is low, averaging less than 10% of the total landed catch since 1989–90.</p> <p>Incidental captures of seabirds occur and there is a risk of incidental capture of Hector's dolphins.</p> |

• **GUR 7**


Stock Structure Assumptions

Stock boundaries are unknown, but for the purpose of this summary, GUR 7 is considered to be a single management unit.

Advice for GUR 7 is based on the biomass series for the recruited portion of the total WCSI trawl survey.

| Stock Status | |
|-----------------------------------|---|
| Year of Most Recent Assessment | 2017 |
| Assessment runs presented | West Coast South Island trawl survey |
| Reference Points | Target: B_{MSY} -compatible proxy based on the mean WCSI trawl survey indices from 1992 to 2013, but excluding the 2003 index Soft Limit: 50% target Hard Limit: 25% target Overfishing threshold: F_{MSY} compatible proxy based on the WCSI trawl survey mean relative exploitation rate from 1992 to 2013, excluding the 2003 index |
| Status in relation to Target | Very Likely (> 90%) to be at or above the target |
| Status in relation to Limits | Soft limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below |
| Status in relation to Overfishing | Overfishing is Unlikely (< 40%) to be occurring |


Historical Abundance and Catch Trajectories



Comparison of the GUR WCSI trawl survey indices with the QMR/MHR landings and TACC for GUR 7. The agreed B_{MSY} proxy (geometric average: 1992–2013 (excluding 2003) WCSI survey biomass estimates=460 t) is shown as a green line; the calculated Soft Limit (=0.5x B_{MSY} proxy) is shown as a purple line; the calculated Hard Limit (=0.25x B_{MSY} proxy) is shown as a grey line. The provisional 2017 WCSI biomass estimate is shown in red and the excluded 2003 survey is shown as a hollow marker.

RED GURNARD (GUR)

Fishing Intensity Trajectories



Relative fishing pressure for GUR 7 based on the ratio of QMR/MHR landings relative to the WCSI trawl survey (recruited). Horizontal green line is the geometric mean fishing pressure from 1992 to 2013, excluding 2003. Fishing pressure for the excluded 2003 survey is shown as a hollow marker.

Fishery and Stock Trends

Recent trend in Biomass or Proxy


The West Coast South Island trawl survey relative biomass indices from 2015 and (preliminary) 2017 were by far the highest of the entire time series.

Recent trend in Fishing Intensity or Proxy

Unlikely (< 40%) that overfishing is occurring
Biomass has increased considerably since 2009–10 while there has been only a moderate increase in annual catches.

Other Abundance Indices

WCSI CPUE indices increased from 2009–10 to 2015–16.



Mean WCSI-BT(FLA+MIX) CPUE series compared with WCSI(recruited) trawl survey. Excluded 2003 survey index shown with hollow marker and preliminary 2017 survey index in red.

| | |
|---|---|
| Trends in Other Relevant Indicators or Variables | Estimates of pre-recruit fish from the West Coast South Island inshore trawl survey indicate that recruitment has been increasing since about 2005 and is currently well above average. |
| Projections and Prognosis | |
| Stock Projections or Prognosis | Quantitative stock projections are unavailable. However, above average recruitment is likely to ensure continuing high biomass at current catch levels, at least in the short term. |
| Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits | Soft Limit: Very Unlikely (< 10%) Hard Limit: Very Unlikely (< 10%) Current abundance is at historically high levels and is unlikely to decline below limits in 3–5 years |
| Probability of Current Catch or TACC causing Overfishing to continue or to commence | Unlikely (< 40%) |

| | | |
|--|---|--------------------------------------|
| Assessment Methodology and Evaluation | | |
| Assessment Type | Level 2: Partial Quantitative Stock Assessment | |
| Assessment Method | West Coast South Island trawl survey biomass - Survey length frequency - Standardised CPUE indices | |
| Assessment Dates | Latest assessment: 2017 | Next assessment: 2019 |
| Overall assessment quality rank | 1 – High Quality | |
| Main data inputs | - Survey biomass and length frequencies - CPUE indices | 1 – High Quality 1 – High Quality |
| Changes to Model Structure and Assumptions | - Tasman and Golden Bay survey data combined into the WCSI survey series - WCSI trawl survey series given precedence over the CPUE series for monitoring abundance - Use of the WCSI survey only to derive reference points CPUE used to provide corroboration | |
| Major Sources of Uncertainty | - Choice of the period used to derive reference points | |

| |
|---|
| Qualifying Comments |
| Red gurnard are a survey target of the West Coast South Island trawl survey and the Plenary regards the series as a reliable index of abundance. |
| Trends in CPUE indices are broadly consistent with trends in trawl survey biomass, particularly since the late 2000s, corroborating the recent increase. |
| Fishery Interactions |
| Red gurnard are primarily taken in conjunction with the following QMS species: flatfish, barracouta, stargazer, red cod, tarakihi and other species in the West Coast South Island target bottom trawl fishery. |
| Incidental captures of seabirds occur and there is a risk of incidental capture of Hector's dolphins. |

7. FOR FURTHER INFORMATION

- Beentjes, M P; MacGibbon, D; Lyon, W S (2015) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2014 (KAH1402). *New Zealand Fisheries Assessment Report 2015/14*.
- Beentjes, M.P.; MacGibbon, D.; Parkinson, D. (2016). Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2016 (KAH1605). *New Zealand Fisheries Assessment Report 2016/61*. 135 p.
- Blackwell, R (1988) Red gurnard. New Zealand Fisheries Assessment Research Document 1988/23: 18 p. (Unpublished report held by NIWA library, Wellington.)
- Boyd, R O; Reilly, J L (2002) 1999–00 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)

RED GURNARD (GUR)

- Challenger Finfisheries Management Company (2003) Report to the Adaptive Management Programme Fishery Assessment Working Group. GUR 7 Adaptive Management Proposal for the 2004–05 fishing year. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Cordue, P L (1998) Designing optimal estimators for fish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences* 55: 376–386.
- Elder, R D (1976) Studies on age and growth, reproduction and population dynamics of red gurnard, *Chelidonichthys kumu* (Lesson and Garnot), in the Hauraki Gulf, New Zealand. *Fisheries Research Bulletin* No. 12. 62 p.
- Francis, R I C C (1992) Recommendations concerning the calculation of maximum constant yield (MCY) and current annual yield (CAY). New Zealand Fisheries Assessment Research Document 1992/8. 26 p. (Unpublished report held by NIWA library, Wellington.)
- Hartill, B; Bian, R; Armiger, H; Vaughan, M; Rush, N (2007) Recreational marine harvest estimates of snapper, kahawai, and kingfish in QMA 1 in 2004–05. *New Zealand Fisheries Assessment Report 2007/26*. 44 p.
- Hartill, B; Bian, R; Davies, N M (Drafta) Review of methods used to estimate recreational harvests. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by the Ministry for Primary Industries, Wellington.) Project code: REC2004-06
- Hartill, B; Bian, R; Rush, N; Armiger, H (2013) Aerial-access recreational harvest estimates for snapper, kahawai, red gurnard, tarakihi and trevally in FMA 1 in 2011–12. *New Zealand Fisheries Assessment Report 2013/70*. 44 p.
- Kendrick, T H (2009a) Fishery characterisation and catch-per-unit-effort indices for three sub-stocks of red gurnard in GUR 1; 1989–90 to 2004–05. *New Zealand Fisheries Assessment Report 2009/10*.
- Kendrick, T H (2009b) Updated Catch-per-Unit effort indices for red gurnard in GUR 2; 1989–90 to 2004–05 *New Zealand Fisheries Assessment Report 2009/11*.
- Kendrick, T H; Bentley, N (2011) Fishery characterisations and catch-per-unit-effort indices for three sub-stocks of red gurnard in GUR 1, 1989–90 to 2008–09. *New Zealand Fisheries Assessment Report 2011/4*.
- Kendrick, T H; Bentley, (in prep. a) Updated CPUE Analyses for three substocks of red gurnard in GUR 1. Working group paper held by MPI.
- Kendrick, T H; Bentley (in prep. b) Fishery characterisation and standardised CPUE for FLA 2, GUR 2 and SNA 2. Working group paper held by MPI.
- Kendrick, T H; Bentley, N; Langley, A (2011) Report to the Challenger Fishfish Company: CPUE analyses for FMA 7 Fishstocks of gurnard, tarakihi, blue warehou, and ghost shark. (Unpublished client report held by Trophica Limited, Kaikoura).
- Kendrick, T H; Walker, N (2004) Characterisation of the GUR 2 red gurnard (*Chelidonichthys kumu*) and associated inshore trawl fisheries, 1989–90 to 2000–01. *New Zealand Fisheries Assessment Report 2004/21*. 83 p.
- Langley, A (2011) Characterisation of the Inshore Finfish fisheries of Challenger and South East coast regions (FMAs 3, 5, 7 & 8). (Unpublished client report available from <http://www.seafoodindustry.co.nz/SIFisheries>).
- Langley, A D (2014) Updated CPUE analyses for selected South Island inshore finfish stocks. *New Zealand Fisheries Assessment Report 2014/40*. 116 p.
- Langley, A.D. (2015). Fishery characterisation and Catch-Per-Unit-Effort indices for John dory in JDO 1. *New Zealand Fisheries Assessment Report 2015/47*. 76 p.
- Lydon, G J; Middleton, D A J; Starr, P J (2006) Performance of the GUR 3 Logbook Programme. AMP-WG-06/22. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Lyon, W S; Horn, P L (2011) Length and age of red gurnard (*Chelidonichthys kumu*) from trawl surveys off west coast South Island in 2003, 2005, and 2007, with comparisons to earlier surveys in the time series. *New Zealand Fisheries Assessment Report 2011/46*.
- MacGibbon, D J; Stevenson, M L (2013) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2013 (KAH1305). *New Zealand Fisheries Assessment Report 2013/66*. 115 p.
- Morrison, M A; Francis, M P; Parkinson, D M (2002) Trawl survey of the Hauraki Gulf, 2000 (KAH0012). *New Zealand Fisheries Assessment Report 2002/46*. 48 p.
- Parker, S; Fu, D (2012) Length and age structure of commercial landings of red gurnard (*Chelidonichthys kumu*) in GUR 2 in 2009–10. *New Zealand Fisheries Assessment Report 2012/35*. 36 p.
- Schofield, M I, Langley, A D; Middleton, D A (in prep.) Fisheries characterisation and catch-per-unit-effort analyses FLA 2. (Draft FAR held by MPI.)
- Starr, P J; Kendrick, T H (2012) GUR 3 Fishery Characterisation and CPUE Report. *New Zealand Fisheries Assessment Report 2013/37*. 71 p.
- Starr, P J; Kendrick, T H (in prep.) GUR 7 Fishery Characterisation and CPUE Report. Draft New Zealand Fisheries Assessment Report.
- Starr, P J; Kendrick, T H; Lydon, G J; Bentley, N (2007) Report to the Adaptive Management Fishery Assessment Working Group: Full term review of the GUR 3 Adaptive Management Programme. AMP-WG-07/11v2. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington).
- Stevenson, M L (2000) Assessment of red gurnard (*Chelidonichthys kumu*) stocks GUR 1 and GUR 2. *New Zealand Fisheries Assessment Report 2000/40*. 51 p.
- Stevenson, M L (2004) Trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2003 (KAH0304). *New Zealand Fisheries Assessment Report 2004/4*. 69 p.
- Stevenson, M L (2006) Trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2005 (KAH0503). *New Zealand Fisheries Assessment Report 2006/4*. 69 p.
- Stevenson, M L (2007) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2007 KAH0704. *New Zealand Fisheries Assessment Report 2007/41*. 64 p.
- Stevenson, M L (2009) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2009. *New Zealand Fisheries Assessment Report 2010/11*. 77 p.
- Stevenson, M L (2012) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2011. *New Zealand Fisheries Assessment Report 2012/50*. 77 p.
- Stevenson, M.L.; MacGibbon, D.J. (2015). Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2015 (KAH1503) *New Zealand Fisheries Assessment Report 2015/67*. 94 p.
- Sutton, C P (1997) Growth parameters, and estimates of mortality for red gurnard (*Chelidonichthys kumu*) from off the east and west coasts of the South Island, New Zealand. *New Zealand Fisheries Assessment Research Document 1997/1*. 15 p. (Unpublished report held by NIWA library, Wellington.)
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991–92 to 1993–94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Vignaux, M (1997) CPUE analyses for fishstocks in the adaptive management programme. *New Zealand Fisheries Assessment Research Document 1997/24*. 68 p. (Unpublished report held by NIWA library, Wellington.)
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey Of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.