


DREDGE OYSTERS (OYS 7C) – Challenger Marlborough*(Ostrea chilensis)***1. FISHERY SUMMARY**

OYS 7C encompasses an area from West Head, Tory Channel in the north to Clarence Point in the south including Cloudy Bay and Clifford Bay in the southern part of Cook Strait. OYS 7C is considered a separate fishery from OYS 7 (Golden Bay, Tasman Bay, and Marlborough Sounds) on the basis of differences in habitat and environmental parameters.

OYS 7C was introduced into the QMS on 1 October 2005 with a TAC of 5 t and a TACC of 2 t. Following a survey in April 2007, the TAC was increased to 50 t with a TACC of 43 t on 1 October 2007. In 2009, with information from CPUE and catch data, the TAC was reviewed again and resulted in a TAC increase to 72 t in October 2009 (Table 1). At the time of the review the Shellfish Working Group suggested that raising the TACC by a further 15–20 tonnes was unlikely to be detrimental to the fishery in the short-term, however without improved estimates of mortality, growth, and dredge efficiency, it was difficult to predict the effects that an increased TACC would have on the status of the fishery in the medium to long-term, and that a research strategy for improved assessment was required.

Table 1: Total Allowable Commercial Catch (TACC, t) declared for OYS 7C since introduction into the QMS in 2005.

Fishing year	TAC	TACC	Customary	Recreational	Other
2005–07	5	2	1	1	1
2007–09	50	43	1	1	5
2009 to present	72	63	1	1	7

1.1 Commercial fishery

Commercial landings for OYS 7C are reported in greenweight. The fishing year runs from 1 October to 30 September and fishers can harvest year round (there is no oyster season defined by regulations).

There is historical evidence of limited exploitation of oyster beds within Port Underwood as early as the 1800s (K. Wright pers. comm. in Drummond 1994a). Limited fishing under a special permit took place south of Tory Channel on the east coast of the South Island in 1990 and 1991.

Since 2005, landed catch has been reported via Monthly Harvest Returns (Table 2), although landings were negligible until 2007–08 when the recent commercial operation was initiated. During 2007–08 fishing took place over 30 fishing days from December to February and in 2008–09 fishing took place from January to April. Landings were at about the level of the TACC up to and including 2010–11, but were lower in recent years due to oyster grading and marketing requirements; only 3 t was landed in 2014-15 (Figure 1, Table 2).




Figure 1: Reported landings (t) and TACC for OYS 7C from 2005–06 to present.

Table 2: Reported landings (t) in the OYS 7C fishery since October 2005 (QMS). Reported catch is landed greenweight summarised from Monthly Harvest Returns.

Fishing year	TACC	Reported Landings (MHR)
2005–06	2	0.1
2006–07	2	0
2007–08	43	40.9
2008–09	43	38.2
2009–10	63	62.7
2010–11	63	62.5
2011–12	63	39.9
2012–13	63	5.9
2013–14	63	2.8
2014–15	63	3.1

1.2 Recreational fishery

The recreational catch allowance for OYS 7C is 1 tonne. The recreational daily bag limit for oysters in the Challenger fishery area is 50 per person. Oysters that cannot pass through a 58 mm internal diameter solid ring are deemed legal size. The recreational season for dredge oysters in the Challenger area is all year round. Oysters must be landed in their shells. There are no data available on the recreational catch within OYS 7C.

1.3 Customary fisheries

DREDGE OYSTER (OYS 7C)

The customary catch allowance for OYS 7C is 1 tonne. There are no data available on the customary catch.

1.4 Illegal catch

There is no quantitative information on the level of illegal catch.

1.5 Other sources of mortality

Bonamia exitiosa is a haemocytic, haplosporid parasite (infects mainly haemocytes or blood cells) of flat oysters and is known to infect *Ostrea chilensis* in New Zealand and Chile and various other species of *Ostrea* in other countries. *Bonamia* has caused catastrophic mortality in the Foveaux Strait oyster fishery and is endemic in oysters in the OYS 7 area (Hine pers. comm.). The level of mortality caused by disease is unknown.

An allowance of 7 t for Other Mortality (including incidental fishing mortality, heightened natural mortality such as disease mortality, and illegal harvest) is included in the TAC.

2. BIOLOGY

There are no biological studies of *O. chilensis* specific to the OYS 7C area. In the absence of area-specific estimates, parameters required for management purposes are based on the Foveaux Strait fishery described by Cranfield & Allen (1979) or the OYS 7 (Tasman Bay) fishery. The biology of oysters in the neighbouring area of OYS 7 (Tasman and Golden Bay) was summarised by Handley & Michael (2001), and further biological data was presented in Brown et al (2008). All this work is summarised below.

Oysters in OYS 7C (Cloudy Bay/Clifford Bay) and OYU 5 (Foveaux) both comprise rather discrete patches of oysters on a predominantly sandy substrate whereas OYS 7 (Tasman Bay) oysters tend to be more uniformly distributed at a lower density on muddy habitat. Environmental factors such as hydrodynamics, seasonal water temperature and riverine inputs differ substantially among the OYS 7, OYS 7C and OYU 5 areas and are likely to influence the biological characteristics of those oyster populations. Oysters in OYS 7C are generally more abundant and occur at higher densities than in OYS 7 (Brown & Horn 2007).

The variability in shell shapes and high variability in growth rate between individuals, between areas within the OYS 7 fishery, and between years require careful consideration in describing growth. Assuming the minimum legal size could range in diameter (1/2 length + height) from 58 mm to 65 mm, data from Drummond (1994b) indicated that Tasman Bay oysters could grow to legal size in two to three years. Modelling of limited data from Tasman Bay in Brown et al (2008) indicated that 77% of three year old oysters and 82% of 4 year old oysters would attain lengths greater than the minimum legal size of 58 mm length at the start of the fishing season. Osborne (1999) used results from a MAF Fisheries study conducted between 1990 and 1994 to construct a von Bertalanffy equation describing oyster growth in the OYS 7 fishery. Estimated biological parameters including instantaneous natural mortality (M) from Drummond (1993, 1994b) and growth parameters for von Bertalanffy equations from Osborne (1999) and from Brown et al (2008) are given in Table 3. Mortality estimates by Drummond (1994b) and growth parameters in Osborne (1999) were derived from a tagging study conducted in Tasman Bay between 1990 and 1992 (Drummond 1993). von Bertalanffy growth parameters in Brown et al (2008) were estimated based on a limited data set from enhanced habitat experiments, and describe growth of young oysters. Estimates of M based on experimental data from Foveaux and Tasman Bay ranged from 0.042 (Dunn et al 1998b) to 0.92 (Drummond et al 1994a). However, after some discussion the Shellfish Working Group concluded that those figures were not realistic, and that M was more likely to lie between 0.1 and 0.3.

Table 3: Estimated biological parameters for oysters in OYS 7 and OYU 5. In the absence of data specific to OYS 7C these estimates are used for management purposes in OYS 7C. [Continued on next page]

1. Natural Mortality (M)

Area	Estimate	Source
Tasman Bay	0.920	Drummond (1994b)
Tasman Bay	0.200	Drummond (1993)
Foveaux Strait	0.042	Dunn et al (1998b)
Foveaux Strait	0.100	Allen (1979)

Table 3 [Continued]2. von Bertalanffy growth (change in diameter mm) parameter estimates from OYS 7 t_0 not provided by Osborne (1999).

K	L_{inf}	t_0	Source
0.597	85.43	-	Osborne (1999)
0.99 +/- 0.16 (sd)	67.52	0.11	Brown et al (2008)

3. STOCKS AND AREAS

Fishing within OYS 7C has been limited to two discrete areas; one in parts of Clifford and Cloudy Bays and the other immediately south of Tory Channel, and commercial oyster fishing has not extended south of Cape Campbell. The oyster population in OYS 7C is likely to be biologically isolated from populations in Foveaux Strait (OYU 5) and at the Chatham Islands (OYS 4) on the basis of geographical distance. The populations in OYS 7C and OYS 7 could also be biologically distinct due to their geographical separation which quite likely leads to limited dispersal of larvae between the two areas.

4. STOCK ASSESSMENT**4.1 Estimates of fishery parameters and abundance**

A survey of OYS 7C was carried out in 2007 (Brown & Horn 2007) and estimates of the number of recruits (oysters unable to pass through a 58 mm ring) and pre-recruits (less than 58 mm) from Clifford and Cloudy Bay are given in Table 4. Dredge efficiency was assumed to be 100% for the purposes of the survey.

Table 4: Estimate of number of recruit and pre-recruit oysters from Brown & Horn (2007).

Year	Area (Ha)	Recruit No.		Pre-recruit No.	
		Estimate	CV %	Estimate	CV %
2007	43 709	19.5 million	19	14 million	19

4.2 Biomass estimates

Estimates of recruited biomass, from the 2007 survey are given in Table 5.

Table 5: Estimate of relative recruited (≥ 58 mm) oyster biomass (t greenweight) in OYS 7C (Brown & Horn 2007).

Year	Area (Ha)	Biomass (t)	CV
2007	43 709	1 778	0.19

4.3 Yield estimates and projections

For new fisheries where there are insufficient data to conduct a yield per recruit analysis, yield can be estimated using the formula from Mace (1988) recommended by the Ministry of Fisheries Science Group (MFish 2008) for calculation of Maximum Constant Yield (MCY).

$$MCY = 0.25MB_0$$

DREDGE OYSTER (OYS 7C)

Where B_0 is an estimate of virgin recruited biomass (here assumed to be equal to the recruited biomass estimate from the 2007 survey (1778 t, Brown & Horn 2007) divided by dredge efficiency) and M is an estimate of natural mortality. A range of MCY estimates are given in Table 6 using values for dredge efficiency of 100% and 64% (Bull 1989), and values for M ranging from 0.1 to 0.3 taken from studies conducted in the Foveaux and Nelson-Marlborough oyster fisheries.

Table 6: Estimates of MCY for M of 0.1–0.3. MCY 1 was estimated using a dredge efficiency of 64% from Bull (1989) and MCY 2 was estimated assuming a dredge efficiency of 100%.

M	MCY 1	MCY 2
0.1	69	44
0.2	139	89
0.3	208	133

There are no CAY estimates for OYS 7C.

4.4 Other Yield Estimates

There are no other yield estimates for OYS 7C.

4.5 Other Factors

Dredging for oysters will have an impact on the soft sediment habitats within Cloudy and Clifford Bays, and will affect both the dredge oyster beds and other species found in association with these beds. In addition, various areas within the fishery (mainly around coastal rocky reefs) are understood to support a range of sensitive invertebrate species including soft corals, large erect and divaricating bryozoans, starfish, horse mussels, and crabs. The impacts of dredging are likely to be more severe on these habitats than on soft sediments, and will increase with increasing fishing effort, but there is insufficient information to quantify the degree of impact under any given TAC. There may be some overlap with other fisheries that contact the bottom in this area, but this has not been quantified.

Industry has proposed to voluntarily restrict fishing to two discrete areas to mitigate the effects of fishing. These areas are where oyster densities are highest. By-catch of benthic invertebrates was collected during the biomass survey and could be analysed to help to determine the distribution of sensitive habitats.

5. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was reviewed by the Aquatic Environment Working Group for inclusion in the Fishery Assessment Plenary November 2014. A broader summary of information on a range of issues related to the environmental effects of fishing and aspects of the marine environment and biodiversity of relevance to fish and fisheries is available in the Aquatic Environment & Biodiversity Annual Review (Ministry for Primary Industries 2013a).

5.1 Role in the ecosystem

Dredge oysters (*Ostrea chilensis*) are benthic, epifaunal, sessile bivalve molluscs that have a relatively limited pelagic larval dispersal phase. They are patchily distributed around the New Zealand coast on a variety of substrates (biogenic reef, gravel, sand, mud) in intertidal to subtidal inshore waters, commonly in depths of up to 60 m or more. Commercially exploited beds of oysters occur in Foveaux Strait (OYU 5), Tasman Bay (OYS 7), and Cloudy and Clifford Bays (OYS 7C). Beds at the Chatham Islands (OYS 4) have potential for commercial exploitation. Oysters play important roles in the ecosystem that include influencing water quality by filtering phytoplankton and other suspended particles from the seawater, linking primary production with higher trophic levels, and acting as ecosystem engineers by stabilising sediments and providing

structural habitat (biogenic reef) for other taxa (e.g., algae, ascidians, bryozoans, sponges, echinoderms, worms, molluscs, crustaceans, fish).

5.1.1 Trophic interactions

Oysters are active suspension feeders, consuming phytoplankton suspended in the water column. Their diet is the same as or similar to that of many other suspension feeding taxa, including other bivalves such as scallops, clams, and mussels. Oysters are probably prey for a wide range of invertebrate and fish predators, but published records of known or suspected predators are limited. Reported invertebrate predators of *O. chilensis* include brittlestars (*Ophiopsammus maculata*) (Stead 1971), starfish (*Coscinasterias calamaria* and *Astrostele scabra*) (Cranfield 1979) and flatworms (*Enterogonia orbicularis*) (Handley 2002); suspected invertebrate predators include octopus (*Pinnoctopus cordiformis*) and shell boring gastropods (*Poirieria zelandica*, *Xymeme ambiguous*, and *Xymenella pusillis*) (Brown 2012). Predators of oysters probably change with oyster size. Most mortality of oyster spat (small juveniles) during their first winter appears to result from predation by polychaetes, crabs, and gastropods (Ministry for Primary Industries 2013b).

5.2 Incidental catch (fish and invertebrates)

A range of non-target fish and invertebrate species are caught and discarded by dredge fisheries for *O. chilensis*. No data are available on the level or effect of this incidental catch (bycatch) and discarding by the fisheries. Invertebrate bycatch data are available from dredge surveys of the oyster stocks, and the bycatch of the fisheries is likely to be similar to that of the survey tows conducted in areas that support commercial fishing. Fish bycatch data are generally not recorded on surveys, presumably because fish constitute a small fraction of the total bycatch.

In OYU 5 (Foveaux Strait), Cranfield et al (1999) summarised the results of Fleming (1952) who sampled the macrofaunal bycatch of oyster fishing in a ‘near virgin’ area of the fishery in 1950; the bycatch was dominated by the frame-building bryozoan *Cinctipora elegans* (and oysters *O. chilensis*) and included a diverse range of other epifaunal organisms. More recently, presence-absence data on the bycatch of oyster dredging have been recorded during surveys and in fishers’ logbooks (Michael 2007). In a specific study of the benthic macrofauna bycatch of the 2001 oyster dredge survey in Foveaux Strait, Rowden et al (2007) identified at least 190 putative species representing 82 families and 12 phyla; ‘Commercial’ survey strata were principally characterised by the families Balanidae (barnacles), Mytilidae (mussels), Ophiodermatidae (brittle stars), Ostreidae (oysters), and Pyuridae (tunicates). For the 2007 survey of OYU 5, Michael (2007) listed the percentage occurrence of sessile and motile species caught as bycatch in the survey dredge tows. The five most commonly caught sessile species (excluding oysters) were hairy mussels *Modiolus areolatus* (80% occurrence), barnacles *Balanus* sp. (61%), kina *Evechinus chloroticus* (61%), nesting mussels *Modiolarca impacta* (53%), and ascidians *Pyura pulla* (51%). The five most commonly occurring motile bycatch species were brittlestars *Ophiopsammus maculata* (90% occurrence), circular saw shells (gastropods) *Astraea heliotropium* (80%), hermit crabs *Pagurus novizelandiae* (80%), eight armed starfish *Coscinasterias muricata* (63%), and brown dipple starfish *Pentagonaster pulchellus* (54%). Common bycatch species of oyster dredge surveys in Foveaux Strait were reported by Michael (2007) and are listed below in Table 7.

Table 7: Invertebrate species commonly caught as bycatch in dredge surveys of oysters (*O. chilensis*) in Foveaux Strait. Sourced from Michael (2007). [Continued on next page]

Type	Species
Infaunal bivalves	<i>Glycymeris modesta</i> (small dog cockle), <i>Tawera spissa</i> (morning star shell), <i>Tucetona laticostata</i> (large dog cockle), <i>Pseudoxyperas elongata</i> (‘tuatua’), <i>Venericardia purpurata</i> (purple cockle)
Epifaunal bivalves	<i>Modiolus areolatus</i> (hairy mussel), <i>Modiolarca impacta</i> (nesting mussel), <i>Aulacomya atra maoriana</i> (ribbed mussel), <i>Barbatia novaezelandiae</i> (ark shell), <i>Pecten novaezelandiae</i> (scallop), <i>Chlamys zelandiae</i> (lions paw scallop), <i>Neothyris lenticularis</i> (large lantern shell), <i>N. compressa</i>

DREDGE OYSTER (OYS 7C)

Table 7 [Continued]

	(compressed lantern shell)
Sponges	<i>Chondropsis topsentii</i> (cream sponge), <i>Crella incrustans</i> (red-orange sponge), <i>Dactylia palmata</i> (finger sponge)
Ascidians	<i>Pyura pachydermatina</i> (kaeo), <i>P. pulla</i>
Bryozoans	<i>Celleporaria agglutinans</i> (hard/plate coral), <i>Cinctipora elegans</i> (reef-building bryozoan), <i>Horera foliacea</i> (lace coral), <i>Hippomenella vellicata</i> (paper coral), <i>Tetrocycloecia neozelanica</i> (staghorn coral), <i>Orthoscuticella fusiformis</i> (soft orange bryozoan)
Barnacles and chitons	<i>Balanus decorus</i> (large pink barnacle), <i>Cryptochonchus porosus</i> (butterfly chiton), <i>Eudoxochiton nobilis</i> (noble chiton), <i>Rhyssoplax canaliculata</i> (pink chiton)
Starfish, brittlestars, and holothurians	<i>Coscinasterias muricata</i> (eight armed starfish), <i>Pentagonaster pulchellus</i> (brown dipple starfish), <i>Ophiosammus maculata</i> (snaketail brittlestar), <i>Australostichopus mollis</i> (sea cucumber)
Crabs	<i>Pagurus novaezelandiae</i> (hermit crab), <i>Eurynolambrus australis</i> (triangle crab), <i>Metacarcinus novaezelandiae</i> (cancer crab), <i>Nectocarcinus</i> sp. (red crab)
Urchins	<i>Evechinus chloroticus</i> (kina), <i>Apatopygus recens</i> (heart urchin), <i>Goniocidaris umbraculum</i> (coarse-spined urchin), <i>Pseudechinus novaezelandiae</i> (green urchin), <i>P. huttoni</i> (white urchin), <i>P. albocinctus</i> (red urchin)
Gastropods	<i>Astraea heliotropium</i> (circular saw shell), <i>Alcithoe arabica</i> (volute), <i>Argobuccinum pustulosum tumidum</i> , <i>Turbo granosus</i> , <i>Cabestana spengleri</i> , <i>Charonia lampras</i>
Octopuses	<i>Pinnoctopus cordiformis</i> (common octopus), <i>Octopus huttoni</i> (small octopus)

In OYS 7 (Tasman/Golden Bays), data on the bycatch of the 1994–2014 dredge surveys have been collected but not analysed, except for preliminary estimation of the 1998–2013 bycatch trajectories (Williams et al 2014b). The surveys record the bycatch of other target species of scallops (*Pecten novaezelandiae*) and green-lipped mussels (*Perna canaliculus*), and various other non-target bycatch in nine categories (Williams et al 2014b). Observation of the 2014 survey sampling identified a problem with the way these categorical bycatch data have been recorded which limits their utility (Williams et al 2014a).

In OYS 7C, a dredge survey of oysters in Cloudy and Clifford Bays was conducted in 2006, and the survey skipper recorded qualitative comments on the bycatch of each tow, which included ‘coral’, ‘sticks and seaweed’, shells, volutes, ‘red weed’, horse mussels, shell with worm, small crabs, mussels, and scallop (Brown & Horn 2006).

In OYS 4 (Chatham Islands), data on the bycatch of a 2013 dredge survey of oysters off the north coast of Chatham Island were recorded (as estimated volumes of different bycatch categories) but not analysed (Williams et al 2013).

5.3 Incidental catch (seabirds, mammals, and protected fish)

There is no known bycatch of seabirds, mammals or protected fish species from *O. chilensis* oyster fisheries.

5.4 Benthic interactions

There are a variety of benthic habitats in the different oyster fisheries areas, which generally occur either on coarse substrates usually found in areas of high natural disturbance (Foveaux Strait, Cloudy/Clifford Bays and the Chatham Islands) or on fine substrates typical of sheltered areas (Tasman Bay). Benthic habitats within the Foveaux Strait oyster fishery area were classified by Michael (2007) and comprise a variety of sand/gravel/shell flats and waves, rocky patch reef, and biogenic areas. Cranfield et al (1999) referred to the latter as epifaunal reefs that he defined as “tidally-oriented, linear aggregations of patch reefs formed by the bryozoan *Cinctipora elegans*, cemented by encrusting bryozoans, ascidians, sponges and polychaetes”. Cranfield et al’s papers (Cranfield et al 1999, Cranfield et al 2001, Cranfield et al 2003) suggested that epifaunal reefs are oyster habitat, but Michael’s reports (Michael 2007, 2010) state that commercial fishing for

oysters is mainly based on sand, gravel, and shell habitats with little epifauna. In Foveaux Strait, commercial oyster dredging occurs within an area of about 1000 km² (although only a portion of this is dredged each year), which is about one-third of the overall OYU 5 stock area (Michael 2010). Habitats within the Cloudy/Clifford Bays and the Chatham Islands fisheries areas have not been defined. The benthic habitat within the Tasman Bay oyster fishery area is predominately mud, although to some extent this may have been affected by land-based sedimentation into the bay and homogenisation of the substrate by dredging and trawling (Brown 2012).

It is well known that fishing with mobile bottom contact gears such as dredges has impacts on benthic populations, communities, and their habitats (e.g., see Kaiser et al 2006, Rice 2006). The effects are not uniform, but depend on at least: “the specific features of the seafloor habitats, including the natural disturbance regime; the species present; the type of gear used, the methods and timing of deployment of the gear, and the frequency with which a site is impacted by specific gears; and the history of human activities, especially past fishing, in the area of concern” (Department of Fisheries and Oceans 2006). In New Zealand, the effects of oyster dredging on the benthos have been studied in Foveaux Strait (OYU 5) (Cranfield et al 1999, Cranfield et al 2001, Cranfield et al 2003, Michael 2007) and Tasman/Golden Bays (OYS 7) (Tuck et al 2011). The results of these studies are summarised in the Aquatic Environment & Biodiversity Annual Review (Ministry for Primary Industries 2013a), and are consistent with the global literature: generally, with increasing fishing intensity there are decreases in the density and diversity of benthic communities and, especially, the density of emergent epifauna that provide structured habitat for other fauna.

The effects of dredging (Ministry for Primary Industries 2013a) may be more severe in sheltered areas (e.g., Tasman Bay) than in exposed areas (e.g., Foveaux Strait, Cloudy/Clifford Bays, Chatham Islands). Dredging damages epifauna, and erect, structured habitats, such as biogenic/epifaunal reefs, are the most sensitive to dredging disturbance. Dredging destabilises sediment/shell substrates, suspends sediments and increases water turbidity; the sensitivity of habitats to suspended sediments and their deposition probably varies depending on the prevailing natural flow regime, being greater in muddy sheltered areas than in high flow environments. Habitats disturbed by dredging tend to become simpler, more homogenous areas typically dominated by opportunistic species. Dredging generally results in reduced habitat structure and the loss of long-lived species.

For studies of the effects of oyster dredging in Foveaux Strait, interpretation of the authors differ (Ministry for Primary Industries 2013a). Cranfield et al’s papers (Cranfield et al 1999, Cranfield et al 2001, Cranfield et al 2003) concluded that dredging biogenic reefs for their oysters damages their structure, removes epifauna, and exposes associated sediments to resuspension such that, by 1998, none of the original bryozoan reefs remained. Michael (2007) concluded that there are no experimental estimates of the effect of dredging in the strait or on the cumulative effects of fishing or regeneration, that environmental drivers should be included in any assessment, and that the previous conclusions cannot be supported. The authors agree that biogenic bycatch in the fishery has declined over time in regularly-fished areas, that there may have been a reduction in biogenic reefs in the strait since the 1970s, and that simple biogenic reefs appear able to regenerate in areas that are no longer fished (dominated by byssally attached mussels or reef-building bryozoans). There is no consensus that reefs in Foveaux Strait were (or were not) extensive or dominated by the bryozoan *Cinctipora*.

Some areas of the Foveaux Strait (OYU 5) oyster fishery are also commercially fished (potted) for blue cod (*Parapercis colias*), and Cranfield et al (2001) presented some evidence to suggest that dredged benthic habitats and blue cod densities regenerated in the absence of oyster dredging. Bottom trawling also occurs within the OYU 5 area, but there is little overlap with the main areas fished for oysters. In OYS 7, other benthic fisheries (e.g. bottom trawl, scallop, green-lipped mussel) occur and probably also interact with oysters and their habitats.

5.5 Other considerations

5.5.1 Spawning disruption

Fishing during spawning may disrupt spawning activity or success. In the Foveaux Strait fishery, the traditional harvesting period (1 March to 31 August) occurs after the main spring and summer peaks in oyster spawning activity (Jeffer & Hickman 2000). Fishing-induced damage to oysters incurred during the period before spawning could interrupt gamete maturation. Oyster fishing also targets high-density beds of oysters, which are disproportionately more important for fertilisation success during spawning.

5.5.2 Habitat of particular significance for fisheries management

None currently identified.

6. STOCK STATUS

Stock Structure Assumptions

Current management assumes that the OYS 7C oyster fishery is separate from the other oyster fisheries (i.e., Challenger (OYS 7), Foveaux Strait (OYS 5), and the Chatham Islands (OYS 4)). The stock structure of OYS 7C is assumed to be a single biological stock. Survey data show that oysters are patchily distributed in the commercial fishery area of OYS 7C and it has been suggested that the oyster populations may be mainly self-recruiting.

Stock Status	
Year of Most Recent Assessment	2009
Reference Points	Target: Default = 40% B_0 , with at least a 50% probability of achieving the target Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: F_{MSY}
Status in relation to Target	Very Likely (> 90%) to be at or above the target
Status in relation to Limits	Based on annual commercial oyster removals of less than 4% of the estimated 2007 stock size, the status is likely to be close to virgin size and is Very Unlikely (< 10%) to be below the soft and hard limits.
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring.
Historical Stock Status Trajectory and Current Status	
OYS7C	
<p>Estimated relative biomass (t greenweight) of recruited oysters (≥ 58 mm) (green point and error bars denoting CV), TACC (solid red line), and reported landings (blue line) in t greenweight since 1998. The biomass estimate is from a 2007 survey and is uncorrected for dredge efficiency. Landings data from MHRs. Fishing year beginning 2005-06 to 2014-15</p>	

Fishery and Stock Trends	
Recent trend in Biomass or Proxy	Only one biomass survey has been conducted, in 2007, from which the recruited biomass was estimated to be 1778 t (assuming 100% dredge efficiency).
Recent trend in Fishing Intensity or Proxy	The OYS 7C commercial fishery got underway in 2007–08; in that fishing year the exploitation rate was an estimated at 0.02 (assuming 100% dredge efficiency).
Other Abundance Indices	-
Trends in Other Relevant Indicator or Variables	Landings were at about the level of the TACC up to and including 2010–11, but were lower in recent years due to oyster grading and marketing requirements.

Projections and Prognosis	
Stock Projections or Prognosis	Quantitative stock projections are unavailable
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (<10%)
Assessment Methodology and Evaluation	
Assessment Type	Level 2: Partial Quantitative Stock Assessment
Assessment Method	Yields are estimated as a proportion of the survey biomass for a range of assumed values of natural mortality and dredge efficiency.
Assessment Dates	Latest assessment: 2009 Next assessment: Unknown
Overall Assessment Quality Rank	1 - High Quality
Main data inputs (rank)	Biomass survey: 2007 1 – High Quality
Period of Assessment	Latest assessment: 2009 Next assessment: Unknown
Data not used (rank)	N/A
Changes to Model Structure and Assumptions	
Major Sources of Uncertainty	There has been only a single biomass survey of this fishstock and repeat surveys should be scheduled at regular intervals. Natural mortality (M) and dredge efficiency are poorly known but are integral parameters of the method used to estimate yield. There is also major uncertainty about the response of localised populations to fishing.
Qualifying Comments	
Some of the surveyed area was not actively fished up to 2009. There are areas of potential oyster habitat which are not fished due to sanitation concerns and substrate which is marginal for fishing. In 2009, the Shellfish FAWG was asked to evaluate the implications of raising the TACC (of 50 t) by 15–20 t. In 2009 it was considered Very Unlikely (< 10%) that an increase in the TACC of this amount would cause the biomass to decline below the Soft Limit in the next 3 to 5 years. On 1 October 2009 the TACC was changed to 63 t.	

Fishery Interactions
There may be some overlap with other fisheries that contact the bottom in this area, but this has not been quantified.

7. FOR FURTHER INFORMATION

- Allen, R L (1979) A yield model for the Foveaux Strait oyster (*Ostrea lutaria*) fishery. *Rapports et Proces-verbaux des reunions. Conseil Permanent International pour l'Exploration de la mer* (175): 70–79A.
- Brown, S; Campbell, M; Handley, S; Michael, K; Cairney, D; Bradley, A; Moremede, S (2008) Challenger oyster enhancement research programme. NIWA Client Report NEL2008-032. 77 p.
- Brown, S; Horn, P (2006) Preliminary estimate of dredge oyster abundance in Cloudy Bay. NIWA Client Report NEL2006-012 prepared for Talleys Fisheries Ltd. NIWA project TAL07401. 5 p. (Unpublished report held by NIWA, Nelson.)
- Brown, S; Horn, P L (2007) Survey of scallops and oysters in Golden Bay, Tasman Bay, and the Marlborough Sounds, in May 2007. NIWA Client Report NEL2007-012. 41 p.
- Brown, S N (2012) Ecology and enhancement of the flat oyster *Ostrea chilensis* (Philippi, 1845) in central New Zealand. Unpublished PhD thesis. University of Canterbury, Christchurch, New Zealand. 218 p.
- Bull, M F (1989) Dredge Oyster Survey, Tasman Bay. October–November, 1989. MAFFish Central Region unpublished report. (Unpublished report held by Ministry for Primary Industries.)
- Cranfield, H J (1979) The biology of the oyster, *Ostrea lutaria*, and the oyster fishery of Foveaux Strait. *Rapports et Proces-verbaux des reunions. Conseil Permanent International pour l'Exploration de la mer* 175: 44–49.
- Cranfield, H J; Allen, R L (1979) Mark-recapture surveys of the Foveaux Strait dredge oyster (*Ostrea lutaria*) population. *Rapports et Proces-verbaux des reunions. Conseil Permanent International pour l'Exploration de la Mer* (175): 63–69.
- Cranfield, H J; Carbines, G; Michael, K P; Dunn, A; Stotter, D R; Smith, D J (2001) Promising signs of regeneration of blue cod and oyster habitat changed by dredging in Foveaux Strait, southern New Zealand. *New Zealand Journal of Marine and Freshwater Research* 35: 897–908.
- Cranfield, H J; Manighetti, B; Michael, K P; Hill, A (2003) Effects of oyster dredging on the distribution of bryozoans biogenic reefs and associated sediments in Foveaux Strait, southern New Zealand. *Continental Shelf Research* 23: 1337–1357.
- Cranfield, H J; Michael, K P; Doonan, I J (1999) Changes in the distribution of epifaunal reefs and oysters during 130 years of dredging for oysters in Foveaux Strait, southern New Zealand. *Aquatic Conservation of Marine and Freshwater Ecosystems* 9: 461–483.
- Department of Fisheries and Oceans. (2006) Impacts of trawl gear and scallop dredges on benthic habitats, populations and communities. *DFO Canadian Science Advisory Secretariat Science Advisory Report 2006/025*. 13 p.
- Drummond, K (1987) The Nelson Marlborough dredge oyster (*Tiostrea lutaria*) fishery. MAF Fisheries. (Unpublished report held by the Ministry for Primary Industries.)
- Drummond, K (1994a) The Challenger dredge oyster *Tiostrea chilensis* fishery. (Unpublished draft FARD held by Ministry for Primary Industries.)
- Drummond, K L (1994b) Report on Tasman Bay dredge oyster productivity trials. MAF Fisheries Central Region Internal Report.23. (Unpublished report held by Ministry for Primary Industries.)
- Drummond, K; Bull, M (1993) Oyster enhancement research trials. MAF Fisheries internal memorandum, 18 August 1993. (Unpublished report held by Ministry for Primary Industries.)
- Dunn, A; Cranfield, H J; Michael, K P (1998b) Estimates of natural mortality of the Foveaux Strait oyster (*Tiostrea chilensis*) between 1974 and 1986. Final Research Report to the Ministry of Fisheries for Project OYS9801. 12 p. (Unpublished report held by the Ministry for Primary Industries, Wellington.)
- Dunn, A; Doonan, I J D; Cranfield, H J; Michael, K P; Stotter, D R (1998) Modelling Growth of the Foveaux Strait Oyster (*Tiostrea chilensis*) Final research report for Ministry of Fisheries Research Project OYS9701. 16 p. (Unpublished report held by the Ministry for Primary Industries.)
- Fleming, C A (1952) A Foveaux Strait oyster bed. *New Zealand Journal of Science and Technology* 34B: 73–85.
- Handley, S; Michael, K (2001) Research plan for the Challenger dredge oyster fishery: A summary of existing knowledge and options for further research. NIWA client report no. CSE01402. 42 p.
- Handley, S J (2002) Optimizing intertidal Pacific oyster (Thunberg) culture, Houhora Harbour, northern New Zealand. *Aquaculture Research* 33: 1019–1030.
- Jeffs, A; Hickman, R W (2000) Reproductive activity in a pre-epizootic wild population of the Chilean oyster, *Ostrea chilensis*, from southern New Zealand. *Aquaculture* 183: 241–253.
- Kaiser, M J; Clarke, K R; Hinz, H; Austen, M C V; Somerfield, P J; Karakassis, I (2006) Global analysis of the response and recovery of benthic biota to fishing. *Marine Ecology Progress Series* 311: 1–14.
- Mace, P M (1988) The relevance of MSY and other biological reference points to stock assessments in New Zealand. New Zealand Fisheries Assessment Research Document 1988/30. (Unpublished document held at NIWA library, Wellington.)
- Michael, K P (2007) Summary of information in support of the Foveaux Strait Oyster Fishery Plan: the Foveaux Strait ecosystem and effects of oyster dredging. Final Research Report for Ministry of Fisheries project ZBD2005-04. 103 p. (Unpublished report held by the Ministry for Primary Industries, Wellington.)
- Michael, K P (2010) Summary of information from Foveaux Strait oyster (*Ostrea chilensis*, OYU 5) strategic research 2000–09: context for the 2010 strategic research plan. *New Zealand Fisheries Assessment Report 2010/20*. 37 p.
- Ministry of Fisheries Science Group (2008) Report from the Fishery Assessment Plenary, May 2008: stock assessments and yield estimates. Part 1 Wellington, Ministry of Fisheries. 348 p.
- Ministry for Primary Industries (2013a) Aquatic Environment and Biodiversity Annual Review 2013. Compiled by the Fisheries Management Science Team, Ministry for Primary Industries, Wellington, New Zealand. 538 p. Available from: www.mpi.govt.nz/news-resources/publications
- Ministry for Primary Industries (2013b) Fisheries Assessment Plenary, November 2013: stock assessments and yield estimates. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 611 p.
- Osborne, T A (1998) Relative Biomass of Oysters in Golden and Tasman Bays, 1998. Osborne Research Company Ltd. Client Report ORCO Report 1003.
- Osborne, T A (1999) Nelson Dredge Oyster Stock Assessment. Osborne Research Company Ltd. Client Report ORCO Report 7.
- Rice, J (2006) Impacts of mobile bottom gears on seafloor habitats, species, and communities: a review and synthesis of selected international reviews. *Canadian Science Advisory Secretariat Research Document 2006/057*. 35 p.
- Rowden, A A; Cranfield, H J; Mann, P J; Wood, A C L (2007) Benthic macrofauna bycatch of oyster dredging in Foveaux Strait. *New Zealand Fisheries Assessment Report 2007/10*. 26 p.
- Stead, D H (1971) Observations on the biology and ecology of the Foveaux Strait dredge oyster (*Ostrea lutaria*, Hutton). *Fisheries Technical Report, New Zealand Marine Department* 68. 49 p.

- Stead, D H (1976) The Dredge Fishery for Scallops, Mussels, and Oysters in Golden Bay and Tasman Bay: Survey Results 1969–1975. Fisheries Management Division, Ministry of Agriculture and Fisheries, New Zealand.
- Sullivan, K J; Mace, P M; Smith, N W; Griffiths, M H; Todd, P R; Livingston, M E; Harley, S J; Key, J M; Connell, A M (2005) Report from the Fishery Assessment Plenary, May 2005: stock assessments and yield estimates. Wellington, Ministry of Fisheries. 364 p.
- Tuck, I; Hewitt, J; Handley, S; Willis, T; Carter, M; Hadfield, M; Gorman, R; Cairney, D; Brown, S; Palmer, A (2011) Assessing the effects of fishing on soft sediment habitat, fauna and processes. Progress Report for Ministry of Fisheries project BEN2007-01. 30 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Tunbridge, B R (1962) Occurrence and distribution of the dredge oyster (*Ostrea sinuata*) in Tasman and Golden Bays. *Fisheries Technical Report* 6. 42 p.
- Williams, J R; Bian, R; Williams, C L (2014a) Survey of scallops in Golden Bay, Tasman Bay, and Marlborough Sounds, 2014. NIWA Client Report AKL2014-027 prepared for Challenger Scallop Enhancement Company Ltd. NIWA Project CSE14301. 63 p. (Unpublished report held by NIWA, Auckland.)
- Williams, J R; Hartill, B; Bian, R; Williams, C L (2014b) Review of the Southern scallop fishery (SCA 7). *New Zealand Fisheries Assessment Report 2014/07*. 71 p.
- Williams, J R; Tuck, I D; Williams, C L (2013) "Chatham Islands dredge oyster survey, March 2013." Presented at the MPI Shellfish Fisheries Assessment Working Group, Wellington, 14 May 2013 for NIWA Project FIPR1301.