


HORSE MUSSEL (HOR)

(*Atrina zelandica*)
Kukuroroa, Kupa, Hururoa

**1. FISHERY SUMMARY****1.1 Commercial fisheries**

Horse mussels (*Atrina zelandica*) were introduced into the Quota Management System on 1 April 2004, with a combined TAC of 105 t and TACC of 29 t. Customary non-commercial and recreational allowances are 9 t each, and 58 t was allowed for other sources of mortality. The fishing year is from 1 April to 31 March and commercial catches are measured in greenweight. TACCs have been allocated in HOR 1–HOR 9. Most reported landings have been from HOR 1, and, apart from 1994–95 and 2002–03 when catches of about 5 t and 7 t respectively were reported, reported landings have all been small (Table 1). About 90% of the catch is taken as a bycatch during bottom trawling and the remainder is taken as a bycatch of dredge and Danish seine. It is likely that there is a reasonably high level of unreported discarded horse mussel catch.

1.2 Recreational fisheries

A. zelandica do not appear in records from recreational fishing surveys (Bradford 1998) but are nevertheless taken from time to time by recreational fishers. There are no estimates of recreational take for this species.

1.3 Customary non-commercial fisheries

A traditional food of Māori, this mussel is probably under-represented in midden shell counts because of the fragile and short-lived nature of the shell.

Māori customary fishers can utilise the provisions under both the recreational fishing regulations and the various customary regulations. Tangata whenua can harvest horse mussels under their recreational allowance and these are not included in records of customary catch. Customary reporting requirements vary around the country. Customary fishing authorisations issued in the South Island and Stewart Island would be under the Fisheries (South Island Customary Fishing) Regulations 1999. Many rohe moana / areas of the coastline in the North Island and Chatham Islands are gazetted under the Fisheries (Kaimoana Customary Fishing) Regulations 1998 which require reporting on authorisations. In the areas not gazetted, customary fishing permits would be issued would be under the Fisheries (Amateur Fishing) Regulations 2013, where there is no requirement to report catch.

HORSE MUSSEL (HOR)

Table 1: TACCs and reported landings (t) of horse mussel by Fishstock from 1990–91 to present from CELR and CLR data. There have never been any reported landings in HOR 4, 5, 6, or 8. These fishstocks each have a TACC of 1 t and are not reported in here.

Fishstock	HOR 1		HOR 2		HOR 3		HOR 7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1990–91	0.834	–	0	–	0	–	0	–
1991–92	0	–	0	–	0	–	0	–
1992–93	0	–	0	–	0	–	0	–
1993–94	0.003	–	0	–	0.016	–	0	–
1994–95	5.525	–	0	–	0	–	0	–
1995–96	0	–	0.019	–	0	–	0	–
1996–97	0.024	–	0	–	0	–	0	–
1997–98	0	–	0	–	0	–	0	–
1998–99	0	–	0	–	0	–	0	–
1999–00	0	–	0	–	0	–	0.81	–
2000–01	0	–	0	–	0	–	0.128	–
2001–02	0	–	0.002	–	0	–	0	–
2002–03	7.153	–	0	–	0	–	0	–
2003–04	0.026	4	0	2	0	2	0	16
2004–05	0.217	4	0	2	0	2	1.017	16
2005–06	0.026	4	0	2	0	2	0	16
2006–07	0	4	0	2	0	2	0.06	16
2007–08	0	4	0	2	0	2	0.451	16
2008–09	0.068	4	0	2	0	2	0	16
2009–10	0.289	4	0	2	0	2	0.112	16
2010–11	0	4	0	2	0	2	0.857	16
2011–12	0	4	0	2	0	2	0.605	16
2012–13	0	4	0	2	0	2	0	16
2013–14	0	4	0	2	0	2	0.214	16
2014–15	0	4	0	2	0	2	0.117	16
2015–16	0	4	0	2	0.005	2	0.380	16
2016–17	0	4	0	2	0.018	2	0.630	16
2017–18	0	4	0	2	0.018	2	0.211	16
2018–19	0	4	0	2	0.090	2	0	16
2019–20	0	4	0	2	0.500	2	0	16
2020–21	0	4	0	2	0	2	0.03	16

Fishstock	HOR 9		Total	
	Landings	TACC	Landings	TACC
1990–91	0	–	0.834	–
1991–92	0	–	0	–
1992–93	0	–	0	–
1993–94	0	–	0.019	–
1994–95	0	–	5.525	–
1995–96	0	–	0.019	–
1996–97	0	–	0.024	–
1997–98	0	–	0.128	–
1998–99	0	–	0	–
1999–00	0	–	0.1	–
2000–01	0	–	0.128	–
2001–02	0	–	0	–
2002–03	0	–	7.155	–
2003–04	0	1	0.026	29
2004–05	0.065	1	1.299	29
2005–06	0.942	1	0.968	29
2006–07	0.261	1	0.321	29
2007–08	0	1	0.451	29
2008–09	0	1	0.068	29
2009–10	0	1	0.401	29
2010–11	0	1	1	29
2011–12	0	1	0.605	29
2012–13	0	1	0	29
2013–14	0	1	0.214	29
2014–15	0	1	0.117	29
2015–16	0	1	0.385	29
2016–17	0	1	0.0648	29
2017–18	0	1	0.329	29
2018–19	0	1	0.090	29
2019–20	0	1	0.500	29
2020–21	0	1	0.030	29

The information on Māori customary harvest under the provisions made for customary fishing can be limited (Table 2). These numbers are likely to be an underestimate of customary harvest as only the catch approved and harvested in kilograms and numbers are reported in the table.

Table 2: Fisheries New Zealand records of customary harvest of horse mussel (approved and reported as weight (kg) and in numbers) since 2005–06. – no data.

Stock	Fishing year	Weight (kg)		Numbers	
		Approved	Harvested	Approved	Harvested
HOR 1	2005–06	–	–	2 000	150
	2006–07	220	220	150	150
	2007–08	200	150	–	–
	2008–09	150	70	90	90
	2009–10	–	–	–	–
	2010–11	–	–	100	0
	2011–12	–	–	50	0
	2012–13	–	–	–	–
	2013–14	–	–	–	–
	2014–15	–	–	–	–
	2015–16	–	–	–	–
	2016–17	100	50	80	0
	2017–18	40	40	–	–
	2018–19	–	–	–	–
	2019–20	–	–	–	–
	2020–21	–	–	–	–

1.4 Illegal catch

There is no known illegal catch of this mussel.

1.5 Other sources of mortality

There is no quantitative information on other sources of mortality, although widespread die-offs appear to be characteristic of this species. Storm scour, shell damage and subsequent predation, and exceeding carrying capacity have been suggested as possible reasons for this.

2. BIOLOGY

The horse (or fan) mussel, *Atrina zelandica*, is a widespread endemic bivalve that lives mainly on muddy-sand substrates in the lowest inter-tidal and sub-tidal shallows of mainly sheltered waters. Horse mussels are also found in deeper waters (to 50 m) off open coasts. The horse mussel is a flattened, emergent, filter-feeding mollusc, particularly conspicuous because of its size and abundance. Although more usually 260–300 mm long (110–120 mm wide) it can reach 400 mm in length and is New Zealand's largest bivalve. Horse mussels often live in groups, forming patches of up to 10 m² or more. The shell remains firmly embedded in the substrate by its pointed anterior end, the animal anchored to particles in the sediment by its byssus. The crenellated posterior edge projects a few centimetres above the substrate, keeping the water intake clear of surface deposits and providing attachment for an array of algae and invertebrates such as sponges and sea squirts.

Horse mussels are dioecious broadcast spawners. Although spawning may take place throughout much of the year it is probably mainly during summer. There is no information on the size or age at which breeding begins. A pelagic larva is free swimming for several days or weeks but nothing is known of its primary settlement locations, which may not necessarily be within the adult beds (some bivalves including soft sediment ones such as pipi settle in one area but later migrate to another where adult beds develop). Recruitment events can be sporadic and short-lived.

There is little published information on age, growth, and mortality for horse mussels. It appears that *Atrina* grows rapidly for at least the first 2–4 years: shells about 120 mm long in a northern bed increased about 40 mm per year until 166 mm, after which growth slowed dramatically (Hay C. pers. comm. in Hayward et al. 1999). Large shells are at least 5 y and possibly up to 15 y old. Widespread die-offs seem to be a feature of this species (Allan & Walshe 1984, Hayward et al. 1999). For example, in the Rangitoto Channel, densities of 200–300 per m² reduced to 1–35 per m² over 2–3 y, with storm scour, shell damage and subsequent predation, and exceeding carrying capacity being possible reasons (Hayward et al 1999).

Horse mussels have widespread effects on ecosystem structure and function (Lohrer et al. 2013). They provide shelter and refuge for invertebrates and fish (Townsend et al. 2015) and act as substrata for the settlement of epifauna such as sponges and soft corals. They also affect boundary layer dynamics and

HORSE MUSSEL (HOR)

facilitate productivity and biodiversity by depositing pseudofaeces. The horse mussel community in most northern harbours is almost entirely subtidal, in medium to fine muddy, but fairly stable, sand with moderate current velocities and no wave action. Similar communities have been observed in the Hauraki Gulf and Marlborough Sounds. Scallops, dredge oysters, and green lipped mussels are the main commercial shellfish species with beds that sometimes broadly overlap with the horse mussel distribution.

3. STOCKS AND AREAS

For management purposes stock boundaries are based on FMAs; however, there is no biological information on stock structure, recruitment patterns, or other biological characteristics which might indicate stock boundaries.

4. STOCK ASSESSMENT

4.1 Estimates of fishery parameters and abundance

There are no estimates of fishery parameters or abundance for any horse mussel fishstock.

4.2 Biomass estimates

There are no biomass estimates for any horse mussel fishstock.

4.3 Yield estimates and projections

There are no estimates of *MCY* for any horse mussel fishstock.

There are no estimates of *CAY* for any horse mussel fishstock.

5. STATUS OF THE STOCKS

There are no estimates of reference or current biomass for any horse mussel fishstock. It is not known whether horse mussel stocks are at, above, or below a level that can produce *MSY*.

6. FOR FURTHER INFORMATION

- Allan, L; Walshe, K (1984) Update on New Zealand horse mussel research. *Catch* '84 11(8): 14.
- Booth, J D (1983) Studies on twelve common bivalve larvae, with notes on bivalve spawning seasons in New Zealand. *New Zealand Journal of Marine and Freshwater Research* 17: 231–265.
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Cummings, V J; Thrush, S F; Hewitt, J E; Turner, S J (1998) The influence of the pinnid bivalve *Atrina zelandica* Gray on benthic macroinvertebrate communities in soft-sediment habitats. *Journal of Experimental Marine Biology and Ecology* 228: 227–240.
- Estcourt, I N (1967) Distributions and associations of benthic invertebrates in a sheltered water soft-bottom environment (Marlborough Sounds, New Zealand). *New Zealand Journal of Marine and Freshwater Research* 1: 352–370.
- Hayward, B W; Morley, M S; Hayward, J J; Stephenson, A B; Blom, W M; Hayward, K A; Grenfell, H R (1999) Monitoring studies of the benthic ecology of Waitemata Harbour, New Zealand. *Records of the Auckland Museum* 36: 95–117.
- Lohrer, A M; Rodil, I F; Townsend, M; Chiaroni, L D; Hewitt, J E; Thrush, S F (2013) Biogenic habitat transitions influence facilitation in a marine soft-sediment ecosystem. *Ecology* 94(1): 136 – 145.
- McKnight, D G (1969) An outline distribution of the New Zealand shelf fauna. Benthos survey, station list, and distribution of the Echinoidea. *New Zealand Oceanographic Institute Memoir* No. 47.
- Paul, L J (1966) Observations on past and present distribution of mollusc beds in Ohiwa Harbour, Bay of Plenty. *New Zealand Journal of Science* 9: 30–40.
- Townsend, N; Lohrer, A M; Rodil, I F; Chiaroni, L D (2015) The targeting of large-sized benthic macrofauna by an invasive portunid predator: evidence from a caging study. *Biological Invasions* 17(1): 231–244.
- Warwick, R M; McEvoy, A J; Thrush, S F (1997) The influence of *Atrina zelandica* Gray on meiobenthic nematode diversity and community structure. *Journal of Experimental Marine Biology and Ecology* 214: 231–247.