


INTRODUCTION - SURF CLAMS

Surf clam is a generic term used here to cover the following seven species:

Deepwater tuatua	<i>Paphies donacina</i>	(PDO)
Fine (silky) dosinia	<i>Dosinia subrosea</i>	(DSU)
Frilled venus shell	<i>Bassina yatei</i>	(BYA)
Large trough shell	<i>Mactra murchisoni</i>	(MMI)
Ringed dosinia	<i>Dosinia anus</i>	(DAN)
Triangle shell	<i>Crassula aequilatera</i>	(SAE)
Trough shell	<i>Mactra discors</i>	(MDI)

The same FMAs apply to all these species and this introduction will cover issues common to all of these species. Each species has its own chapter.



1. INTRODUCTION

All surf clams were introduced into the Quota Management System on 1 April 2004. The fishing year is from 1 April to 31 March and commercial catches are measured in greenweight. There is no minimum legal size (MLS) for surf clams. Surf clams are managed under Schedule 6 of the Fisheries Act 1996. This allows them to be returned to the sea soon after they are taken, provided they are likely to survive.

The development of hydraulic dredges, facilitated by the New Zealand Fishing Industry Board in 1985, enabled the potential for surf clam fisheries on exposed surf beaches to be investigated. Exploratory surveys between 1988 and 1992 found high densities of surf clams off beaches in Poverty Bay, the Kapiti and Manawatu coasts, Marlborough, Pegasus Bay, and Oreti Beach in Southland.

Before surf clams were introduced into the Quota Management System, commercial surf clam fisheries were managed using special permits (sections 63,64 (1) (c), and 66 of the Fisheries Act 1983). Four special permits for exploratory fishing were issued between 1986 and 1992. Permit holders had a sole right to fish a defined length of coastline and had an annual quota for each surf clam species. A moratorium was placed on the issue of new permits in 1993 to allow structured and informed development of surf clam fisheries. A special permit to fish for surf clams in Southland was granted in 2000. Commercial fishing trials began in FMA 8 in 1986, and FMAs 3 and 7 in 1987.

New Zealand operates a mandatory shellfish quality assurance programme for all bivalve shellfish grown and harvested in areas for human consumption. Shellfish caught outside this programme can only be sold for bait. This programme is based on international best practice and is managed by New Zealand Food Safety, in cooperation with the District Health Board Public Health Units and the shellfish industry¹. This involves surveying the water catchment area for pollution, sampling water and shellfish microbiologically over at least 12 months, classifying and listing areas for harvest, regular monitoring of the water and shellfish, biotoxin testing, and closure after rainfall and when biotoxins are detected. Products are traceable by source and time of harvest in case of contamination.

2. BIOLOGY

Three families of surf clams dominate the biomass in different regions of New Zealand. At the northern locations, the venerids *D. anus* and *D. subrosea* make up the major proportion of the surf clam biomass, and *D. anus* is abundant at all other North Island locations. The mactrids and mesodesmatid become increasingly abundant south of Ohope (Bay of Plenty). The mesodesmatid *P. donacina* is most abundant around central New Zealand from Nuhaka on the east coast south to the Kapiti Coast, Cloudy Bay, and as far south as Pegasus Bay. The mactrids *M. murchisoni* and *M. discors* dominate in southern New Zealand (Blueskin Bay, Te Waewae, and Oreti), where they account for more than 80% of the total biomass (Cranfield et al. 1994, Cranfield & Michael 2001).

Each species grows to a larger size in the South Island than in the North Island (Cranfield & Michael 2002). Growth parameters are available for several surf clam species from two locations, Cloudy Bay (FMA 7) and the Kapiti Coast (FMA 8). Length frequencies of sequential population samples were analysed by Cranfield et al. (1993) using MULTIFAN to estimate the von Bertalanffy growth parameters (Table 1). MULTIFAN simultaneously analyses multiple sets of length frequency samples using a maximum likelihood method to estimate the proportion of clams in each age class and the von Bertalanffy growth parameters (see Fournier et al. 1990). Less confidence should be placed in the estimates from MULTIFAN for Cloudy Bay relative to the Kapiti Coast because length sample sizes were smaller from Cloudy Bay, and there was a lack of juvenile surf clams in samples.

Incremental growth of recaptured marked clams at Cloudy Bay was analysed using GROTAG to corroborate the MULTIFAN estimates (Cranfield et al. 1993). GROTAG uses a maximum-likelihood method to estimate growth rate (Francis 1988). The estimates and annual mean growth estimates at lengths α and β are shown in Table 2. Low numbers of marked clams were recaptured from the Kapiti Coast, indicative estimates are given in Cranfield & Michael (2001).

Table 1: Von Bertalanffy growth parameter estimates from Cranfield et al. (1993) for surf clams estimated using MULTIFAN (SE in parentheses). – Indicates where estimates were not generated.

Stock	Site	L_{∞} (mm)	K
BYA 7	Cloudy Bay	–	–
BYA 8	Kapiti Coast	–	–
DAN 7	Cloudy Bay	0.10 (0.03)	77.5 (0.71)
DAN 8	Kapiti Coast	0.13 (0.02)	58.7 (0.28)
DSU 7	Cloudy Bay	–	–
DSU 8	Kapiti Coast	–	–
MDI 7	Cloudy Bay	0.41 (0.03)	68.0 (0.35)
MDI 8	Kapiti Coast	0.42 (0.02)	56.0 (0.95)
MMI 7	Cloudy Bay	0.57 (0.01)	88.0 (0.44)
MMI 8	Kapiti Coast	0.35 (0.01)	75.2 (0.30)
PDO 7	Cloudy Bay	0.33 (0.01)	94.1 (0.29)
PDO 8	Kapiti Coast	–	–
SAE 7	Cloudy Bay	1.01 (0.02)	60.3 (0.92)
SAE 8	Kapiti Coast	0.80 (0.03)	52.1(0.25)

¹ For full details of this programme, refer to the Animal Products (Regulated Control Scheme-Bivalve Molluscan Shellfish) Regulations 2006 and the Animal Products (Specifications for Bivalve Molluscan Shellfish) Notice 2006 (both referred to as the BMSRCS) at: <http://www.nzfsa.govt.nz/industry/sectors/seafood/bms/page-01.htm>

Table 2: Mean annual growth estimates (mm/year) at lengths α and β (95% confidence intervals in parentheses for mean growth values) from Cloudy Bay (Cranfield et al. 1996). L^* is the transitional length, at which point the model allows an asymptotic reduction in growth rate and values of L_∞ are included for reference.

Species	α (mm)	g_α (mm year ⁻¹)	β (mm)	g_β (mm year ⁻¹)	L^* (mm)	L_∞ (mm)	Residual error (mm)
<i>Paphies donacina</i>	50.0	10.26 (9.7 – 10.8)	80.0	1.41 (1.1 – 1.7)	80.0	84.8	1.25
<i>Crassula aequilatera</i>	30.0	22.71 (22.2 – 23.0)	50.0	6.23 (6.0 – 6.4)	55.0	57.6	2.04
<i>Mactra murchisoni</i>	40.0	17.83 (17.4 – 18.2)	70.0	4.65 (4.3 – 4.9)	80.0	80.6	1.42
<i>Mactra discors</i>	35.0	11.01 (10.5 – 11.7)	55.0	2.69 (2.4 – 2.9)	62.0	61.5	0.63
<i>Dosinia anus</i>	20.0	12.5 (12.0 – 13.2)	55.0	1.99 (1.8 – 2.2)	63.0	61.6	0.44

The maximum ages for these species were estimated from the number of age classes indicated in MULTIFAN analyses, and from shell sections. Estimates of natural mortality come from age estimates (Table 3). Higher mortality is seen where the surf clams are subject to higher wave energies, e.g., *C. aequilatera* and *M. murchisoni* are distributed within the primary wave break and hence show higher mortality (Cranfield et al. 1993). The maximum age of Kapiti coast shells suggests higher mortality than Cloudy Bay, perhaps because the Kapiti Coast is more exposed and surf clams there have a higher chance of being eroded out of the seabed by storms (Cranfield et al. 1993). Surf clam populations are subject to catastrophic mortality from being dislodged from within the seabed and being washed ashore during storms, high nearshore seawater temperatures and low oxygen levels during calm summer periods, blooms of toxic algae, and long periods of reduced salinity from high freshwater outflow (Eggleston & Hickman 1972, Cranfield & Michael 2001).

Table 3: Estimates of the instantaneous natural mortality rate, M . A = minimum number of year classes indicated by MULTIFAN; B = maximum age indicated by shell sections; M1 = mortality range estimated from using two equations: $\ln M = 1.23 - 0.832 \ln(t_{max})$ and $\ln M = 1.44 - 0.982 \ln(t_{max})$, (Hoenig 1983); M2 mortality estimated from $M = \ln 100 / (t_{max})$; t_{max} is the estimate of maximum age.

	A	B	M1	M2
Cloudy Bay				
<i>Mactra murchisoni</i>	8	11	0.40–0.46	0.42
<i>Mactra discors</i>	7	14	0.32–0.38	0.33
<i>Crassula aequilatera</i>	5	7	0.63–0.68	0.66
<i>Paphies donacina</i>	10	17	0.26–0.32	0.27
<i>Dosinia anus</i>	16	22	0.20–0.26	0.21
Kapiti Coast				
<i>Mactra murchisoni</i>	8	11*	0.40–0.46	0.42
<i>Mactra discors</i>	8	16*	0.28–0.34	0.29
<i>Crassula aequilatera</i>	3	5*	0.87–0.89	0.92
<i>Paphies donacina</i> [†]				
<i>Dosinia anus</i>	19	26*	0.17–0.23	0.18

*Shell sections not yet examined. Ages are inferred from Cloudy Bay data.

[†]Growth data could not be analysed.

3. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was first introduced to the May 2011 Plenary after review by the Aquatic Environment Working Group and is updated as relevant research is undertaken and published. It was last updated in 2021. This summary is from the perspective of the surf clam fisheries; a more detailed summary from an issue-by-issue perspective is available in the Aquatic Environment and Biodiversity Annual Review 2021 (Fisheries New Zealand 2021), online at <https://www.mpi.govt.nz/dmsdocument/51472-Aquatic-Environment-and-Biodiversity-Annual-Review-AEBAR-2021-A-summary-of-environmental-interactions-between-the-seafood-sector-and-the-aquatic-environment>.

3.1 Ecosystem role

Worldwide, the ecosystem roles of subtidal communities on exposed surf beaches are difficult to study. The physical environments of surf beaches are defined by their sand, wave, and tidal regimes. Surf beaches range from narrow and steep (reflective) beaches with no surf zone and coarse sediments (e.g., Cloudy Bay, FMA 7) to wide and flat (dissipative) beaches with extensive surf zones, finer sand and larger waves and tides (e.g., Pegasus Bay, (FMA 3), after Defo et al. 2009). The surf zone systems of New Zealand dissipative beaches are similar to those described in California (Morin et al. 1985) and South Africa (McLachlan et al. 1984). There is distinct zonation of subtidal, beach communities driven by an environmental gradient: the nearshore swash zone characterized by almost continual sweeping and shifting sands churned up by strong, fluctuating wave surge at one extreme, to a relatively benign environment with reduced surge, and finer sediments with almost no sand movement at the other extreme (Morin et al. 1985). As common on most exposed surf beaches, suspension feeding bivalves dominate the nearshore biomass. Beyond the primary wave break where the bottom becomes more stable and finer, there is a rapid rise in species diversity, but in New Zealand, generally a decreased in overall biomass.

Only three published papers examine aspects of the role of surf clams in the ecosystem in New Zealand. Juvenile surf clams, mainly *Paphies* spp., are important prey for the paddle crab *Ovalipes catharus* in central New Zealand (Wear & Haddon 1987). Predation of *Dosinia* spp. by rock lobsters has been documented from the reef/soft sediment interface zones (Langlois et al. 2005, Langlois et al. 2006), notably surf clams are usually harvested from exposed beaches, not reef/soft sediment interface zones.

Surf clams are filter-feeders; recent research suggests that most of their food is obtained from microalgae from the top 2 cm of the sediment and the bottom 2–3 cm of the water column (Sasaki et al. 2004). The effects of predation are difficult to study on exposed sandy beaches and it is believed internationally that there are no keystone species in this environment and predation is not important in structuring the community (McLachlan & Brown 2006).

3.2 Fishery captures (fish and invertebrates)

The only bycatch caught in large quantities associated with surf clam dredging in New Zealand is *Fellaster zelandiae* — the sand dollar or sea biscuit (Haddon et al. 1996, Triantafillos 2008a, Triantafillos 2008b, White et al. 2012). Other species caught in association with surf clams include paddle crabs (*Ovalipes catharus*), a number of bivalves including the lance shell (*Resania lanceolata*), otter clams (*Zenatia acinaces*), battle axe (*Myadora striata*), olive tellinid (*Hiatula nitidia*), the wedge shell (*Peronaea gairmadi*), and the gastropods the olive shell (*Baryspira australis*) and ostrich foot shell (*Struthiolaria papulosa*). Fish are rarely caught but include juvenile common soles (*Peltorhamphus novaezeelandiae*) and stargazers (*Kathetostoma* spp.) (NIWA, unpublished data).

3.3 Fishery captures (seabirds and mammals)

Not relevant to surf clam fisheries.

3.4 Benthic impacts

Surf clams mainly inhabit the surf zone, a high-energy environment characterised by high sand mobility (Michael et al. 1990). Divers observed that the survey dredge (used for surf clam surveys) formed a well-defined track in the substrate, but within 24 hours (probably much less time) the track could not be distinguished, indicating that physical recovery of the substrate was rapid (Michael et al. 1990). A different dredge is used for commercial fishing, and the impacts of this dredge have not been tested. Shallow water environments such as the surf zone or those subjected to frequent natural disturbance tend to recover faster from the effects of mobile fishing gears compared with those in deeper water (Kaiser et al. 1996, Collie et al. 2000, Hiddink et al. 2006, Kaiser et al. 2006).

Incidental effects of hydraulic dredging on the target species within New Zealand fisheries have not been studied. Any effects will be species, size, dredge, and location specific. For the Italian surf clam (*Chamelea gallina*), laboratory experiments found that 50% of the surf clams had reburied after 4 hours (95% CI 3.6–4.4) and 90% after 8 hours (95% CI 7.7–8.2) (Bargione et al. 2021). There are no data on

reburial times of New Zealand surf clams encountering hydraulic dredges. Given the time it takes for *Chamelea gallina* to rebury, and New Zealand surf conditions, surf clams may be washed ashore. In the same research Bargione et al (2021) found survival of disturbed clams in both laboratory experiments and caged sea trials was high. A laboratory study by Cranfield and Michael (1995) found that surf clams (PDO, SAE, and MMI) float on liquified sand, and based on laboratory observations suggest that some surf clams that encounter hydraulic dredges may suffer some damage or stress. The survival of New Zealand surf clams encountering hydraulic dredges and the return of surf clams to sea is currently unknown.

Surf clam species show zonation by sediment type and mobility which is generally, although not always, correlated with depth and wave exposure. Species with good burrowing ability are generally found in shallow, mobile sediment zones (for example, *Paphies donacina*), and those species less able to burrow (for example, *Dosinia subrosea* and *Bassina yatei*) are generally found in softer, more stable sediments. The present high-value species (*Crassula aequilatera*, *Macra murchisoni*, *Paphies donacina* and *Macra discors*) generally occur in transition zones from highly mobile sediments to mostly stable sediment. Mobile fishing gear effects will be primarily determined by the characteristics of the beach and target species. Little fishing presently takes place in the most vulnerable areas characterised by stable, soft fine sediment communities because of the lower densities of surf clams and the presently lower value of species there.

An Italian study showed that widespread intensive hydraulic dredging can adversely modify habitat at depths of 4–6 m within the surf zone environment, although recovery in this study occurred within six months (Morello et al. 2006). Scottish (Tuck et al. 2000; Hauton et al. 2003a, 2003b) and American (Mercaldo-Allen et al. 2017) studies detected benthic community and physical impacts, but survival of discarded bycatch was generally high. The applicability of these studies' finding to New Zealand is unknown, given the relatively sheltered nature of some of the habitats examined compared to surf beaches. In New Zealand, the effects of hydraulic dredging are likely to vary with site specific environmental conditions and depending on whether beaches have reflective or dissipative beach profiles.

3.5 Other considerations

None.

3.6 Key information gaps

The distribution of hydraulic dredging in New Zealand is localised at fishery scale, and generally not widespread geographically. The impacts of hydraulic dredging at the intensities fished are unknown. Key information gaps include population connectivity and stock structure, and spatial and temporal variability in key population parameters, e.g., recruitment patterns varied on Wellington west coast beaches with juveniles of *Macra discors* and *M. murchisoni* dominating in alternate years (Conroy et al. 1996).

A key information gap is the survival of surf calms encountered by the hydraulic dredge and that pass under or through the dredge on the seabed, and of those landed onboard and returned to sea.

4. FOR FURTHER INFORMATION

- Bargione, G., Petetta, A., Vasapollo, C. et al. Reburial potential and survivability of the striped venus clam (*Chamelea gallina*) in hydraulic dredge fisheries. *Sci Rep* 11, 9109 (2021). <https://doi.org/10.1038/s41598-021-88542-8>
- Brierley, P.J.; Cranfield, J.; Knedler, K.; Bauckham, A.; Town, J.; Stevens, P.; Sullivan, M.; Blank, F.; Martin, N. (1990). Management and development of the New Zealand sub-tidal clam fishery. Report of the Surf Clam Working Group. (Unpunished report held by MAF Fisheries. New Zealand). 57 p.
- Collie, J.; Escanero, G.; Valentine, P (2000) Photographic evaluation of the impacts of bottom fishing on benthic epifauna. *ICES Journal of Marine Science* 57, 987–1001.
- Conroy, A.M.; Smith, P.J.; Michael, K.P.; Stotter, D.R. (1993). Identification and recruitment patterns of juvenile surf clams, *Macra discors* and *M. murchisoni* from central New Zealand. *New Zealand Journal of Marine and Freshwater Research* 27(3): 279-285.
- Constantino, R; Caspar, M B; Tata-Regala, J; Carvalho, S; Curdia, J; Drago, T; Tabora, R; Monteiro, C C (2009) Clam dredging effects and subsequent recovery of benthic communities at different depth ranges. *Marine Environmental Research* 67(2): 89–99.

- Cranfield, H.J.; Michael, K.P. (1950). A laboratory study of the behaviour of surf clams (*Paphies donacina*, *Spisula aequilatera*, *Macra murchisoni*). (Unpublished report to Kai Moana (Pacific) Ltd., held by NIWA).
- Cranfield, H.J.; Michael, K.P. (2001). Growth rates of five species of surf clams on a southern North Island beach, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 35: 909-924.
- Cranfield, H.; Michael, K (2001) The surf clam fishery in New Zealand: description of the fishery, its management, and the biology of surf clams. *New Zealand Fisheries Assessment Report 2001/62*. 24 p.
- Cranfield, H; Michael, K (2002) Potential area boundaries and indicative TACs for the seven species of surf clam. (Unpublished report to the Ministry of Fisheries, held by Fisheries New Zealand).
- Cranfield, H; Michael, K; Francis, R (1996) Growth Rates of Five Species of Subtidal Clam on a Beach in the South Island, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 47: 773-784.
- Cranfield, H J; Michael, K P; Stotter, D R (1993) Estimates of growth, mortality, and yield per recruit for New Zealand surf clams. New Zealand Fisheries Assessment Research Document 1993/20. 26 p. (Unpublished document held by NIWA library, Wellington).
- Cranfield, H J; Michael, K P; Stotter, D R; Doonan, I J (1994) Distribution, biomass and yield estimates of surf clams off New Zealand beaches. New Zealand Fisheries Assessment Research Document 94/11. 17p. (Unpublished document held in NIWA library, Wellington).
- Defeo, O.; McLachlan, A.; Schoeman, D.S.; Schlacher, T.A.; Dugan, J.; Jones, A.; Lastra, M.; Scapini, F. (2009). Threats to sandy beach ecosystems: a review. *Estuarine, Coastal and Shelf Science* 81(1): 1-12.
- Eggleston, D.; Hickman, R.W. (1972). Mass Strandings of Molluscs at Te Waewae Bay, Southland, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 6(3): 379-382.
- Fisheries New Zealand (2021) Aquatic Environment and Biodiversity Annual Review 2021. Compiled by the Aquatic Environment Team, Fisheries Science and Information, Fisheries New Zealand, Wellington, New Zealand. 779 p.
- Fournier, D A; Sibert, J R; Majkowski, J; Hampton, J (1990) MULTIFAN a likelihood-based method for estimating growth parameters and age composition from multiple length frequency data sets illustrated using data for southern bluefin tuna (*Thunnus maccoyii*). *Canadian Journal of Fisheries and Aquatic Sciences* 47: 301-317.
- Francis, R I C (1988) Maximum likelihood estimation of growth and growth variability from tagging data. *New Zealand Journal of Marine and Freshwater Research* 22: 42-51.
- Haddon, M; Willis, T J; Wear, R G; Anderlini, V C (1996) Biomass and distribution of five species of surf clam off an exposed west coast North Island beach, New Zealand. *Journal of Shellfish Research* 15(2): 331-339.
- Hauton, C.; Atkinson, R.J.A.; Moore, P.G. (2003a) The impact of hydraulic blade dredging on a benthic megafaunal community in the Clyde Sea area, Scotland. *Journal of Sea Research* 50, 45-56.
- Hauton, C.; Hall-Spencer, J.M.; Moore, P.G. (2003b) An experimental study of the ecological impacts of hydraulic bivalve dredging on maerl. *ICES Journal of Marine Science* 60(2), 381-392.
- Hiddink, J; Jennings, S; Kaiser, M; Queiros, A; Duplisea, D; Piet, G (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences* 63, 721-736.
- Hoenig, J M (1983) Empirical use of longevity data to estimate mortality rates. *Fishery Bulletin* 82: 898-903.
- Kaiser, M; Clarke, K; Hinz, H; Austen, M; Somerfield, P; Karakassis, I (2006) Global analysis of response and recovery of benthic biota to fishing. *Marine Ecology-Progress Series* 311: 1-14.
- Kaiser, M; Hill, A; Ramsay, K; Spencer, B; Brand, A; Veale, L; Prudden, K; Rees, E; Munday, B; Ball, B; Hawkins, S (1996) Benthic disturbance by fishing gear in the Irish Sea: a comparison of beam trawling and scallop dredging. *Aquatic Conservation: Marine and Freshwater Ecosystems* 6: 269-285.
- Langlois, T; Anderson, M J; Babcock, R; Kato, S (2005) Marine reserves demonstrate trophic interactions across habitats. *Oecologia* 147(1): 134-140.
- Langlois, T J; Anderson, M J; Brock, M; Murman, G (2006) Importance of rock lobster size-structure for trophic interactions: choice of soft-sediment bivalve prey. *Marine Biology* 149(3): 447-454.
- McLachlan, A; Brown, A (2006) *The Ecology of Sandy Shores*. Academic Press.
- McLachlan, A.; Cockcroft, A.; Malan, D. (1984). Benthic faunal response to a high energy gradient. *Marine Ecology Progress Series*. 16(1): 51-63.
- Mercaldo-Allen, R.; Meseck, S.; Goldberg, R.; Clark, P.; Kuropat, C.; Rose, J.M. (2017) Effects of clam dredging on benthic ecology of two cultivated northern quahog beds with different harvest histories and sediment grain sizes. *Aquaculture International* 25: 1971-1985.
- Michael, K; Olsen, G; Hvid, B; Cranfield, H (1990) Design and performance of two hydraulic subtidal clam dredges in New Zealand. *New Zealand Fisheries Technical Report* No. 21. 16 p.
- Minami, T; Sasaki, K; Ito, K (2009) Recent research topics in biological production processes of marine bio-resources in the coastal waters and estuaries. *Tohoku Journal of Agricultural Research* 59(3-4): 67-70.
- Morello, E B; Frogliia, C; Atkinson, R G A; Moore, P G (2005) Impacts of hydraulic dredging on a macrobenthic community of the Adriatic Sea, Italy. *Canadian Journal of Fisheries and Aquatic Sciences* 62(9): 2076-2087.
- Morello, E B; Frogliia, C; Atkinson, R G A; Moore, P G (2006) Medium-term impacts of hydraulic clam dredgers on a macrobenthic community of the Adriatic Sea (Italy). *Marine Biology* 149(2): 401-413.
- Morin, J.; Kastendiek, J.; Harrington, A.; Davis, N. (1985). Organization and patterns of interactions in a subtidal sand community on an exposed coast. *Marine Ecology Progress Series* 27: 163-185.
- Reise, K (1984) *Tidal flat ecology*. Berlin, Springer-Verlag.
- Sasaki, K; Sanematsu, A; Kato Y., Ito K. (2004) Dependence of the Surf Clam *Pseudocardium sachalinense* (Bivalvia: Mactridae) On the Near-Bottom Layer for Food Supply. *Journal of Molluscan Studies* 70(3): 207-212.
- Triantafillos, L (2008a) Survey of subtidal surf clams in Pegasus Bay, November-December 2007. Prepared by NIWA for Seafood Innovations Limited and SurfCo. Limited. 43 p. (Unpublished document held by Fisheries New Zealand.).
- Triantafillos, L (2008b) Survey of subtidal surf clams in Quota Management Area 2, June-August 2008. Prepared by NIWA for Seafood Innovations Limited and SurfCo. Limited. 40 p. (Unpublished document held by Fisheries New Zealand).
- Tuck, I.D.; Bailey, N.; Harding, M.; Sangster, G.; Howell, T.; Graham, N.; Breen, M. (2000) The impact of water jet dredging for razor clams, *Ensis* spp., in a shallow sandy subtidal environment. *Journal of Sea Research* 43, 65-81.
- White, W. L., Millar, R. B., Breen, B. and Farrington, G. (2012) Survey of subtidal surf clams from the Manawatu Coast (FMA8), October-November 2012. Report for the Shellfish Working Group Meeting 19th November 2012, 35 p.+ Addendum.
- Wear, R.; Haddon, M. (1987). Natural diet of the crab *Ovalipes catharus* (Crustacea, Portunidae) around central and northern New Zealand. *Marine Ecology Progress Series*: 39-49.