


JACK MACKERELS (JMA)

(*Trachurus declivis*, *Trachurus novaezelandiae*, *Trachurus murphyi*)
Hauture



1. FISHERY SUMMARY

The jack mackerel fisheries catch three species: two endemic species, *Trachurus declivis* and *T. novaezelandiae*, and *T. murphyi* which appeared in New Zealand in the 1980s.

Jack mackerels have been fully included in the QMS since 1 October 1996, with four QMAs. From 1 October 1987 jack mackerels were considered part of the QMS, although ITQs were issued only in JMA 7. In JMA 1 and JMA 3 quota for the fishery was fully allocated as IQs by regulation, with the exception of the 20% allocated to customary non-commercial catch. Before the 1995 jack mackerel regulations were issued, catch in JMA 1 taken in the Muriwhenua area north of 36° S to the limit of the Territorial Sea was not covered by the JMA 1 regulations. Allowances for customary non-commercial fishers, recreational fishers, and an allowance for other sources of mortality have only been set in JMA 3 (Table 1).

Table 1: TACs, TACCs, and allowances (t) for jack mackerels by fishstock.

Fishstock	TAC	TACC	Customary allowance	Recreational allowance	Other mortality
JMA 1	–	10 000	–	–	–
JMA 3	9 000	8 780	20	20	180
JMA 7	–	32 537	–	–	–
JMA 10	–	10	–	–	–

1.1 Commercial fisheries

In JMA 1, the jack mackerel catch is largely taken by the target purse seine fishery operating in the Bay of Plenty in Statistical Area 009 during March–November, with minor catches taken as a bycatch of kahawai and blue mackerel purse seine fisheries, and as a bycatch from trawl fisheries. In most years, relatively small catches were taken from off the east Northland coast (Statistical Areas 002 and 003), although this area accounted for a substantial proportion of the total catch in 1993–94 and 1994–95.

Since 1991–92, jack mackerel targeted landings in JMA 1 have represented more than 80% of total catch. The highest rates of bycatch are from kahawai and blue mackerel targeted operations which each account for about 7% of the total jack mackerel catch. The majority of JMA 1 catch over these years

JACK MACKERELS (JMA)

has been taken from Statistical Areas 008 and 009 (Bay of Plenty) between June and November; considerably less has been taken in Statistical Areas 002 and 003, although high catches were recorded from these areas in 1993–94 and 1994–95.

In JMA 3 little targeting occurred before 1992–93. During the 1990s targeting increased and accounted for the majority of catch (about 50% between 1991–92 and 1996–97), but, after a peak of more than 80% in 1997–98 and 1998–99, the target catch has decreased again to about 50–60% in recent years. The balance of the catch in this area comes from trawl bycatch (squid 15–30%, barracouta 15–20%) on the Chatham Rise and in the Southland/Sub-Antarctic region. A purse seine fishery has operated between the Clarence River mouth and the Kaikōura Peninsula, which peaked at 4400 t in 1992–93 and averaged more than 3000 t between 1989–90 and 1993–94. Purse seine catches have shown a steady decline since, dropping from 1000 t in 1994–95, to 100 t in 2001–02 and 2002–03; no catch was recorded for 2003–04, and purse seine catch has subsequently been rare.

Increased availability of jack mackerels caused by the influx of *T. murphyi* resulted in increased TACCs in JMA 1 and JMA 3, to 8000 t and 9000 t, respectively, for the 1993–94 fishing year, and a further increase to 10 000 t and 18 000 t, respectively, for the 1994–95 year. The latter increases were made under the proviso that they be accounted for by increased catches of *T. murphyi* only; combined landings of *T. declivis* and *T. novaezelandiae* in JMA 1 and JMA 3 must not exceed the original quotas of 5970 t and 2700 t, respectively. Industry agreed to these limits and voluntarily introduced monitoring programmes to provide the information necessary for them to be met.

From the 2016–17 fishing year, the TACC for JMA 3 was reduced to 8780 t, approximating the 1993–94 TACC level, on the basis that recent catches had been considerably lower than the TACC and that catches of *T. murphyi* were minimal, indicating low abundance of the species in New Zealand waters in recent years.

The three species occur in each of the Fishstocks but are not individually identified in catch reporting. Historical estimated and recent reported jack mackerel landings and TACCs are shown in Tables 1 and 2, and Figure 1 shows the historical landings and TACC values for the main JMA stocks. Total annual landings have ranged between 21 059 t and 50 388 t since 1986–87 (Table 3).

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	JMA 1	JMA 3	JMA 7	Year	JMA 1	JMA 3	JMA 7
1931–32	0	0	0	1957	0	0	6
1932–33	0	0	0	1958	0	0	9
1933–34	0	0	0	1959	2	0	0
1934–35	0	0	0	1960	2	0	5
1935–36	0	0	0	1961	1	0	5
1936–37	0	0	0	1962	5	0	5
1937–38	0	0	0	1963	7	2	13
1938–39	0	0	0	1964	5	4	10
1939–40	1	0	0	1965	14	0	8
1940–41	1	1	2	1966	47	0	54
1941–42	0	0	2	1967	213	0	250
1942–43	3	0	2	1968	172	505	4 558
1943–44	0	0	0	1969	128	388	7 065
1944	9	0	0	1970	75	1 029	7 274
1945	7	0	0	1971	473	776	12 684
1946	3	0	6	1972	350	5 450	15 581
1947	14	0	4	1973	395	1 238	14 648
1948	3	0	6	1974	1 236	2 016	16 943
1949	5	0	22	1975	204	3 615	10 043
1950	7	6	3	1976	838	5 690	14 228
1951	4	4	1	1977	1 317	5 228	13 729
1952	1	4	7	1978	1 250	1 547	4 657
1953	0	3	9	1979	2 158	516	4 475
1954	3	0	1	1980	2 504	104	3 533
1955	3	0	12	1981	2 815	110	8 665
1956	1	0	2	1982	1 607	119	8 364

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data include both foreign and domestic landings.

Table 3: Reported landings (t) of jack mackerel by Fishstock from 1983–84 to present and actual TACCs (t) for 1986–87 to present. QMS data from 1986 to present.

	JMA 1		JMA 3		JMA 7		JMA 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings§	TACC
1983–84*	3 682	–	715	–	12 464	–	0	–	16 861	–
1984–85*	1 857	–	1 223	–	16 013	–	0	–	19 093	–
1985–86*	1 173	–	2 228	–	10 002	–	0	–	13 403	–
1986–87	4 056	5 970	1 638	2 700	19 815	20 000	0	10	25 509	28 680
1987–88	3 108	5 970	1 883	2 700	17 879	22 697	0	10	22 870	31 377
1988–89	2 986	5 970	1 919	2 700	17 403	26 008	0	10	22 308	34 688
1989–90	4 226	5 970	4 013	2 700	21 776	32 027	0	10	30 015	40 707
1990–91	6 472	5 970	6 403	2 700	17 786	32 069	0	10	30 661	40 749
1991–92	7 017	5 970	5 779	2 700	25 880	32 069	0	10	38 676	40 749
1992–93	7 529	5 970	15 399	2 700	24 659	32 537	0	10	47 587	41 216
1993–94‡	14 256	8 000	9 115	9 000	22 377	32 537	0	10	45 748	49 546
1994–95‡	7 832	10 000	11 519	18 000	18 912	32 537	0	10	38 263	60 547
1995–96	6 874	10 000	19 803	18 000	12 270	32 537	0	10	38 947	60 547
1996–97	6 912	10 000	15 687	18 000	12 056	32 537	0	10	34 655	60 547
1997–98	7 695	10 000	15 452	18 000	14 293	32 537	0	10	37 440	60 547
1998–99	5 641	10 000	15 111	18 000	13 629	32 537	0	10	34 381	60 547
1999–00	2 864	10 000	10 306	18 000	7 889	32 537	0	10	21 059	60 547
2000–01	8 360	10 000	2 744	18 000	15 703	32 537	0	10	26 807	60 547
2001–02	5 247	10 000	5 000	18 000	22 338	32 537	0	10	32 585	60 547
2002–03	6 172	10 000	2 225	18 000	26 084	32 537	0	10	34 481	60 547
2003–04	7 396	10 000	705	18 000	28 888	32 537	0	10	36 989	60 547
2004–05	9 418	10 000	716	18 000	36 507	32 537	0	10	46 641	60 547
2005–06	9 924	10 000	5 000	18 000	27 782	32 537	0	10	42 706	60 547
2006–07	5 293	10 000	1 857	18 000	32 039	32 537	0	10	39 189	60 547
2007–08	11 167	10 000	2 629	18 000	34 059	32 537	0	10	47 855	60 547
2008–09	9 791	10 000	1 964	18 000	28 828	32 537	0	10	40 583	60 547
2009–10	9 086	10 000	2 706	18 000	31 152	32 537	0	10	42 944	60 547
2010–11	8 262	10 000	3 592	18 000	28 177	32 537	0	10	40 031	60 547
2011–12	8 911	10 000	3 085	18 000	28 266	32 537	0	10	40 261	60 547
2012–13	8 054	10 000	3 830	18 000	31 776	32 537	0	10	43 659	60 547
2013–14	10 520	10 000	4 693	18 000	35 175	32 537	0	10	50 388	60 547
2014–15	10 177	10 000	4 115	18 000	33 970	32 537	0	10	48 262	60 547
2015–16	6 989	10 000	2 756	18 000	30 875	32 537	0	10	40 621	60 547
2016–17	8 890	10 000	4 665	8 780	33 802	32 537	0	10	47 357	51 327
2017–18	5 553	10 000	5 559	8 780	34 190	32 537	0	10	45 302	51 327
2018–19	4 332	10 000	4 651	8 780	31 752	32 537	0	10	40 735	51 327
2019–20	6 478	10 000	5 355	8 780	31 451	32 537	0	10	43 284	51 327
2020–21	6 777	10 000	5 601	8 780	31 810	32 537	0	10	44 188	51 327
2021–22	3 455	10 000	4 858	8 780	27 782	32 537	0	10	36 095	51 327

* FSU data.

§ Includes landings from unknown areas before 1986–87.

‡ JMA 1 & 3 landings are totals from CLR and CELR data.

Landings in JMA 1 before 1989–90 were generally well below the quota of 5970 t (Table 3), with the maximum in 1986–87 only slightly above 4000 t. Landings increased to 7529 t in 1992–93, followed by a substantial increase to the highest recorded value of 14 256 t in 1993–94, which was more than twice the original quota and exceeded the quota of 8000 t set for that year. In 1994–95 reported landings (7832 t) were half those of 1993–94. Landings from 1994–95 to 1997–98 were around 7000 t. Over the period 1997–98 to 2004–05, annual catches from JMA 1 increased to near the level of the TACC (10 000 t) and, until 2014–15, annual catches fluctuated about 8000–10 000 t, with the exception of a considerably lower catch in 2006–07 and a peak catch of 11 200 t in 2007–08. JMA 1 landings since 2015–16 have been consistently less than the TACC of 10 000 t. The 2018–19 JMA 1 landings were the lowest since 1999–00 (at 4332 t), then increased to 6777 t in 2020–21, but decreased to 3455 t in 2021–22.

Estimates of the species composition of the JMA 1 purse seine catches are available from 1989–90 to 2019–20 (Figure 2, Table 4). During 1989–90 and 1990–91, annual catches were dominated by *T. novaezelandiae*, but included a small component of *T. declivis*. The proportion of *T. murphyi* in the catch increased considerably over the following years, accounting for 65% of the total catch in 1993–94 and continued to account for a considerable proportion of the JMA 1 catch during 1994–95 to 1998–99. Since 1999–00, annual catches of *T. murphyi* have been small. From 1999–00 to 2016–17, annual catches from JMA 1 were generally dominated by *T. novaezelandiae*. The annual catch of this species increased from about 2000 t to 5000 t during the 1990s to an average of 8150 t in 2007–08 to 2016–17. Correspondingly, cumulative catches of *T. declivis* and *T. murphyi* were low during this period (7% and 2%, respectively). *Trachurus novaezelandiae* annual catches dominated the JMA 1 purse seine fishery

JACK MACKERELS (JMA)

from 2014–15 to 2016–17, ranging from 6488 t to 8858 t, but dropped to 2432 t and 52% of the catch in 2017–18. Catches of *T. declivis* increased in 2017–18 and ranged from 1521 t to 2313 t from 2017–18 to 2019–20.




Figure 1: Reported commercial landings and TACC for the three main JMA stocks. From top: JMA 1 (Auckland East, Central East), JMA 3 (South East coast, South East Chatham Rise, Sub-Antarctic, Southland), and JMA 7 (Challenger, Central Egmont, Auckland West).




Figure 2: The time series of annual species catch estimates from the JMA 1 purse seine fishery (JMN, *T. novaezelandiae*; JMD, *T. declivis*; JMM, *T. murphyi*).

Table 4: Total JMA 1 purse seine catches and the time series of annual estimates of the species composition of the catch (JMN, *T. novaezelandiae*; JMD, *T. declivis*; JMM, *T. murphyi*) (compiled from various sources, see appendix 5 of Langley et al 2016 and Middleton in prep).

Fishing year	Catch (t)	Species proportion		
		JMD	JMM	JMN
1989–90	1 433	0.15	0.04	0.81
1990–91	7 147	0.15	0.10	0.76
1991–92	6 921	0.11	0.32	0.58
1992–93	8 629	0.11	0.33	0.56
1993–94	13 710	0.17	0.65	0.18
1994–95	8 530	0.13	0.45	0.42
1995–96	5 643	0.03	0.13	0.84
1996–97	6 256	0.05	0.30	0.65
1997–98	7 009	0.05	0.42	0.53
1998–99	5 077	0.14	0.30	0.56
1999–00	2 416	0.01	0.01	0.98
2000–01	7 896	0.02	0.01	0.97
2001–02	5 146	0.17	0.01	0.82
2002–03	5 518	0.30	0.02	0.68
2003–04	6 838	0.46	0.11	0.43
2004–05	8 919	0.11	0.07	0.82
2005–06	9 568	0.11	0.00	0.89
2006–07	4 803	0.44	0.26	0.31
2007–08	11 270	0.23	0.01	0.76
2008–09	9 579	0.06	0.07	0.87
2009–10	8 714	0.00	0.00	1.00
2010–11	7 936	0.00	0.00	1.00
2011–12	8 765	0.13	0.00	0.86
2012–13	7 841	0.06	0.01	0.93
2013–14	10 543	0.07	0.01	0.92
2014–15	9 968	0.05	0.01	0.94
2015–16	6 721	0.01	0.00	0.99
2016–17	8 439	0.00	0.00	1.00
2017–18	5 140	0.46	0.03	0.52
2018–19	4 111	0.37	0.01	0.62
2019–20	6 208	0.34	0.00	0.66

Total landings in JMA 3 over the period 1984–85 to 1988–89 were relatively constant, at a level below the quota of 2700 t. Landings increased over subsequent years to peak in 1992–93 at almost three times that of the preceding year and more than five times the quota. Under the first of two consecutive annual

JACK MACKERELS (JMA)

increases to the JMA 3 TACC in 1993–94, landings were slightly above the limit set, but dropped well below the higher TACC level in 1994–95. The lower 1994–95 catch relative to that in 1992–93 has been attributed to the delayed implementation of the quota, less targeting of jack mackerel, and low bycatch in the squid trawl fishery. Landings in JMA 3 increased markedly in 1995–96 (19 803 t) to a value exceeding the quota, with catches remaining stable around 15 500 t over three subsequent years. More recently, landings have decreased to levels well below the TACC, fluctuating between 700 t and about 5000 t since 2000–01. Declines in landings are attributed to declining abundance of *T. murphyi*, which historically comprised the bulk of JMA 3 landings. JMA 3 landings have been stable in recent years (average 5115 t for 2016–17 to 2021–22).

The largest jack mackerel fishery is in JMA 7. In the early 1990s, JMA 7 landings were mainly taken by bottom trawlers but, following a fleet transition in the mid-1990s to early 2000s, are now mainly taken by midwater trawlers. Landings fluctuated between 17 403 t and 25 880 t from 1986–87 to 1994–95. From 1995–96 to 1998–99, landings were in the range of 12 056–14 293 t. Subsequently, landings increased steadily from 15 703 t in 2000–01, peaking at 36 507 t in 2004–05 (exceeding the TACC by 3971 t). From 2005–06 to 2021–22, catches have fluctuated around the TACC.

A number of factors have been identified that can influence landing volumes in the jack mackerel fisheries. In the purse seine fishery during the 1990s, jack mackerel was often mixed with kahawai. Fishing companies tend to avoid these mixed schools to conserve kahawai quota, particularly at the beginning of the fishing year. When mixing of the two species is prevalent, a low kahawai TACC can result in the targeting of jack mackerel being inhibited. Both skipjack tuna and blue mackerel have been fished in preference to jack mackerel in the purse seine fishery, with the jack mackerel season being influenced by the availability of these species. However, global increases in the market price for jack mackerel have increased its importance in the purse seine fishery to a level similar to that for blue mackerel, and, as a result, the seasonal catch for jack mackerel has broadened considerably in recent years. This has provided fishers with a cost-effective alternative to traditional purse seine targets, particularly skipjack tuna, which incurs higher costs related to onboard storage and handling.

In recent years, there has been a change in the operation of the JMA 1 purse-seine fleet. In response to market requirements, fish are no longer stored in brine on board the vessel. This has resulted in shorter trip durations and consequently a concentration of fishing effort in the Bay of Plenty (where *T. novaezelandiae* dominate) near the processing facilities in Tauranga. Market requirements for fish size also affect the jack mackerel species targeted, and consequently the areas fished.

In the trawl fisheries (JMA 3 and 7) the majority of Japanese vessels (and hence the destination market) exited the fishery in the mid to late 1990s. The influx of *T. murphyi* also altered the value mix because the New Zealand species had a higher market value (at least in Japan). Since 2000, the target trawl fishery has been dominated by BATM vessels (specifically Pulkovskiy Meridian/BATM-1288 vessels, a class of 104 m factory trawlers constructed in eastern Europe and designed for distant-water fisheries) which have high processing capacity focussed on whole (GRE) or Headed and Gutted (H&G) product. These vessels operate across a range of bulk fisheries, often seasonal (e.g., SQU SBW, HOK), which means the apparent seasonality of JMA effort is primarily driven by the need to operate in the biologically seasonal (e.g., spawn) fisheries of various species as required, with JMA targeted in the temporal gaps between these events through the year.

Vessels over 46 m (i.e., all vessels in the JMA 7 midwater fleet) are excluded from nearshore areas throughout the JMA 7 QMA. These areas may represent a significant part of the habitat of jack mackerels. Vessels in this fishery have also worked to minimise bycatch of kingfish, snapper, and also common dolphins. Nevertheless, these restrictions do not appear to have constrained the fishery from taking the TACC in most years.

1.2 Recreational fisheries

Jack mackerels do not rate highly as a recreational target species although they are popular as bait.

Recreational catch in the northern region (JMA 1) was estimated at 333 000 fish (CV 0.13) by a diary survey in 1993–94 (Bradford 1996), 79 000 fish (CV 0.16) in a national recreational survey in 1996 (Bradford 1998), 349 000 fish (CV 39%) in the 2000 survey (Boyd & Reilly 2004) and 295 000 fish 658

(CV 0.2%) in the 2001 survey (Boyd et al 2004). The surveys suggest a harvest of 80–110 t per year for JMA 1, insignificant in the context of the commercial catch. Estimates from other areas are very low (between 500 and 47 000 fish) and are insignificant in the context of the commercial catch

The harvest estimates provided by telephone/diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a national panel survey was conducted for the first time throughout the 2011–12 fishing year (Wynne-Jones et al 2014). The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. The national panel survey was repeated during the 2017–18 fishing year using very similar methods to produce directly comparable results (Wynne-Jones et al 2019). Recreational catch estimates from the two national panel surveys are given in Table 5. Note that national panel survey estimates do not include recreational harvest taken under s111 general approvals.

Table 5: Recreational harvest estimates for jack mackerel stocks (Wynne-Jones et al 2014, 2019). Mean fish weights were obtained from boat ramp surveys (Hartill & Davey 2015, Davey et al 2019).

Stock	Year	Method	Number of fish	Total weight (t)	CV
JMA 1	2011–12	Panel survey	101 076	32.2	0.20
	2017–18	Panel survey	62 710	18.6	0.24
JMA 3	2011–12	Panel survey	50	<1	1.01
	2017–18	Panel survey	0	0	–
JMA 7	2011–12	Panel survey	11 194	10.2	0.57
	2017–18	Panel survey	20 026	6.2	0.51

1.3 Customary non-commercial fisheries

Quantitative information on the current level of Māori customary non-commercial catch is not available.

1.4 Illegal catch

There is no information on illegal activity or catch but it is considered to be insignificant.

1.5 Other sources of mortality

There is no information on other sources of mortality.

2. BIOLOGY

The three species of jack mackerel in New Zealand have different geographical distributions, but their ranges partially overlap. *T. novaezelandiae* predominates in waters shallower than 150 m and warmer than 13 °C; it is uncommon south of latitude 42° S. *Trachurus declivis* generally occurs in deeper (but less than 300 m) waters cooler than 16 °C, north of latitude 45° S (Robertson 1978). *Trachurus murphyi* occurs to depths of least 500 m and has a wide latitudinal range (0° at the Galapagos Islands and coastal Ecuador, to south of 40° S off the Chilean coast) (Kawahara et al 1988).

Trachurus murphyi was first described from New Zealand waters in 1987 (Kawahara et al 1988). Its presence was recorded off the south and east coasts of the South Island. Its distribution expanded to off the west coast of the South Island and the North and South Taranaki bights by the late 1980s, reaching the Bay of Plenty in appreciable quantities by 1992, and becoming common off the east coast of Northland by June 1994. However, this extensive distribution has decreased in more recent years and, since the late 1990s, its presence north of Cook Strait has been sporadic with occasional landings in the JMA 1 purse seine fishery north of East Cape and from the JMA 1 inshore trawl fishery south of East Cape. The total range of *T. murphyi* extends along the west coast of South America, across the South Pacific, to the New Zealand EEZ, and into waters off south-eastern Australia.

JACK MACKERELS (JMA)

All species can be caught by bottom trawl, midwater trawl, or by purse seine nets targeting surface schools. The vertical and horizontal movement patterns are poorly understood. Jack mackerels are presumed to be generally off the bottom at night, and surface schools can be quite common during the day.

Jack mackerels have a protracted spring-summer spawning season. *T. novaezelandiae* probably matures at about 26–30 cm fork length (FL) at an age of 3–4 years, and *T. declivis* matures when about 26–30 cm FL at an age of 2–4 years. Spawning occurs in the North and South Taranaki bights, and probably in other areas as well.

The reproductive biology of *T. murphyi* in New Zealand waters is not well understood. Pre- and post-spawning fish have been recorded from the Chatham Rise, Stewart-Snares shelf, Northland east coast, and off Kaikoura in summer, but it is unknown whether there has been any resulting recruitment in New Zealand waters. A study by Taylor (2002a) showed that older size/age groups become increasingly dominant in catches westward from the South American coast, suggesting that an eastward migration of oceanic spawned larvae and juveniles occurs in the South Pacific Ocean.

Horn (1993) fitted von Bertalanffy growth curves for *T. novaezelandiae* and *T. declivis* to samples collected from the central west coast and in the Bay of Plenty between 1974 and 1991. No significant sexual differences in growth were apparent for either species but geographical differences were indicated for *T. novaezelandiae*. Hartill et al (2022) have provided updated estimates of growth parameters for *T. novaezelandiae* in JMA 1. Updated estimates for *T. novaezelandiae* and *T. declivis* in have been provided by Neubauer & Middleton (in prep) using a mixture modelling approach that simultaneously estimated growth parameters and the correct species for otoliths collected by observers in the JMA 7 fishery (see Section 5.1 below). *Trachurus novaezelandiae* and *T. declivis* have moderate initial growth rates that slow after about 6 years. Both species reach a maximum age of 25+ years.

Trachurus murphyi has proven difficult to age with markedly varying growth parameters estimated from fish sampled off South America (Horn & Ó Maolagáin 2021). Using otolith samples from JMA 7 in the period 2006–07 to 2018–19, Horn & Ó Maolagáin (2021) estimate that the mean asymptotic length of male *T. murphyi* in New Zealand is larger than that of females. Initial growth is rapid, slowing at 6–7 years, and *T. murphyi* is a moderately long-lived species with a maximum observed age of 38 years.

The best available estimate of M for *T. novaezelandiae* and *T. declivis* is 0.18 based on the age-frequency distributions of lightly exploited populations in the Bay of Plenty. Assuming $M = 0.18$, estimates of Z made in 1989 suggest that F is less than 0.05 for both endemic species off the central west coast (the main jack mackerel fishing ground). Biological parameters relevant to the stock assessment are shown in Table 6.

Table 6: Estimates of biological parameters.

Fishstock		Estimate	Source
<u>1. Natural mortality (M)</u>			
All		0.18	
	Considered best estimate for both endemic species from all areas.		Horn (1991a)
<u>2. Weight = $a(\text{length})^b$ (Weight in g, length in cm fork length)</u>			
		<u>All</u>	
	a	b	
<i>T. declivis</i>	0.023	2.84	Horn (1991a)
<i>T. novaezelandiae</i>	0.0093	3.10	Hartill et al (2022)
<i>T. murphyi</i>	0.016	2.85	Basten (1981)
<u>3. von Bertalanffy growth parameters</u>			
		<u>All</u>	
	L_∞	k	t_0
<i>T. declivis</i> (JMA 7)	46 cm	0.29	-1.06
<i>T. novaezelandiae</i> (JMA 1)	34.5 cm	0.34	0.1
<i>T. novaezelandiae</i> (JMA 7)	34.5 cm	0.30	-0.72
<i>T. murphyi</i> (JMA 7, females)	50.7 cm	0.231	-0.5 (fixed)
<i>T. murphyi</i> (JMA 7, males)	52.9 cm	0.216	-0.5 (fixed)
			Horn & Ó Maolagáin (2021)

3. STOCKS AND AREAS

There has been no work to investigate the stock boundaries for the native jack mackerels since their introduction to the QMS. For assessment purposes the three jack mackerel species are treated separately where possible.

Differences in growth rate of *T. novaezelandiae* and *T. declivis* between JMA 1 and JMA 7 may indicate stock separation.

There are two possible hypotheses on the stock structure of *T. murphyi* in New Zealand waters: it is either a separate stock established by fish migrating from South America, or part of a single, extensive trans-Pacific stock. Although successful recruitment in New Zealand waters would indicate the establishment of a separate stock, current evidence favours the latter hypothesis with an extensive stock between latitudes 35° and 50° S linking the coasts of Chile and New Zealand across what has been described as ‘the jack mackerel belt’. Few detailed data are available to document the process of range expansion by *T. murphyi* or indicate the relative abundance of the three species in particular areas. As a requirement of the increased TACCs introduced in 1994–95, improvements to jack mackerel catch monitoring were made to provide adequate data for quantifying species composition and relative abundance in JMA 1 and JMA 3.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

This section was updated for the 2022 Fisheries Assessment Plenary based on Fisheries New Zealand data updates for jack mackerel fisheries interaction tables in this section. Fishery interactions are described more fully issue-by-issue in the Aquatic Environment and Biodiversity Annual Review 2021 (Fisheries New Zealand 2021), online at <https://www.mpi.govt.nz/dmsdocument/51472-Aquatic-Environment-and-Biodiversity-Annual-Review-AEBAR-2021-A-summary-of-environmental-interactions-between-the-seafood-sector-and-the-aquatic-environment>.

4.1 Role in the ecosystem

A study of fish assemblages using research trawls suggested that *Trachurus novaezelandiae* is part of an inshore assemblage that prefers shallow northern waters (centred on about 60 m depth and latitude about 38.7° S). All three species overlap spatially, but *T. declivis* is part of a deeper assemblage around central New Zealand (centred on about 130 m and about 40.1° S), and *T. murphyi* occurs deeper still and further south (centred on about 220 m and about 44.7° S) (Francis et al 2002). *T. novaezelandiae* and *T. declivis* range through the water column from surface to the sea floor. The behaviour of *T. murphyi* in New Zealand is less well known but studies off Chile suggest that this species tends to aggregate at night and that this could reflect nocturnal foraging (Bertrand et al 2004, 2006). The effect on the ecosystem of extracting, for example, between 5000 and 10 000 t of jack mackerels from JMA 1 and about 30 000 t from JMA 3 per year over the past decade is unknown.

4.1.1 Trophic interactions

Stevens et al (2011) reported the diet of *T. novaezelandiae* and *T. declivis* from the Bay of Plenty, Northland, and off the west coast South Island to be predominantly euphausiids with fewer amphipods and fish (see also Hurst 1980). Crustaceans (several groups) were the dominant prey of *T. novaezelandiae* in the Hauraki Gulf, with fewer fish and polychaetes (Godfriaux 1968, 1970). The diet of *T. murphyi* from research trawls on shelf areas around New Zealand, mainly down to 500 m depth, included: crustaceans (55%, mainly euphausiids 38%, amphipods 12%, and *Munida* 6%); salps (36%); and teleosts (11% frequency of occurrence in non-empty stomachs, Stevens et al 2011).

Predators of jack mackerels are likely to include many fishes, seabirds, and marine mammals given the relatively high abundance of jack mackerels. The diet of gemfish from research trawls in Southland included *Trachurus* spp. (6% of total, Stevens et al 2011). *T. declivis* and *T. murphyi* were identified from the stomachs of leafscale gulper shark and Plunket’s shark and *T. declivis* from the stomachs of school shark (Dunn et al 2010). The diet of spiny dogfish included scavenged jack mackerel (Dunn et al 2013).

4.2 Bycatch (fish and invertebrates)

Between 2009 and 2011, *T. novaezealandiae* dominated 97% of purse seine landings in JMA 1 (Walsh et al 2012). The estimated proportions by year were 1–17% for *T. declivis*, 0–3% for *T. murphyi*, and 81–99% for *T. novaezealandiae*. There was spatial and temporal heterogeneity in size and abundance; *T. novaezealandiae* dominated landings from the Bay of Plenty throughout the year and large *T. declivis* and *T. murphyi* were common in east Northland during winter (Walsh et al 2016).

Finucci et al (2020) used data from scientific observers and commercial catch-effort returns to estimate the rates and annual levels of fish and invertebrate bycatch and discards in the jack mackerel trawl fisheries, from 2002–03 to 2018–19. Jack mackerel species (*Trachurus* spp.) accounted for 78% of the total estimated catch from trawls targeting jack mackerels between 1 October 2002 and 30 September 2019. The remaining 22% comprised mostly other commercial species, including barracouta (*Thyrsites atun*, 11%), blue mackerel (*Scomber australasicus*, 3.1%), and frostfish (*Lepidopus caudatus*, 3.0%) (Table 7). Over 90% of reported catch was of QMS species, although altogether 370 taxa were identified by observers. Species with notable levels of discards included spiny dogfish (68%), kingfish (50%), porcupine fish (83%), and sunfish (100%).

Table 7: Bycatch and discards from all observer records for the target trawl fishery for jack mackerel from 1 October 2002 to 30 September 2019 for species or species groups with a total catch of 100 kg or more, ordered by decreasing percentage of catch (Finucci et al in 2020).

Species code	Common name	Scientific name	Estimated catch (kg)	% of catch	% discarded
JMA/JMD/JMM/JMN	Jack mackerel	<i>Trachurus declivis</i> , <i>T. murphyi</i> , <i>T. novaezealandiae</i>	279 209.8	77.7	0.0
BAR	Barracouta	<i>Thyrsites atun</i>	40 004.0	11.1	0.1
EMA	Blue mackerel	<i>Scomber australasicus</i>	11 140.8	3.1	0.0
FRO	Frostfish	<i>Lepidopus caudatus</i>	10 776.2	3.0	0.3
RBT	Redbait	<i>Emmelichthys nitidus</i>	8451.9	2.4	0.5
STU	Slender tuna	<i>Allothenus fallai</i>	1057.6	0.3	3.1
SPD	Spiny dogfish	<i>Squalus acanthias</i>	845.6	0.2	68.1
SWA	Silver warehou	<i>Seriola punctata</i>	786.5	0.2	0.0
PIL	Pilchard	<i>Sardinops sagax</i>	747.7	0.2	3.6
RBM	Ray's bream	<i>Brama brama</i>	698.2	0.2	0.0
KIN	Kingfish	<i>Seriola lalandi</i>	682.4	0.2	50.2
WAR	Blue warehou	<i>Seriola brama</i>	525.5	0.1	0.0
SNA	Snapper	<i>Chrysophrys auratus</i>	485.4	0.1	0.3
SDO	Silver dory	<i>Cyttus novaezealandiae</i>	285.2	0.1	1.2
TRE	Trevally	<i>Pseudocaranx georgianus</i>	246.6	0.1	0.0
JDO	John dory	<i>Zeus faber</i>	225.9	0.1	0.0
POP	Porcupine fish	<i>Allomycterus jaculiferus</i>	219.0	0.1	82.7
HOK	Hoki	<i>Macruronus novaezealandiae</i>	193.3	0.1	0.1
GUR	Gurnard	<i>Chelidonichthys kumu</i>	178.0	<0.1	0.1
ATT	Kahawai	<i>Arripis trutta</i> , <i>A. xylabion</i>	160.2	<0.1	0.0
MAK	Mako shark	<i>Isurus oxyrinchus</i>	145.4	<0.1	34.4
NMP	Tarakihi	<i>Nemadactylus macropterus</i> & <i>N. rex</i>	144.9	<0.1	0.2
SUN	Sunfish	<i>Mola mola</i>	136.5	<0.1	100.0
THR	Thresher shark	<i>Alopias vulpinus</i>	129.2	<0.1	100.0

4.3 Incidental capture of protected species (mammals, seabirds, and protected fish)

For protected species, capture estimates presented here include all animals recovered to the deck (alive, injured, or dead) of fishing vessels but do not include any cryptic mortality, e.g., seabirds that are struck by a warp but not brought onboard the vessel (Middleton & Abraham 2007).

4.3.1 Marine mammal captures

Jack mackerel trawlers occasionally catch marine mammals, primarily common dolphin, long-finned pilot whale, and New Zealand fur seal (which are all classified as 'Not Threatened' under the New Zealand Threat Classification System in 2019 (Baker et al 2019)). Between 2002–03 and 2019–20, there were 198 observed captures of whales and dolphins in jack mackerel trawl fisheries: common dolphin (183), long-finned pilot whale (13), dusky dolphin (1), and long-beaked common dolphin (1). Estimated captures for 2002–03 to 2019–20 are shown in Table 8, and show a strong declining trend. Common dolphins were observed captured off the Taranaki coast or off the west coast of the North

Island (Abraham et al 2016, 2021). The 2002–03 to 2017–18 average of the estimated capture rate for common dolphins is 1.5 captures per 100 tows (range 0 to 4.62) in the jack mackerel fishery.

4.3.2 Seabird captures

Annual observed seabird capture rates ranged from 0 to 1.4 per 100 tows in jack mackerel fisheries between 2002–03 and 2019–20 (Abraham & Thompson 2009, Abraham & Thompson 2011, Thompson et al 2013, Abraham et al 2016). Capture rates have fluctuated without obvious trend at this low level (Table 9). Total estimated seabird captures in the jack mackerel trawl fishery varied from 3 to 27 between 2002–03 and 2019–20 (Table 9).

Observed seabird captures since 2002–03 have been mostly prions, shearwaters, and petrels (83 of the 111 observed seabird captures), with 28 observed albatross captures (Table 10). Seabird captures in the jack mackerel fishery have been observed mostly on the Stewart-Snares shelf, off Taranaki, and off the east coast South Island. These numbers should be regarded as only a general guide on the distribution of captures because the numbers are small, and the observer coverage is not uniform across areas and may not be representative.

The jack mackerel target trawl fishery contributes to the total risk posed by New Zealand commercial fishing to seabirds (Table 11). The species to which the fishery poses the most risk is Southern Buller’s albatross; this target fishery posing 0.002 of PST (Table 11). Southern Buller’s albatross was assessed at high risk (Richard et al 2017).

Mitigation methods such as streamer (tori) lines, Brady bird bafflers, warp deflectors, and offal management are used in the jack mackerel trawl fishery. Warp mitigation was voluntarily introduced from about 2004 and made mandatory in April 2006 (Department of Internal Affairs 2006). The 2006 Notice mandated that all trawlers over 28 m in length use a seabird scaring device while trawling (“paired streamer lines”, “bird baffler” or “warp deflector” as defined in the Notice).

Table 8: Number of tows by fishing year and observed common dolphin captures in jack mackerel trawl fisheries, 2002–03 to 2019–20. Annual fishing effort (tows), number of observed tows and observer coverage (%) in jack mackerel trawl fisheries; number of observed captures and observed capture rate (captures per hundred tows) of common dolphin; estimated captures and capture rate of common dolphin (mean and 95% credible interval). Estimates are based on methods described by Abraham et al (2021), available online at <https://protectedspeciescaptures.nz/PSCv6/released/>. Observed and estimated protected species captures in this table derive from the PSC database version PSCV6.

Fishing year	Fishing effort			Obs. captures		Est. captures		Est. capture rate	
	Tows	No. Obs	% obs	Captures	Rate	Mean	95% c.i.	Mean	95% c.i.
2002–03	3 067	346	11.3	21	6.07	141	60-259	4.59	1.96-8.44
2003–04	2 383	152	6.4	17	11.18	99	45-181	4.17	1.89-7.6
2004–05	2 509	558	22.2	21	3.76	85	46-139	3.39	1.83-5.54
2005–06	2 809	709	25.2	2	0.28	12	2-33	0.43	0.07-1.17
2006–07	2 711	802	29.6	11	1.37	55	23-102	2.04	0.85-3.76
2007–08	2 652	818	30.8	20	2.44	42	24-70	1.60	0.9-2.64
2008–09	2 169	813	37.5	11	1.35	23	11-43	1.05	0.51-1.98
2009–10	2 406	786	32.7	4	0.51	17	4-42	0.69	0.17-1.75
2010–11	1 882	593	31.5	7	1.18	53	18-108	2.82	0.96-5.74
2011–12	2 032	1 548	76.2	5	0.32	7	5-13	0.32	0.25-0.64
2012–13	2 213	1 940	87.7	15	0.77	16	15-20	0.71	0.68-0.9
2013–14	2 447	2 187	89.4	28	1.28	29	28-35	1.21	1.14-1.43
2014–15	1 750	1 512	86.4	19	1.26	21	19-28	1.21	1.09-1.6
2015–16	1 544	1 383	89.6	2	0.14	3	2-7	0.17	0.13-0.45
2016–17	1 407	1 024	72.8	0	0.00	1	0-5	0.05	0-0.36
2017–18	1 688	1 474	87.3	0	0.00	0	0-4	0.03	0-0.24
2018–19	1 627	1 278	78.5	0	0.00				
2019–20	1 747	1 352	77.4	0	0.00				

JACK MACKERELS (JMA)

Table 9: Number of tows by fishing year and observed seabird captures in jack mackerel trawl fisheries, 2002–03 to 2019–20. No. obs, number of observed tows; % obs, percentage of tows observed; Rate, number of captures per 100 observed tows. Estimates are based on methods described by Abraham & Richard (2020) and are available online at <https://protectedspeciescaptures.nz/PSCv6/released/>. Observed and estimated protected species captures in this table derive from the PSC database version PSCV6.

Fishing year	Fishing effort			Obs. captures		Est. captures		Est. capture rate	
	Tows	No. Obs	% obs	Captures	Rate	Mean	95% c.i.	Mean	95% c.i.
2002–03	3 067	346	11.3	4	1.16	22	10-42	0.72	0.33-1.37
2003–04	2 383	152	6.4	0	0.00	7	1-17	0.29	0.04-0.71
2004–05	2 509	558	22.2	8	1.43	16	9-27	0.63	0.36-1.08
2005–06	2 809	709	25.2	0	0.00	20	5-45	0.70	0.18-1.6
2006–07	2 711	802	29.6	1	0.12	8	2-19	0.31	0.07-0.7
2007–08	2 652	818	30.8	1	0.12	9	2-20	0.32	0.08-0.75
2008–09	2 169	813	37.5	6	0.74	14	7-26	0.63	0.32-1.2
2009–10	2 406	786	32.7	9	1.15	15	9-27	0.63	0.37-1.12
2010–11	1 882	593	31.5	7	1.18	15	8-28	0.78	0.43-1.49
2011–12	2 032	1 548	76.2	6	0.39	9	6-18	0.47	0.3-0.89
2012–13	2 213	1 940	87.7	26	1.34	27	26-31	1.22	1.17-1.4
2013–14	2 447	2 187	89.4	7	0.32	7	6-13	0.30	0.25-0.53
2014–15	1 750	1 512	86.4	12	0.79	14	12-22	0.81	0.69-1.26
2015–16	1 544	1 383	89.6	6	0.43	7	6-12	0.47	0.39-0.78
2016–17	1 407	1 024	72.8	4	0.39	6	4-13	0.45	0.28-0.92
2017–18	1 688	1 474	87.3	10	0.68	11	10-16	0.67	0.59-0.95
2018–19	1 627	1 278	78.5	3	0.23	5	3-10	0.28	0.18-0.61
2019–20	1 747	1 352	77.4	1	0.07	3	1-9	0.16	0.06-0.52

Table 10: Number of observed seabird captures in jack mackerel trawl fisheries, 2002–03 to 2019–20, by species and area. Observed protected species captures in this table derive from the PSC database version PSCV6.

Species	Risk category	Taranaki	WCNI	Chatham Rise	Stewart-Snares shelf	ECSI	WCSI	Total
Salvin's albatross	High	0	0	0	0	3	0	3
Southern Buller's albatross	High	0	0	1	3	2	0	6
New Zealand white-capped albatross	Medium	5	0	0	10	4	0	19
Total albatrosses	–	5	0	1	13	9	0	28
Westland petrel	High	0	0	0	0	0	1	1
White-chinned petrel	Negligible	0	0	1	32	5	0	38
Sooty shearwater	Negligible	1	0	0	10	2	0	13
Common diving petrel	Negligible	0	0	0	1	0	1	2
White-faced storm petrels	Negligible	0	3	1	0	0	0	4
Australasian gannet	Negligible	1	0	0	0	0	0	1
Fairy prion	Negligible	5	0	0	1	1	0	7
Cape petrels	–	2	0	0	0	0	1	3
Cook's petrel	–	1	0	0	0	0	0	1
Fulmar prion	–	10	0	0	0	0	0	10
Grey-backed storm petrel	–	1	0	1	0	0	0	2
Large seabird	–	1	0	0	0	0	0	1
Total other birds	–	22	3	3	44	8	3	83

Table 11: Risk ratio of seabirds predicted by the level two risk assessment for the jack mackerel and all fisheries included in the level two risk assessment, 2006–07 to 2016–17, showing seabird species with a risk ratio of at least 0.001 of PST (Richards et al 2020). The risk ratio is an estimate of aggregate potential fatalities across trawl and longline fisheries relative to the Population Sustainability Threshold, PST (from Richard et al 2017, where full details of the risk assessment approach can be found). The DOC threat classifications are shown (Robertson et al 2017 at <http://www.doc.govt.nz/documents/science-and-technical/nztc19entire.pdf>).

Species name	PST (mean)	Risk ratio			DOC Threat Classification
		MAC risk ratio	Total	Risk category	
Southern Buller's albatross	1 368.4	0.002	0.392	High	At Risk: Naturally Uncommon
New Zealand white-capped albatross	10 900.3	0.001	0.353	High	At Risk: Declining

4.3.3 Protected fish species captures

Mobulid rays (spinetail devilrays, *Mobula mobular*, and manta rays, *Mobula birostris*, both protected since 2010 under the Wildlife Act 1953) occur mainly in north-eastern North Island waters during summer and could potentially be caught in purse seine nets along the north-east coast of North Island. However, observers monitoring mackerel purse seine fisheries (coverage 0–17.8% per year, 2002–18) have not reported any captures of mobulid rays to date.

4.4 Benthic interactions

Jack mackerel are taken using trawls that are sometimes fished on or near the seabed. The spatial extent of seabed contact by trawl fishing gear in New Zealand's EEZ and Territorial Sea has been estimated and mapped in numerous studies for trawl fisheries targeting deepwater species (Baird et al 2011, Black et al 2013, Black & Tilney 2015, Black & Tilney 2017, Baird & Wood 2018, and Baird & Mules 2019, 2021a, 2021b), species in waters shallower than 250 m (Baird et al. 2015, Baird & Mules 2021a, 2021b), and all trawl fisheries combined (Baird & Mules 2021a, 2021b). The most recent assessment of the deepwater trawl footprint was for the period 1989–90 to 2018–19 (Baird & Mules 2021b).

During 1989–90 to 2018–19, about 55 100 bottom-contacting jack mackerel trawls were reported on TCEPRs and ERS (Baird & Mules 2021b); this represents about 1200–3300 tows in most years up to 2013–14 and an average of 880 tows per year from 2014–15 to 2018–19. The total footprint generated from these tows was estimated at about 46 697 km². This footprint represented coverage of 1.1% of the seafloor of the combined EEZ and the Territorial Sea areas; 3.4% of the 'fishable area', that is, the seafloor area open to trawling, in depths of less than 1600 m. For the 2018–19 fishing year, 870 jack mackerel bottom-contacting tows had an estimated footprint of 2825 km² which represented coverage of 0.1% of the EEZ and Territorial Sea and 0.2% of the fishable area (Baird & Mules 2021b).

The overall trawl footprint for jack mackerel (1989–90 to 2018–19) covered 16% of the seafloor in < 200 m, 6% of 200–400 m seafloor, and < 0.05% of the 400–600 m seafloor (Baird & Mules 2021b). The jack mackerel footprint contacted 1%, 0.1%, and < 0.01% of those depth ranges, respectively, in 2018–19 (Baird & Mules 2021b). The BOMECS class C (off the west coast of the North Island) had the highest proportion of area covered by the jack mackerel footprint in 2018–19 (4%), with the remainder of the footprint covering about 0.3% of the 61 000 km² of class E (Stewart-Snares shelf) and 0.2% of the 138 550 km² of class H (Chatham Rise) (Baird & Mules 2021b).

Trawling for jack mackerel with some or all of the gear contacting the bottom, like trawling for other species, is likely to have effects on benthic community structure and function (e.g., Rice 2006) and there may be consequences for benthic productivity (e.g., Jennings et al 2001, Hermsen et al 2003, Hiddink et al 2006, Reiss et al 2009). These consequences are not considered in detail here but are discussed in the Aquatic Environment and Biodiversity Annual Review 2021 (Fisheries New Zealand 2021).

4.5 Other considerations

4.5.1 Spawning disruption

Fishing may disrupt spawning activity or success. Canadian research carried out on Atlantic cod (*Gadus morhua*) concluded that "Cod exposed to a chronic stressor are able to spawn successfully, but there appears to be a negative impact of this stress on their reproductive output, particularly through the production of abnormal larvae" (Morgan et al 1999). Morgan et al (1997) also reported disruption of a spawning shoal of Atlantic cod: "Following passage of the trawl, a 300-m-wide "hole" in the aggregation spanned the trawl track. Disturbance was detected for 77 min after passage of the trawl." There have been no specific studies for jack mackerel in New Zealand waters, but information on the timing and location of spawning and fishing exists. *T. declivis* and *T. novaezelandiae* are serial spawners with a protracted spring-summer spawning season (Hurst et al 2000). *T. murphyi* appears to spawn from late winter through to summer (Horn 1991b, Hurst et al 2000). The JMA 7 trawl fishery has peaks of catch and effort in spring-summer (October–March) and in winter (April–September) (McKenzie 2008), the former overlapping with spawning. Most of the purse seine catch from the Bay of Plenty is taken in September–October, but an increasing proportion has been caught in November–December since 2005–06 (Walsh et al 2012), also overlapping the spring-summer spawning.

4.5.2 Habitat of particular significance to fisheries management

Habitat of particular significance for fisheries management (HPSFM) does not have a policy definition (Ministry for Primary Industries 2016), although work is underway to generate one. Studies of potential relevance have identified areas of importance for spawning and juveniles (Hurst et al 2000). *T. declivis* spawning was found to be common on the southwest and northwest North Island outer shelf, and moderate to high abundance of juveniles was recorded from northwest North Island, Hauraki Gulf, and Bay of Plenty outer shelf. *T. novaezealandiae* spawning was found to be common on the southwest and northwest inner and outer shelf of the North Island, and moderate to high abundance of juveniles was recorded from Hauraki Gulf and Bay of Plenty inner and outer shelf, East Cape inner shelf, and Tasman Bay/Golden Bay. *T. murphyi* spawning was found to be common on the southwest outer shelf and only low abundance of juveniles was recorded from the outer Southland shelf and at 300–600 m on the Chatham Rise.

4.5.3 Genetic effects

Fishing and environmental changes, including those caused by climate change or pollution, could alter the genetic composition or diversity of a species. There are no known studies of the genetic diversity of jack mackerels in New Zealand.

4.5.4 Marine heatwave

The effects of the marine heatwave on jack mackerel fisheries that was experienced in New Zealand waters in the summer months of 2017–18 are unknown.

5. STOCK ASSESSMENT

Stock assessments for jack mackerel are complicated by the reporting and management of three species under a single code.

5.1 Challenger, Central West, and Auckland West (JMA 7)

Preliminary stock assessments for *T. declivis* and *T. novaezealandiae* in JMA 7 were undertaken in 2007 (McKenzie 2009). These relied on CPUE indices (McKenzie, 2008) where an average species composition was applied to individual tows within a stratum. Webber & Starr (2022) noted that applying average species proportions across relatively large strata to individual tows, instead of using the actual species proportions in the tow, will reduce the variability in a CPUE analysis and will likely lead to biased coefficient estimates because the model does not have access to the real variation by species relative to the covariate values. As a result, these previous CPUE analyses and the preliminary assessments are considered unreliable.

Webber & Starr (2022) also identified a range of concerns about the data from observer sampling in JMA 7. Consequently, a set of CPUE standardisations of all three species combined was carried out, for positive catches of JMA only.

Species proportion sampling

The key concern identified by Webber & Starr (2022) was that observer sampling for catch composition in the JMA 7 trawl fishery had not always been implemented as specified in the observer manual. There were reports which suggested that observers often took a single sample at the beginning of processing instead of several throughout the processing procedure. They suggested that if stratification of the catch within the trawl persisted through to when the catch was processed, then sampling at a single time would result in biased estimates of species composition.

The existing data collection procedures did not retain sub-sample information for jack mackerel length-frequency samples by species, so there was no way of using these data to assess whether the fish comprising a sample were collected at one time or as a series of sub-samples. To address concerns about potential stratification in the sampled catch, two observer trips were carried out where observers took up to four speciated length-frequency sub-samples of jack mackerel during the processing of catches and recorded the sub-sample data separately. Middleton (2022) found differences between some sub-sample and tow-level estimates of species proportions, but these were not related to the time during processing when sub-samples were taken. As a result, it was concluded by the WG that there was only

a limited risk of bias in tow-level estimates of species composition arising from sampling at a single point during processing.

Misidentification of species

Misidentification of the three jack mackerel species in observer sampling of otoliths for ageing has been apparent in catch-at-age data from the JMA 7 fishery for some time. Based on the subset of fish aged, Saunders et al (2022) reported that the rate of misidentification by observers was at least 8% for *T. novaezelandiae* and at least 3% for *T. declivis*. While potential misidentification was sometimes apparent in length sampling data (e.g., a small number of suspiciously large *T. novaezelandiae*), instances of misidentification tended to stand out most obviously in plots of length at age, from age sampling. However, even in these data, misidentification was only clear for older fish (i.e., when the growth curves were obviously separated) whereas some level of misidentification would also be expected for smaller fish.

Moore et al (in press) have demonstrated that analyses of otolith shape showed promise for confirming the species for fish selected for ageing. To address this issue using the existing data, Neubauer & Middleton (in prep) developed a Bayesian mixture model of growth and fitted this model to age-length data for jack mackerels sampled in JMA 7 from 1995 to 2020. The 18 987 observations had a measured fork length, an age from otolith reading, and an observer-assigned species. The mixture model estimated the von Bertalanffy growth parameters for each species (combined sexes) and made predictions of the true species for each observation, accounting for potential ageing error (Figure 3).

The mixture model results have been used to provide updated estimates of growth parameters for *T. novaezelandiae* and *T. declivis* in JMA 7 (Table 6) but also provided information on the probability that the observer-assigned species was correct (Figure 4). The latter estimates allowed an investigation of the sensitivity of species-specific CPUE indices to species misidentification (see below). For each fish, the model provided an estimate of the probability that a particular fish should be assigned to a particular species. Because the growth curve for *T. declivis* lies between that of the other two species, the model tended to be less confident about species assignment for these fish. Using a threshold for likely misidentification of < 50% probability (that the observer-assigned species was correct, given the model) for *T. novaezelandiae* and *T. murphyi*, and < 25% for *T. declivis*, led to estimated misidentification rates of 8.3% for *T. declivis*, 5.9% for *T. novaezelandiae* and < 1% for *T. murphyi*.




Figure 3: Estimated growth curves for jack mackerels in JMA 7 (JMN, *T. novaezelandiae*; JMD, *T. declivis*; JMM, *T. murphyi*). Solid lines indicate the median and 95% interval for predicted size at age, in comparison to the growth curves based on parameters previously reported in the Plenary (dashed lines). The three panels group the data according to the observer assigned species, and points are coloured by the model predicted species (JMN = green, JMD = blue, JMM = red), and shaded by the estimated probability of that prediction.

JACK MACKERELS (JMA)




Figure 4: Top: estimated growth curves for jack mackerel species in JMA 7 plotted as agreed age, from the otolith readings, at measured size and with panels grouping the data by observer-assigned species (JMN, *T. novaezealandiae*; JMD, *T. declivis*; JMM, *T. murphyi*). Solid lines indicate the median and 95% interval for predicted size at age, in comparison to the growth curves based on parameters previously reported in the Plenary (dashed lines), and points are coloured by the model predicted species (JMN = green, JMD = blue, JMM = red), and shaded by the estimated probability of that prediction. Bottom: median (points) and 95% interval (lines) for the probability that the observer assigned species label is correct, given the model predicted species, grouped by 1 cm length intervals.

CPUE

Species-specific CPUE indices for *T. declivis* and *T. novaezealandiae* in JMA 7 were revisited in 2022, following the work to address concerns about the impacts of observer sampling on tow-level species composition estimates (Middleton 2022).

For observed tows with length-frequency sampling of jack mackerels, the observer sampling was used to estimate the proportion of each species in the tow, by weight after applying the standard length-weight relationships (Table 6) to estimate the weight of each sampled fish. These proportions were then applied to disaggregate the observer-recorded total weight of jack mackerels in the tow into a tow-by-tow catch by species (Figure 5). Where an observer had recorded catch weights using a species-specific code (i.e., JMD, JMM, or JMN), these weights were first recoded to the generic jack mackerel code (JMA) and aggregated at the tow level.

Standardised CPUE indices were calculated using standard maximum-likelihood methods for each species individually. Data were restricted to tows where the observer recorded that jack mackerel was the target species. Two series were considered: (i) midwater tows only (MW series), which allowed annual indices to be estimated from 1999, and (ii) using both bottom and midwater tows (TR series), which allowed indices to be estimated from 1991.

For *T. novaezealandiae* tows were restricted to the Taranaki Bight and west coast of the North Island (Statistical Areas 037, 039, 040, 041, 042, and 045) while the datasets for *T. declivis* also included tows off the South Island west coast (Statistical Areas 034, 035, and 036). Core fleet selection required vessels to have at least one observed trip in a minimum of four fishing years for the midwater series, but—due to lower observer coverage in the early 1990s—this was reduced to a minimum of two years for the longer TR series in order to maximise the number of observations. Stepwise selection on the basis of AIC, retaining terms only if they explained an additional 1% of the deviance, was used to select covariates, with fishing year, vessel, month, and statistical area offered as categorical terms (with

method added for the TR series) and duration, start time, seabed depth, headline height, and fishing speed offered as continuous terms using natural spline smoothers.




Figure 5: Annual boxplots of tow-level proportions of each jack mackerel species (JMN, *T. novaezealandiae*; JMD, *T. declivis*; JMM, *T. murphyi*) from observer sampling in JMA 7. Tows are plotted if the length frequency sample contained at least 30 fish.

Trachurus declivis occurred in a high proportion of tows, reaching 100% in some years and precluding estimation of an occurrence model. For *T. novaezealandiae* both an occurrence and positive catch models were fitted, and a combined index was calculated. For the MW series, vessel was not selected as an explanatory variable for catches of either species, an outcome which was likely attributable to the fact that the core fleets comprised only the very similar BATM-class vessels. The temporal trends evident in the indices for *T. declivis* and *T. novaezealandiae* were similar among the statistical areas included in the models for each species, indicating that it was unlikely that there was an area-year interaction in this analysis.

The following sensitivity analyses were considered:

- the impact of including or excluding tows where the length-frequency sample size was unusually small (less than 30 fish);
- the effect of excluding trips with apparent observer misidentifications of jack mackerel species;
- whether the subset of observer data with length-frequency sampling produced a biased index relative to the entire fleet data; and
- whether there was any evidence for 'effort creep' (i.e., increased catching efficiency over time) in the core fleet vessels.

None of these sensitivities indicated significant problems with the standardised CPUE analyses, and the MW series beginning in 1999 was accepted by the Deepwater Working Group as indices of abundance for *T. declivis* and *T. novaezealandiae* in JMA 7 (Figure 6). The longer TR (MW+BT) series, beginning

in 1991, was not considered sufficiently reliable to provide a longer term series, primarily because the level of observer coverage was much lower in the early 1990s but also because of the fact that the vessels fishing in the early 1990s had largely exited the fishery by the end of that decade with almost no overlap with the post-1999 fishery.




Figure 6: Standardised catch per unit effort (CPUE) indices for *T. declivis* (JMD) and *T. novaezealandiae* (JMN) in JMA 7 calculated from observer data using the subset of tows where length-frequency sampling permitted estimation of catch by species. A positive catch index is presented for JMD and a combined index (binomial/positive) index for JMN. The series are scaled to have a geometric mean of 1.

Establishing B_{MSY} -compatible reference points

In 2022, the Deepwater Working Group (DWWG) considered that there was little information available to allow equating a particular period with a specific stock status for either of the two native jack mackerel species. Commercial fisheries became established in JMA 7 in the 1970s, but catches were lower than in recent decades, and there was no indication that the stocks were depleted when jack mackerels entered the QMS. The DWWG noted that the Harvest Strategy Standard Operational Guidelines propose that, in cases where the relationship between CPUE and abundance can be assumed to be more or less proportional, it was reasonable to select an appropriate historical period when both CPUE and catches were relatively high and stable and to use this CPUE level as a target (conceptual B_{MSY}). Catches in JMA 7 were considered to have been stable since the early-mid 2000s, with the establishment of the BATM fleet. Periods of stable abundance were identified as 2004–2011 for *T. declivis* and 2006–2012 for *T. novaezealandiae*. Noting that both species had subsequently increased in abundance while combined species catches remained stable, the Plenary adopted the geometric mean CPUE in each of these periods as the target reference level for the appropriate species (equated to 35% B_0 for these medium-productivity species). The soft limit (20% B_0) was set at 4/7 of the mean value for each period and the hard limit (10% B_0) was set at 2/7 of the mean value by period.

Catch at Age

Catch-at-age distributions for all three jack mackerel species JMA 7 are available for 1989–90, 1990–91, 1995–96, 2004–05, and 2005–06 to 2020–21 (e.g., Saunders et al 2022). However, the DWWG noted that apparent strong year classes evident in the length -frequency distributions did not clearly translate to the age distributions, which were prepared using a standard method developed in 2011 with a stratification updated in 2016. The Working Group considered that the methodology for preparing age frequency distributions would require re-evaluation before these were used for stock assessment.

Future Research Considerations – JMA7

- Extend the mixture modelling of length at age data to allow evaluation of whether significant sex or area specific differences in growth are present for any species.
- Explore using otolith shape analysis to validate the potential misidentification from the mixture modelling.
- Explore whether there are temporal trends in species misidentification.
- Evaluate the methodology and stratification, and provide time series of catch by species (e.g., bootstrapping vs model-based approaches) and consider whether the estimation of species composition and CPUE can be usefully integrated.
- Review the methodology used to create annual age frequency distributions, including the stratification used.
- Review available data on JMA 7 from west coast inshore trawl surveys to determine what additional information on trends in abundance and recruitment this can provide.
- Explore available information to investigate whether there are any temporal patterns in mean size for each species, and consider how these can inform our understanding of the stocks.
- Given the likely correlation between observer samples collected within the same trip, explore the representativeness of observer sampling at the trip level rather than event level.

5.2 Biomass estimates

Estimates of current biomass are not available.

5.3 Other yield estimates and stock assessment results

For *T. declivis* and *T. novaezelandiae* catch-at-age proportions were used to estimate instantaneous total mortality Z for the years 2006–07 to 2008–09 in JMA 7 (Smith 2011). The methods used to produce species-specific catch-at-age distributions for JMA 7 are not currently considered sufficiently reliable to use these results.

5.4 Other factors

Trachurus murphyi has been known at times to comprise a substantial proportion of the purse seine catches in the area between Cook Strait and Kaikōura, in the Bay of Plenty, and off the east Northland coast, although the proportion of this component has declined considerably since the late 1990s. *Trachurus murphyi* has also been an important component of the west coast North Island jack mackerel trawl fishery but has declined in recent years. Thus, there has been a contraction in the range of this species in New Zealand waters, although it is unknown yet whether this represents a decrease in its overall abundance here. The effect of *T. murphyi* on the range and abundance of the other two species is unknown.

Aerial sightings data were used to produce a time series of relative abundance indices for jack mackerel. The time series covered the period from the beginning of the purse seine fishery in 1976 to 1993. It indicated an increase in abundance in JMA 1 from the early 1990s, and, although the result was not as clear, a similar trend in JMA 3 and JMA 7. These increases were attributed to the invasion of *T. murphyi*.

The validity of this early aerial sightings abundance index is uncertain. Further analysis of these data has been the focus of considerable effort in recent years and the Northern Inshore Working Group has not yet accepted revised abundance indices due to data and model concerns.

The stipulation that catches in JMA 1 and JMA 3 above the original TACs (5970 t and 2700 t, respectively) be accounted for by increases in *T. murphyi* only, is a method of managing this species independently of the other two. This approach was introduced as a means of maintaining stocks of the endemic species while allowing exploitation of increased stocks of *T. murphyi* resulting from its invasion.

The increase in *T. novaezelandiae* catch has predominantly occurred within the Bay of Plenty fishery area. There has been a small decrease in the length of fish caught from the fishery since 2006–07 to 2008–09, although it is unknown whether the decline in fish size is attributable to an increase in fishing mortality rates, changes in fishing operation, or variation in annual recruitment. Age composition data are available for the *T. novaezelandiae* catch from 2006–07 to 2008–09, but age-based sampling was

JACK MACKERELS (JMA)

discontinued due to the relatively high inter-annual variability in the age compositions, with the fishery targeting size classes based on market demand.

Future Research Considerations for all stocks

- The instruction that observers (for JMA 3 and JMA 7) routinely aim to spread the collection of length-frequency samples of jack mackerels throughout the processing period should be maintained; the observer data base should be extended to capture information about the sub-sampling achieved from the processing of each tow.
- Consider the introduction of simpler species composition sampling (proportion by weight) for observed tows that are not selected for length-frequency sampling.
- Review of JMD and JMN stock structure and evaluation of the occurrence and extent of *T. murphyi* incursions into NZ waters, including a national scale characterisation of JMA. This should also review older documents describing the JMA fisheries and stocks.

6. STATUS OF THE STOCKS

Stock Structure Assumptions

Separate stocks of *T. declivis* (JMD) and *T. novaezealandiae* (JMN) are assumed to occur in each jack mackerel Quota Management Area (other than JMA 10). *T. murphyi* (JMM) in New Zealand waters is not considered to represent an independent biological stock.

- **JMA 1**

Stock Status	
Year of Most Recent Assessment	-
Reference Points	Target(s): Not established but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not established
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status
-

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	An index for JMA 1 is not available at this time. Recent work and discussions concerning the use of aerial sightings data for annual relative abundance indices concluded that the inter-annual variation was too great for these data to provide a reliable index.
Recent Trend in Fishing Mortality or Proxy	-
Trends in other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	It is not known whether catches at the level of the current TACCs or recent catch levels are sustainable in the long-term.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown

Probability of Current Catch or TACC causing Overfishing to continue or to commence	-
---	---

Assessment Methodology and Evaluation	
Assessment Type	Level 3—Qualitative Evaluation: Fishery characterisation with evaluation of fishery trends (e.g., catch, effort and nominal CPUE, length frequency information) – there is no agreed index of abundance
Assessment Method	-
Assessment Dates	Latest assessment: - Next assessment: Unknown
Overall assessment quality rank	-
Main data inputs (rank)	Species proportions estimates
Data not used (rank)	
Changes to Model Structure and Assumptions	-
Major Sources of Uncertainty	-

Qualifying Comments
-

Fishery Interactions
JMA 1 catches are primarily taken by targeted purse seine. Because jack mackerel often occur in mixed schools with kahawai, particularly towards the end of the fishing year, this can inhibit jack mackerel targeting in this fishery at this time. Interactions with other species are currently being characterised.

- **JMA 3**

Stock Status	
Year of Most Recent Assessment	-
Reference Points	Management Target: 40% B_0 Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: Not established
Status in relation to Target	Unknown
Status in relation to Limits	Unknown
Status in relation to Overfishing	-

Historical Stock Status Trajectory and Current Status
-

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	-
Recent Trend in Fishing Intensity or Proxy	-
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	It is not known whether catches at the level of the current TACCs or recent catch levels are sustainable in the long-term.

JACK MACKERELS (JMA)

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	-

Assessment Methodology and Evaluation	
Assessment Type	Level 4: Low information evaluation—there are only data on catch and TACC, with no other fishery indicators. Catch is qualified with species proportions estimates from MPI observer data. Some length frequency information is available.
Assessment Method	-
Assessment Dates	Latest assessment: - Next assessment: -
Overall assessment quality rank	
Main data inputs (rank)	-
Data not used (rank)	-
Changes to Model Structure and Assumptions	-
Major Sources of Uncertainty	-


Qualifying Comments
-

Fishery Interactions
JMA 3 catches are primarily taken by midwater trawl. Non-target species captured in this fishery include barracouta and redbait. Incidental captures of protected species have been recorded for New Zealand fur seals and cetaceans. Trawls on or near the seabed interact with benthic habitats.

- **JMA 7 - *T. declivis***

Stock Status	
Year of Most Recent Assessment	2023
Assessment Runs Presented	Event resolution CPUE index from observer data from jack mackerel target midwater trawls with length frequency sampling (JMD7 MW observer)
Reference Points	Management Target: 35% B_0 ; the geometric mean CPUE for the period 2004–2011 (a conceptual proxy for B_{MSY}) Soft Limit: 20% B_0 ; scaled from management target Hard Limit: 10% B_0 ; scaled from management target Overfishing threshold: not established
Status in relation to Target	Likely (> 60%) to be at or above the target
Status in relation to Limits	Very Unlikely (< 10%) to be below both the soft and hard limits
Status in relation to Overfishing	Unknown

Historical Stock Status Trajectory and Current Status



Positive catch standardised catch per unit effort (CPUE) index for *T. declivis* in JMA 7 (calculated from observer data using the subset of tows where length frequency sampling permits estimation of catch by species) relative to the agreed conceptual reference points. The green, orange, and red dashed lines represent the interim target, soft limit, and hard limit, respectively.

Fishery and Stock Trends

Recent Trend in Biomass or Proxy	CPUE increased from 2010 to 2017, was stable to 2021, but decreased in 2022 to a level comparable with the mid-2010 period
Recent Trend in Fishing Intensity or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis

Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Current catch Soft Limit: Unlikely (< 40%) in the short term Hard Limit: Unlikely (< 40%) in the short term
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation

Assessment Type	Level 2 – Partial quantitative stock assessment	
Assessment Method	CPUE analysis	
Assessment Dates	Latest assessment: 2023	Next assessment: 2025
Overall assessment quality rank	1 – High quality	
Main data inputs (rank)	- Observer catch records and length frequency samples	1 – High Quality
Data not used (rank)	- Catch at age	2 – Medium or Mixed Quality: the methods used for producing

JACK MACKERELS (JMA)


		scaled catch at age distributions require review
Changes to Model Structure and Assumptions	- Species-specific CPUE has replaced a combined species analysis	
Major Sources of Uncertainty	-	

Qualifying Comments
-

Fishery Interactions
JMA 7 catches are primarily taken by midwater trawl. Availability of ACE for kingfish and snapper potentially influences targeting in some sub-areas. Incidental captures of protected species have been recorded for New Zealand fur seals and cetaceans.

- **JMA 7 - *T. novaezealandiae***

Stock Status	
Year of Most Recent Assessment	2023
Assessment Runs Presented	Event resolution CPUE index from observer data from jack mackerel target midwater trawls with length frequency sampling (JMN7 MW observer)
Reference Points	Management Target: the geometric mean CPUE for the period 2006–2012 (a conceptual proxy for B_{MSY}) Soft Limit: 20% B_0 ; scaled from management target Hard Limit: 10% B_0 ; scaled from management target Overfishing threshold: not established
Status in relation to Target	Very Likely (> 90%) to be at or above the target
Status in relation to Limits	Very Unlikely (< 10%) to be below both the soft and hard limits
Status in relation to Overfishing	Unknown



Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE increased from 2010 to 2018 and has fluctuated without trend to 2022
Recent Trend in Fishing Intensity or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Unknown
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Current catch Soft Limit: Unlikely (< 40%) in the short term Hard Limit: Unlikely (< 40%) in the short term
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown

Assessment Methodology and Evaluation		
Assessment Type	Level 2 – Partial quantitative stock assessment	
Assessment Method	CPUE analysis	
Assessment Dates	Latest assessment: 2023	Next assessment: 2025
Overall assessment quality rank	1 – High quality	
Main data inputs (rank)	- Observer catch records and length frequency samples	1 – High Quality
Data not used (rank)	- Catch at age	2 – Medium or Mixed Quality: the methods used for producing scaled catch at age distributions require re-evaluation
Changes to Model Structure and Assumptions	- Species-specific CPUE has replaced a combined species analysis	
Major Sources of Uncertainty	-	

Qualifying Comments
-

Fishery Interactions
JMA 7 catches are primarily taken by midwater trawl. Availability of ACE for kingfish and snapper potentially influences targeting in some sub-areas. Incidental captures of protected species have been recorded for New Zealand fur seals and cetaceans.

7. FOR FURTHER INFORMATION

- Abraham, E R; Berkenbusch, K; Richard, Y; Thompson, F (2016) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 2002–03 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 169*. 205 p.
- Abraham, E R; Richard, Y (2017) Summary of the capture of seabirds in New Zealand commercial fisheries, 2002–03 to 2013–14. *New Zealand Aquatic Environment and Biodiversity Report No. 184*. 88 p.
- Abraham, E R; Richard, Y (2018) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report No. 197*. 97 p.
- Abraham, E R; Richard, Y (2020) Estimated capture of seabirds in New Zealand trawl and longline fisheries, to 2017–18. *New Zealand Aquatic Environment and Biodiversity Report No. 249*. 86 p.
- Abraham, E R; Tremblay-Boyer, L; Berkenbusch, K (2021) Estimated captures of New Zealand fur seal, common dolphin, and turtles in New Zealand commercial fisheries, 2002–03 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 258*. 98 p.
- Abraham, E R; Thompson, F N (2009) Capture of protected species in New Zealand trawl and longline fisheries, 1998–99 to 2006–07 *New Zealand Aquatic Environment and Biodiversity Report No. 32*. 197 p.

JACK MACKERELS (JMA)

- Abraham, E R; Thompson, F N (2011) Summary of the capture of seabirds, marine mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2008–09. *New Zealand Aquatic Environment and Biodiversity Report No. 80*.
- Abraham, E R; Thompson, F N; Berkenbusch, K (2013) Estimated capture of seabirds in New Zealand trawl and longline fisheries, 2002–03 to 2010–11. (Unpublished Final Research Report for Ministry for Primary Industries project PRO2010-01 held by Fisheries New Zealand, Wellington.)
- Abraham, E R; Thompson, F N; Oliver, M D (2010) Summary of the capture of seabirds, mammals, and turtles in New Zealand commercial fisheries, 1998–99 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 45*. 149 p.
- Anderson, O F; Edwards, C T T; Roux, M-J (2017) Fish and invertebrate bycatch and discards in New Zealand jack mackerel trawl fisheries from 2002–03 until 2013–14. *New Zealand Aquatic Environment and Biodiversity Report No. 177*. 71 p.
- Baird, S J (2001) Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1998–99. *New Zealand Fisheries Assessment Report 2001/14*. 43 p.
- Baird, S J (2004a) Estimation of the incidental capture of seabird and marine mammal species in commercial fisheries in New Zealand waters, 1999–2000. *New Zealand Fisheries Assessment Report 2004/41*. 56 p.
- Baird, S J (2004b) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2000–01. *New Zealand Fisheries Assessment Report 2004/58*. 63 p.
- Baird, S J (2004c) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2001–02. *New Zealand Fisheries Assessment Report 2004/60*. 51 p.
- Baird, S J (2005) Incidental capture of seabird species in commercial fisheries in New Zealand waters, 2002–03. *New Zealand Fisheries Assessment Report 2005/2*. 50 p.
- Baird, S J; Hewitt, J E; Wood, B A (2015) Benthic habitat classes and trawl fishing disturbance in New Zealand waters shallower than 250 m. *New Zealand Aquatic Environment and Biodiversity Report No. 144*. 184 p.
- Baird, S J; Mules, R (2019) Extent of bottom contact by New Zealand commercial trawl fishing for deepwater Tier 1 and Tier 2 target species determined using CatchMapper software, fishing years 2008–17. *New Zealand Aquatic Environment and Biodiversity Report No. 229*. 106 p.
- Baird, S J; Mules, R (2021a) Extent of bottom contact by commercial fishing activity in New Zealand waters, for 1989–90 to 2017–18. *New Zealand Aquatic Environment and Biodiversity Report No. 259*. 143 p.
- Baird, S J; Mules, R (2021b) Extent of bottom contact by commercial trawling and dredging in New Zealand waters, 1989–90 to 2018–19. *New Zealand Aquatic Environment and Biodiversity Report No. 260*. 157 p.
- Baird, S J; Smith, M H (2007) Incidental capture of New Zealand fur seals (*Arctocephalus forsteri*) in commercial fisheries in New Zealand waters, 2003–04 to 2004–05. *New Zealand Aquatic Environment and Biodiversity Report No. 14*. 98 p.
- Baird, S J; Wood, B A (2012) Extent of coverage of 15 environmental classes within the New Zealand EEZ by commercial trawling with seafloor contact. *New Zealand Aquatic Environment and Biodiversity Report No. 89*. 43 p.
- Baird, S J; Wood, B A (2018) Extent of bottom contact by New Zealand commercial trawl fishing for deepwater Tier 1 and Tier 2 target fishstocks, 1989–90 to 2015–16. *New Zealand Aquatic Environment and Biodiversity Report No. 193*. 102 p.
- Baird, S J; Wood, B A; Bagley, N W (2011) Nature and extent of commercial fishing effort on or near the seafloor within the New Zealand 200 n. mile Exclusive Economic Zone, 1989–90 to 2004–05. *New Zealand Aquatic Environmental and Biodiversity Report No. 73*. 143 p.
- Baker, C S; Boren, L; Childerhouse, S; Constantine, R; van Helden, A; Lundquist, D; Rayment, W; Rolfe, J R (2019) Conservation status of New Zealand marine mammals, 2019. *New Zealand Threat Classification Series 29*. Department of Conservation, Wellington. 18 p.
- Baker, C S; Chilvers, B L; Childerhouse, S; Constantine, R; Currey, R; Mattlin, R; van Helden, A; Hitchmough, R; Rolfe, J (2016) Conservation status of New Zealand marine mammals, 2013. *New Zealand Threat Classification Series 14*. Department of Conservation, Wellington. 18 p.
- Baker, C S; Chilvers, B L; Constantine, R; DuFresne, S; Mattlin, R H; van Helden, A; Hitchmough, R (2010) Conservation status of New Zealand marine mammals (suborders Cetacea and Pinnipedia), 2009. *New Zealand Journal of Marine and Freshwater Research 44*: 101–115.
- Ballara, S L; Anderson, O F (2009) Fish discards and non-target fish catch in the trawl fisheries for arrow squid and scampi in New Zealand waters. *New Zealand Aquatic Environment and Biodiversity Report No. 38*. 102 p.
- Basten, J C (1981). The fishery for jack mackerel (*Trachurus murphyi*) off Northern Chile. Theses, Dissertations and Student Research Papers (College of Earth, Ocean and Atmospheric Sciences), School of Oceanography, Oregon State University, June 1981. 65 p
- Bertrand, A; Barbieri, M A; Córdova, J; Hernández, C; Gómez, F; Leiva, F (2004) Diel vertical behaviour, predator-prey relationships, and occupation of space by jack mackerel (*Trachurus murphyi*) off Chile. *ICES Journal of Marine Science 61*: 1105–1112.
- Bertrand, A; Barbieri, M A; Gerlotto, F; Leiva, F; Córdova, J (2006) Determinism and plasticity of fish schooling behaviour as exemplified by the South Pacific jack mackerel *Trachurus murphyi*. *Marine Ecology Progress Series 311*: 145–156.
- Black, J; Tilney, R (2015) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–1990 to 2010–2011. *New Zealand Aquatic Environment and Biodiversity Report No. 142*. 56 p.
- Black, J; Tilney, R (2017) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–90 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 176*. 65 p.
- Black, J; Wood, R; Berthelsen, T; Tilney, R (2013) Monitoring New Zealand's trawl footprint for deepwater fisheries: 1989–1990 to 2009–2010. *New Zealand Aquatic Environment and Biodiversity Report No. 110*. 57 p.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. (Unpublished New Zealand Fisheries Assessment Research Report for the Ministry of Fisheries Project REC2000-01 held by Fisheries New Zealand, Wellington.) 92 p.
- Boyd, R O; Reilly, J L (2004) 1999/2000 National Marine Recreational Fishing Survey: harvest estimates. (Unpublished New Zealand Fisheries Assessment Research Report for the Ministry of Fisheries Project REC9803 held by Fisheries New Zealand, Wellington.) 28 p.
- Bradford, E (1996) A comparison of the 1993–94 diary and boat ramp surveys of recreational fishing in the Ministry of Fisheries North region New Zealand Fisheries Assessment Research Document 96/5. 21 p. (Unpublished report held by NIWA library.)
- Bradford, E (1997) Estimated recreational catches from Ministry of Fisheries North region marine recreational fishing surveys, 1993–94. New Zealand Fisheries Assessment Research Document 1997/7. 16 p. (Unpublished document held by NIWA library, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16. 27 p. (Unpublished report held by NIWA library.)
- Bradford, E; Taylor P R (1995) Trends in pelagic fish abundance from aerial sightings data. New Zealand Fisheries Assessment Research Document 1995/8. 60 p. (Unpublished document held by NIWA library, Wellington.)
- Brothers, N; Duckworth, A R; Safina, C; Gilman, E L (2010) Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PLoS One 5*: e12491. Doi: 10.1371/journal.pone.0012491.
- Chapman, D G; Robson, D S (1960) The analysis of a catch curve. *Biometrics 16*: 354–368.
- Davey, N; Hartill, B; Carter, M (2019) Mean weight estimates for recreational fisheries in 2017–18. *New Zealand Fisheries Assessment Report 2019/25*. 32 p.

- Department of Internal Affairs (2006) Seabird Scaring Devices – Circular Issued Under Authority of the Fisheries (Commercial Fishing) Amendment Regulations 2006 (No. F361). *New Zealand Gazette 6 April 2006*: 842–846.
- Dunn, M R; Stevens, D W; Forman, J S; Connell, A (2013) Trophic Interactions and Distribution of Some Squaliforme Sharks, Including New Diet Descriptions for *Deania calcea* and *Squalus acanthias*. *PLoS ONE 8(3)*: e59938. Doi:10.1371/journal.pone.0059938.
- Dunn, M R; Szabo, A; McVeagh, M S; Smith, P J (2010) The diet of deepwater sharks and the benefits of using DNA identification of prey. *Deep-Sea Research I 57*: 923–930.
- Elizarov, A A; Grechina, A S; Botenev, B N; Kuzetsov, A N (1993) Peruvian jack mackerel, *Trachurus symmetricus*, in the open waters of the South Pacific. *Journal of Ichthyology 33(3)*: 86–104.
- Finucci, B; Anderson, O F; Edwards, C T T (2020) Non-target fish and invertebrate catch and discards in New Zealand jack mackerel trawl fisheries from 2002–03 to 2018–19. *New Zealand Aquatic Environment and Biodiversity Report No. 279*. 81 p.
- Francis, M P; Hurst, R J; McArdle, B; Bagley, N W; Anderson, O F (2002) New Zealand demersal fish assemblages. *Environmental Biology of Fishes 65*: 215–234.
- Fisheries New Zealand (2021) Aquatic Environment and Biodiversity Annual Review 2021. Compiled by the Aquatic Environment Team, Fisheries Science and Information, Fisheries New Zealand, Wellington, New Zealand. 779 p.
- Godfriaux, B L (1968) The food and feeding relationships of snapper (*Chrysophrys auratus*) and some other fish-species trawled in Hauraki Gulf. (Unpublished MSc (Hons) thesis, Victoria University of Wellington.)
- Godfriaux, B L (1970) Food of predatory demersal fish in Hauraki Gulf. II. Five fish species associated with snapper. *New Zealand Journal of Marine and Freshwater Research 4*: 248–266.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*. 37 p.
- Hartill, B; Middleton, D A J; Walsh, C; Spong, K; Ó Maolagáin, C (2022) Catch-at-age sampling of *Trachurus novaezelandiae* from JMA 1 in 2019–20 and 2020–21. *New Zealand Fisheries Assessment Report 2022/37*. 16 p.
- Hermesen, J M; Collie, J S; Valentine, P C (2003) Mobile fishing gear reduces benthic megafaunal production on Georges Bank. *Marine Ecology Progress Series 260*: 97–108.
- Hiddink, J G; Jennings, S; Kaiser, M J; Queiros, A M; Duplisea, D E; Piet, G J (2006) Cumulative impacts of seabed trawl disturbance on benthic biomass, production, and species richness in different habitats. *Canadian Journal of Fisheries and Aquatic Sciences 63*: 721–36.
- Horn, P L (1991a) Assessment of jack mackerel stocks off the central west coast, New Zealand, for the 1990–91 fishing year. New Zealand Fisheries Assessment Research Document 1991/6. 14 p. (Unpublished report held in NIWA library, Wellington.)
- Horn, P L (1991b) Trawl survey of jack mackerels (*Trachurus* spp.) off the central west coast, New Zealand, February–March 1990. *New Zealand Fisheries Technical Report No. 28*. 39 p.
- Horn, P L (1993) Growth, age structure, and productivity of jack mackerels (*Trachurus* spp.) in New Zealand waters. *New Zealand Journal of Marine and Freshwater Research 27*: 145–155.
- Hurst, R J (1980) Studies on the life cycle of some New Zealand Anisakidae (Nematoda). Unpublished PhD thesis, Victoria University of Wellington, New Zealand. 212 p.
- Hurst, R J; Stevenson, M L; Bagley, N W; Griggs, L H; Morrison, M A; Francis, M P (2000) Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand coastal fish. (Unpublished Final Research Report for Ministry of Fisheries Research Project ENV1999/03 held by Fisheries New Zealand, Wellington.)
- Jennings, S; Dinmore, T A; Duplisea, D E; Warr, K J; Lancaster, J E (2001) Trawling disturbance can modify benthic production processes. *Journal of Animal Ecology 70*: 459–475.
- Jones, J B (1988) Jack mackerels. New Zealand Fisheries Assessment Research Document 1988/19. 19 p. (Unpublished report held by NIWA library, Wellington.)
- Jones, J B (1990) Jack mackerels (*Trachurus* spp.) in New Zealand waters. *New Zealand Fisheries Technical Report No. 23*. 28 p.
- Kawahara, S; Uozumi, Y; Yamada, H (1988) First record of a carangid fish, *Trachurus murphyi* from New Zealand. *Japanese Journal of Ichthyology 35(2)*: 212–214.
- Langley, A D; Middleton, D A J (2019) Determination of the species composition of the JMA 1 purse seine catch. NINSWG-2019-04. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Langley, A D; Middleton, D A J; Wilson, O L (2016) Species composition of the jack mackerel (genus *Trachurus*) catch from the JMA 1 purse seine fishery, 2005/06 to 2013/14. *New Zealand Fisheries Assessment Report 2016/11*. 33 p.
- Leathwick, J R; Rowden, A; Nodder, S; Gorman, R; Bardsley, S; Pinkerton, M; Baird, S J; Hadfield, M; Currie, K; Goh, A (2012) A Benthic-optimised Marine Environment Classification (BOMECE) for New Zealand waters. *New Zealand Aquatic Environment and Biodiversity Report No. 88*. 54 p.
- McKenzie, A (2008) Standardised CPUE analyses for *Trachurus declivis* and *Trachurus novaezelandiae* in the JMA 7 jack mackerel fishery to 2004–05. *New Zealand Fisheries Assessment Report 2008/46*. 36 p.
- McKenzie, A (2009). Preliminary stock assessments for *Trachurus declivis* and *T. novaezelandiae* in the JMA 7 jack mackerel fishery to 2005–06. *New Zealand Fisheries Assessment Report 2009/28*. 39 p.
- MacKenzie, D; Fletcher, D (2006) Characterisation of seabird captures in commercial trawl and longline fisheries in New Zealand 1997/98 to 2003/04. Final Research Report for ENV2004/04, held by Ministry of Fisheries, New Zealand. 102 p. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Middleton, D A J (in prep) Characterisation of the JMA 1 fishery and species composition of the jack mackerel catch from the JMA 1 purse seine fishery, 2014/15 to 2019/20. Draft New Zealand Fisheries Assessment Report
- Middleton, D A J (2022) An evaluation of species composition sampling in the JMA 7 trawl fishery. *New Zealand Fisheries Assessment Report 2022/20*. 39 p.
- Middleton, D A J; Abraham, E R (2007) The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. (Unpublished Final Research Report for research project IPA2006/02 held by Fisheries New Zealand, Wellington.)
- Ministry for Primary Industries (2016) Aquatic Environment and Biodiversity Annual Review 2016. Compiled by the Fisheries Management Science. Ministry for Primary Industries, Wellington, New Zealand. 790 p.
- Moore, B R; Ó Maolagáin, C; Spong, K; Saunders, R J (in press) Commercial catch sampling for species proportion, sex, length, and age of jack mackerels in JMA 7 in the 2020–21 fishing year, with a summary of all available datasets. New Zealand Fisheries Assessment Report.
- Morgan, M J; DeBlois, E M; Rose, G A. (1997) An observation on the reaction of Atlantic cod (*Gadus morhua*) in a spawning shoal to bottom trawling. Conference - Symposium on the Biology and Ecology of Northwest Atlantic Cod, St. John's, NF (Canada), 24–28 Oct 1994.
- Morgan, M J; Wilson, C E; Crim, L W (1999) The effect of stress on reproduction in Atlantic cod. *Journal of Fish Biology 54(3)*: 477–488.
- Neubauer, P. & Middleton, D. A. J. (in prep) Bayesian mixture models for simultaneous estimation of growth parameters and species assignment for species that are frequently mis-identified.
- Reiss, H; Greenstreet, S P R; Seibe, K; Ehrlich, S; Piet, G J; Quirijns, F; Robinson, L; Wolff, W K; Kronke, I (2009) Effects of fishing disturbance on benthic communities and secondary production within an intensively fished area. *Marine Ecology Progress Series 394*: 201–213.

JACK MACKERELS (JMA)

- Rice, J (2006) Impacts of Mobile Bottom Gears on Seafloor Habitats, Species, and Communities: A Review and Synthesis of Selected International Reviews. Canadian Science Advisory Secretariat Research Document 2006/057. 35 p. (Available from http://www.dfo-mpo.gc.ca/CSAS/Csas/DocREC/2006/RES2006_057_e.pdf.)
- Richard, Y; Abraham, E R (2013) Risk of commercial fisheries to New Zealand seabird populations. *New Zealand Aquatic Environment and Biodiversity Report No. 109*. 58 p.
- Richard, Y; Abraham, E R (2015) Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2012–13. *New Zealand Aquatic Environment and Biodiversity Report No. 162*. 85 p
- Richard, Y; Abraham, E R; Berkenbusch, K (2017). Assessment of the risk of commercial fisheries to New Zealand seabirds, 2006–07 to 2014–15. *New Zealand Aquatic Environment and Biodiversity Report No. 191*. 133 p.
- Richard, Y; Abraham, E R; Berkenbusch, K (2020) Estimated capture of seabirds in New Zealand trawl and longline fisheries, to 2017–18. *New Zealand Aquatic Environment and Biodiversity Report No. 249*. 90 p.
- Robertson, D A (1978) The New Zealand jack mackerel fishery. In Habib, G. and Roberts, P.E. (Eds.). Proceedings of the pelagic fisheries conference, July 1977. *FRD Occasional Publication No. 15*. pp. 43–47.
- Robertson, H A; Baird, K; Dowding J E; Elliott, G P; Hitchmough, R A; Miskelly, C M; McArthur, N; O'Donnell, C F J; Sagar, P M; Scofield, R P; Taylor, G A (2017) Conservation status of New Zealand birds, 2016. *New Zealand Threat Classification Series 19*.
- Rohan, V M; Mittinty, M; Taylor, P R (2006) Aspects of estimating species proportions in the jack mackerel fishery: JMA 7 estimates using a Bayesian approach and examination of sampling bias in JMA 1 resulting from automated grading. Final Research Report for Specific Objective 7 of project JMA2004/01: Stock monitoring of jack mackerels. PELWG06/09. 26 p. (Unpublished report held by Ministry for Primary Industries, Wellington.)
- Saunders, R J; Ó Maolagáin, C; Hulston D; Spong K (2022) Commercial catch sampling for species proportion, sex, length, and age of jack mackerels in JMA 7 in the 2019–20 fishing year, with a summary of all available data sets. *New Zealand Fisheries Assessment Report 2022/52*. 26 p.
- Smith, M H (2011) Catch curves for JMA species in JMA 7. DWWG-2011/27. (Unpublished report held by Fisheries New Zealand, Wellington.) 6 p.
- Stevens, D W; Hurst, R J; Bagley, N W (2011) Feeding habits of New Zealand fishes: a literature review and summary of research trawl database records 1960 to 2000. *New Zealand Aquatic Environment and Biodiversity Report No. 85*. 218 p.
- Taylor, P R (2000) Species composition and seasonal variability in commercial catches and aerial sightings of jack mackerel, *Trachurus declivis*, *T. symmetricus murphyi*, and *T. novaezelandiae*, in JMA 1, JMA 3, and JMA 7. *New Zealand Fisheries Assessment Report 2000/45*. 25 p
- Taylor, P R (2002a) Stock structure and population biology of the Peruvian jack mackerel, *Trachurus symmetricus murphyi*. *New Zealand Fisheries Assessment Report 2002/21*. 78 p.
- Taylor, P R (2002b) Species composition and seasonality of jack mackerel (*Trachurus declivis*, *T. symmetricus murphyi*, and *T. novaezelandiae*) in commercial catches from JMA 1, 3, and 7 during 1998–99 and 1999–2000, with a summary of biological information from 1990–91 to 1999–2000. *New Zealand Fisheries Assessment Report 2002/51*. 72 p.
- Taylor, P R (2004) Species composition and seasonal variability in commercial catches of jack mackerel (*Trachurus declivis*, *T. symmetricus murphyi*, and *T. novaezelandiae*) in JMA 1, JMA 3, and JMA 7 during 2000–01. *New Zealand Fisheries Assessment Report 2004/28*. 22 p.
- Taylor, P R (2008) Factors affecting fish size and landed volumes in the purse-seine and TCEPR charter-boat fisheries in 2004–05 and 2005–06. *New Zealand Fisheries Assessment Report 2008/32*. 17 p.
- Taylor, P R; Horn, P L; O'Maolagáin, C (2012) Commercial catch sampling for species proportion, sex, length, and age of jack mackerels in JMA 1 in the 2006–07, 2007–08 and 2008–09 fishing years. *New Zealand Fisheries Assessment Report 2012/48*. 36 p.
- Taylor, P R; Julian, K A (2008) Species composition and seasonal variability in commercial catches of jack mackerel (*Trachurus declivis*, *T. murphyi* and *T. novaezelandiae*) in JMA 1, 3, and 7 during 2004–05. *New Zealand Fisheries Assessment Report 2008/25*. 24 p.
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94. New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Thompson, F N; Abraham, E R (2009) Dolphin bycatch in New Zealand trawl fisheries, 1995–96 to 2006–07. *New Zealand Aquatic Environment and Biodiversity Report No. 36*. 24 p.
- Thompson, F N; Abraham, E R; Oliver, M D (2010) Estimation of fur seal bycatch in New Zealand trawl fisheries, 2002–03 to 2007–08. *New Zealand Aquatic Environment and Biodiversity Report No. 56*. 39 p.
- Thompson, F N; Berkenbusch, K; Abraham, E R (2013) Marine mammal bycatch in New Zealand trawl fisheries, 1995–96 to 2010–11. *New Zealand Aquatic Environment and Biodiversity Report No. 105*. 73 p.
- Tuck, I; Cole, R; Devine, J (2009) Ecosystem indicators for New Zealand fisheries. *New Zealand Aquatic Environment and Biodiversity Report No. 42*. 188 p.
- Walsh, C; Bian, R; McKenzie, J M; Miller, A; Spong, K; Armiger, H (2012) Species composition and seasonal variability in commercial purse-seine catches of jack mackerel (*Trachurus declivis*, *T. murphyi*, and *T. novaezelandiae*) in JMA 1 between January 2009 and September 2011. *New Zealand Fisheries Assessment Report 2012/46*. 41 p.
- Walsh, C; Bian, R; McKenzie, J; Spong, K; Armiger, H (2016) Species composition and seasonal variability in commercial purse-seine catches of jack mackerel (*Trachurus declivis*, *T. murphyi*, and *T. novaezelandiae*) in JMA 1 between January 2011 and September 2013. *New Zealand Fisheries Assessment Report 2016/10*. 44 p.
- Webber, D N; Starr, P J (2022) Characterisation and CPUE from 1989–90 to 2018–19 of jack mackerel off the west coast of New Zealand (JMA 7). *New Zealand Fisheries Assessment Report 2022/06*. 65 p
- Wynne-Jones, J; Gray, A; Heinemann, A; Hill, L; Walton, L (2019) National Panel Survey of Marine Recreational Fishers 2017–2018. *New Zealand Fisheries Assessment Report 2019/24*. 104 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.