


GROPER (HPB)

(*Polyprion oxygeneios*, *Polyprion americanus*)
Hāpuku, Moeone



1. FISHERY SUMMARY

1.1 Commercial fisheries

Both groper species, *Polyprion oxygeneios* (hāpuku) and *P. americanus* (bass), occur in shelf and slope waters of the New Zealand mainland and offshore islands, from the Kermadecs to the Auckland Islands. The groper fishery takes both species, but in different proportions by region, depth, fishing method and season, and these have changed over time. Reported landings generally do not distinguish between species, and published data combine them. In earlier years, bluenose (*Hyperoglyphe antarctica*) landings were sometimes also combined with groper. In this document, groper is used as collective term for hāpuku and bass. Historical estimated and recent reported groper landings and TACCs are shown in Tables 1, 2 and 3, while Figure 1 shows the historical and recent landings and TACC values for the main groper stocks.

Table 1: Reported total New Zealand landings (t) of groper from 1948 to 1983.

Year	Landings	Year	Landings	Year	Landings	Year	Landings
1948	1 665	1957	1 368	1966	1 222	1975	1 422
1949	1 969	1958	1 532	1967	1 314	1976	1 512
1950	1 709	1959	1 310	1968	1 073	1977	1 942
1951	1 396	1960	1 223	1969	1 122	1978	1 488
1952	1 430	1961	1 203	1970	1 499	1979	2 078
1953	1 403	1962	1 173	1971	1 346	1980	2 435
1954	1 364	1963	1 194	1972	1 120	1981	2 379
1955	1 305	1964	1 370	1973	1 312	1982	2 218
1956	1 399	1965	1 249	1974	1 393	1983	2 511

Reported foreign catches are included from 1974.
Source: Fisheries data.

The main fishery comprises a number of domestic fishers working small to medium sized vessels - longliners, setnetters and trawlers, at a variety of depths (according to method) out to 500 m (Paul 2002a). Over 90% of early (to 1950) total groper catches were taken by longline. Trawl catches rose from 5–10% during this period to 20–30% by the late 1970s. A setnet fishery developed in the late 1970s and early 1980s, mainly at Kaikoura, taking 14% in 1983 and then subsequently declining. From 1950 to the mid-1980s, line-fishing took 70–80% of the catch. After the introduction of the QMS in 1986, the proportion of the catch taken by lines appeared to drop.

The Cook Strait region has always supported the main groper fishery, followed by the Canterbury Bight; both show the same slow decline from 1949 to 1986 (equivalent regional data from subsequent years are not available). Northland, Bay of Plenty and Hawke Bay fisheries developed at different rates during the 1960s and 1970s. In most other areas, the groper fishery has been small and/or variable.

The first recorded landings of about 1 500 t in 1936 were typical of the range of catches (1000–2000 t) from then until 1978. After a decrease during the war when effort was restricted, landings in the total fishery slowly declined from almost 2000 t in 1949 to about 1300 t in the mid-1970s. They then increased sharply to 2700 t in 1983–84 (Tables 1 and 2). Figure 1 shows the historical landings and TACC values for the main HPB stocks.

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	HPB 1	HPB 2	HPB 3	HPB 4	HPB 5	HPB 7	HPB 8
1931–32	231	0	207	2	130	13	13
1932–33	201	276	242	0	91	98	53
1933–34	198	330	173	25	99	127	53
1934–35	204	304	212	57	115	106	56
1935–36	179	201	146	70	33	109	33
1936–37	129	445	115	12	29	156	50
1937–38	119	523	315	15	29	148	52
1938–39	90	621	479	8	75	156	50
1939–40	118	502	409	12	59	155	43
1940–41	120	444	286	9	54	142	41
1941–42	80	450	302	10	46	150	44
1942–43	69	287	315	9	44	115	35
1943–44	59	316	271	8	42	112	42
1944	55	332	286	9	60	188	117
1945	106	311	271	3	65	173	128
1946	154	326	409	7	83	229	190
1947	98	401	563	5	142	250	175
1948	111	450	526	11	140	275	151
1949	174	498	547	7	142	364	236
1950	141	423	555	9	116	281	184
1951	104	353	381	19	102	267	171
1952	112	368	373	35	100	281	162
1953	105	349	431	33	96	252	137
1954	156	355	397	32	77	235	112
1955	142	351	419	26	82	197	88
1956	106	404	439	32	114	227	77
1957	133	380	419	23	92	246	76
1958	115	473	458	30	96	250	109
1959	147	406	350	54	68	198	87
1960	122	394	331	48	100	150	77
1961	135	369	348	50	82	139	80
1962	163	355	298	40	101	142	75
1963	197	315	321	56	75	159	71
1964	224	397	365	41	76	193	74
1965	212	368	325	68	48	176	52
1966	213	415	315	4	49	163	62
1967	229	448	275	0	49	228	85
1968	139	357	264	0	67	176	70
1969	197	454	220	0	30	138	84
1970	259	670	239	2	54	175	97
1971	191	562	289	4	41	181	78
1972	401	370	188	0	29	99	33
1973	419	481	215	0	30	136	32
1974	356	457	208	2	43	140	72
1975	227	315	213	18	55	379	62
1976	183	220	350	107	101	445	37
1977	277	301	265	87	47	575	113
1978	348	470	194	10	59	280	67
1979	620	487	355	147	113	276	71
1980	956	376	414	40	199	315	105
1981	693	373	457	59	218	381	166
1982	957	336	402	26	133	256	46

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns: Data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data includes both foreign and domestic landings.

Table 3: Reported landings (t) of groper by Fishstock from 1983–84 to present and actual TACCs (t) from 1986–87 to present. QMS data from 1986–present. * FSU data, includes exploratory permit catches. [Continued on next page]

Fishstock FMA (s)	HPB 1		HPB 2		HPB 3		HPB 4		HPB 5	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	974	-	493	-	505	-	55	-	395	-
1984–85*	642	-	388	-	418	-	52	-	228	-
1985–86*	569	-	270	-	391	-	53	-	126	-
1986–87	238	360	179	210	260	270	42	300	131	410
1987–88	248	388	202	219	268	286	43	315	91	414
1988–89	231	405	187	248	259	294	49	315	70	425
1989–90	310	465	179	263	283	318	40	322	127	430
1990–91	350	480	225	263	311	326	77	323	120	436
1991–92	277	480	252	263	298	326	58	323	112	446
1992–93	375	480	273	264	299	327	68	323	128	446
1993–94	363	480	287	264	306	330	90	323	147	446
1994–95	334	481	259	264	274	335	149	323	161	451
1995–96	335	481	214	264	321	335	173	323	144	451
1996–97	331	481	234	264	301	335	131	323	149	451
1997–98	375	481	260	266	329	335	88	323	91	451
1998–99	433	481	256	266	348	335	121	323	97	451
1999–00	471	481	229	266	385	335	66	323	169	451
2000–01	450	481	220	266	381	335	45	323	188	451
2001–02	427	481	226	266	343	335	82	323	169	451
2002–03	442	481	273	266	350	335	79	323	212	451
2003–04	433	481	281	266	335	335	87	323	166	451
2004–05	433	481	263	266	371	335	147	323	208	451
2005–06	425	481	280	266	406	335	185	323	167	451
2006–07	483	481	245	266	394	335	222	323	157	451
2007–08	439	481	253	266	341	335	241	323	138	451
2008–09	415	481	253	266	391	335	138	323	153	451
2009–10	374	481	249	266	358	335	213	323	152	451
2010–11	371	481	222	266	322	335	231	323	128	451
2011–12	312	481	193	266	336	335	265	323	158	451
2012–13	314	481	206	266	337	335	156	323	140	451
2013–14	319	481	224	266	301	335	169	323	143	451
2014–15	314	481	180	266	280	335	156	323	126	451
2015–16	270	481	143	266	315	335	144	323	143	451
2016–17	287	481	162	266	342	335	152	323	156	451
2017–18	276	481	159	266	344	335	142	323	158	451
2018–19	283	481	173	266	347	335	137	323	167	451
2019–20	226	481	126	266	299	335	181	323	161	451
2020–21	180	481	170	266	284	335	197	323	195	451
2021–22	134	140	72	80	264	335	116	323	189	451
2022–23	133	140	84	80	207	335	108	323	124	451
		HPB 7		HPB 8		HPB 10		Total		
		7		8		10		Total		
		Landings		Landings		Landings		Landings		
		TACC		TACC		TACC		TACC		
1983–84*	174	-	46	-	0	-	2 698	-		
1984–85*	207	-	33	-	0	-	2 039	-		
1985–86*	199	-	25	-	0	-	1 697	-		
1986–87	149	210	35	60	0	10	1 036	1 830		
1987–88	158	215	66	76	0	10	1 076	1 923		
1988–89	132	226	39	78	1	10	968	2 001		
1989–90	119	229	43	80	0	10	1 098	2 117		
1990–91	128	235	48	80	23#	10	1 282	2 153		
1991–92	175	235	50	80	83#	10	1 319	2 163		
1992–93	186	236	62	80	22#	10	1 405	2 165		
1993–94	193	236	69	80	0	10	1 455	2 167		
1994–95	192	236	68	80	0	10	1 437	2 179		
1995–96	214	236	78	80	0	10	1 479	2 179		
1996–97	186	236	71	80	15	10	1 418	2 179		
1997–98	147	236	60	80	33#	10	1 406	2 181		
1998–99	218	236	78	80	3#	10	1 562	2 181		
1999–00	165	236	65	80	0#	10	1 561	2 181		
2000–01	171	236	64	80	0#	10	1 519	2 181		
2001–02	204	236	62	80	< 1	10	1 514	2 181		
2002–03	233	236	72	80	0	10	1 661	2 181		
2003–04	239	236	66	80	0	10	1 607	2 181		
2004–05	240	236	80	80	0	10	1 742	2 181		
2005–06	207	236	56	80	0	10	1 728	2 181		
2006–07	206	236	66	80	0	10	1 773	2 181		
2007–08	195	236	44	80	0	10	1 651	2 181		
2008–09	207	236	71	80	0	10	1 628	2 181		
2009–10	221	236	66	80	0	10	1 633	2 181		
2010–11	191	236	80	80	0	10	1 543	2 181		
2011–12	173	236	61	80	0	10	1 187	2 181		
2012–13	209	236	75	80	0	10	1 436	2 181		
2013–14	182	236	63	80	0	10	1 401	2 181		
2014–15	132	236	67	80	0	10	1 254	2 181		
2015–16	148	236	73	80	0	10	1 236	2 181		
2016–17	141	236	69	80	0	10	1 309	2 181		
2017–18	110	236	61	80	0	10	1 250	2 181		
2018–19	105	236	47	80	0	10	1 260	2 182		

Table 3 [Continued]:

Fishstock FMA(s)	HPB 7		HPB 8		HPB 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
2019–20	79	236	33	80	0	10	1 105	2 182
2020–21	78	236	43	80	0	10	1 147	2 182
2021–22	79	236	40	80	0	10	894	1 655
2022–23	79	97	45	55	0	10	780	1 491

Landings and TACCs for all Fishstocks are given in Table 3. Total landings of groper were relatively stable throughout the mid-1990s, remaining below 1500 t until 1998–99. From 1999–2000 onwards, landings have generally ranged between 1200 t and 1700 t. Although the TACC in HPB 3 has been exceeded in some years, landings have generally remained within the quotas for individual Fishstocks and have never exceeded the total TACC.

For the 1991–92 fishing year the conversion factor for headed and gutted groper was increased from 1.40 to 1.45, for fish landed in this state (about 75% of the total), which resulted in a reduction in removals from the stock of 3.5% for the same nominal quota.




Figure 1: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB 1 (Auckland) and HPB 2 (Central East). [Continued on next two pages]




Figure 1 [Continued]: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB HPB 3 (South East Coast), HPB 4 (Chatham Rise), HPB 5 (Southland, Sub-Antarctic).




Figure 1 [Continued]: Total reported landings and TACC for the seven main HPB stocks. From top to bottom: HPB 7 (Challenger), and HPB 8 (Central).

1.2 Recreational fisheries

Groper are taken by handline and setline, and to a lesser extent by setnets. Recreational catch estimates from surveys undertaken in the 1990s are given in Tables 4–6.

Table 4: Estimated number of groper harvested by recreational fishers by Fishstock and survey, the corresponding estimated survey harvest and the estimated Fishstock harvest. Surveys were carried out in different years in the MAF Fisheries regions: South in 1991–92, Central in 1992–93 and North in 1993–94 (Teirney et al 1997).

Fishstock	Survey	Total		Survey harvest (t)
		Number	CV (%)	
HPB 1	North	22 000	17	190–220
HPB 2	North	1 000	-	5–10
HPB 2	Central	10 000	37	45–85
HPB 3	Central	3 000	-	10–30
HPB 3	South	4 000	40	10–30
HPB 5	Central	7 000	36	20–40
HPB 5	South	2 000	-	5–15
HPB 7	Central	12 000	40	45–115
HPB 8	Central	1 000	-	5–10

Table 5: Results of a national diary survey of recreational fishers in 1996, indicating estimated number of groper harvested by recreational fishers by Fishstock and the corresponding harvest tonnage. The mean weights used to convert numbers to catch weight are considered the best available estimates. Estimated harvest is also presented as a range to reflect the uncertainty in the estimates (from Bradford 1998).

Fishstock	Number	CV (%)	Harvest	Point
	caught		range (t)	Estimate (t)
HPB 1	11 000	17	40–60	49
HPB 2	23 000	22	75–125	100
HPB 3	4 000	-	-	-
HPB 5	2 000	-	-	-
HPB 7	9 000	-	-	-
HPB 8	< 500	-	-	-

Table 6: Results of the 1999–2000 national diary survey of recreational fishers (Dec 1999–Nov 2000). Estimated number of groper harvested by recreational fishers by Fishstock, and the corresponding harvest tonnage. Estimated harvest is presented as a range to reflect the uncertainty in the estimates (Boyd & Reilly 2002).

Fishstock	Number	CV (%)	Harvest	Point
	caught		range (t)	estimate (t)
HPB 1	60 000	39	209–476	342
HPB 2	56 000	33	307–608	457
HPB 3	52 000	50	97–293	195
HPB 5	6 000	70	14–80	47
HPB 7	17 000	37	79–172	125
HPB 8	2 000	67	6–32	19

The harvest estimates provided by telephone-diary surveys between 1993 and 2001 are no longer considered reliable for various reasons. A Recreational Technical Working Group concluded that these harvest estimates should be used only with the following qualifications: a) they may be very inaccurate; b) the 1996 and earlier surveys contain a methodological error; and c) the 2000 and 2001 estimates are implausibly high for many important fisheries. In response to these problems and the cost and scale challenges associated with onsite methods, a National Panel Survey was conducted for the first time throughout the 2011–12 fishing year. The panel survey used face-to-face interviews of a random sample of 30 390 New Zealand households to recruit a panel of fishers and non-fishers for a full year (Wynne-Jones et al 2019). The panel members were contacted regularly about their fishing activities and harvest information collected in standardised phone interviews. The national panel survey was repeated during the 2017–18 and 2022–23 fishing years using very similar methods to produce directly comparable results (Wynne-Jones et al 2019; Heinemann & Gray, in prep). Recreational catch estimates from the three national panel surveys are given in Table 7. Note that national panel survey estimates do not include recreational harvest taken on charter vessel trips or under s111 general approvals. For the 2022–23 survey hapuku (HAP) and bass (BAS) were recorded separately (Table 7a).

Table 7: Recreational harvest estimates for groper stocks (Wynne-Jones et al 2014, 2019, Heinemann & Gray, in prep). Mean weights from boat ramp surveys (Hartill & Davey 2015, Davey et al 2019; Davey et al in prep). [Continued on next page]

Stock	Year	Method	Number of fish	Total weight (t)	CV
HPB 1	2011–12	Panel survey	11 754	68.8	0.45
HPB 2	2011–12	Panel survey	8 273	48.4	0.33
HPB 3	2011–12	Panel survey	4 469	26.2	0.33
HPB 5	2011–12	Panel survey	-	-	-
HPB 7	2011–12	Panel survey	1 774	10.4	0.44
HPB 8	2011–12	Panel survey	3 279	19.2	0.61
HPB 1	2017–18	Panel survey	9 857	58.8	0.22
HPB 2	2017–18	Panel survey	7 932	46.6	0.31
HPB 3	2017–18	Panel survey	7 566	44.5	0.4
HPB 5	2017–18	Panel survey	1 185	7.0	0.47
HPB 7	2017–18	Panel survey	4 952	29.1	0.42
HPB 8	2017–18	Panel survey	759	4.5	0.55
HPB 1	2022–23	Panel survey	2 410	16.0	0.37
HPB 2	2022–23	Panel survey	2 340	15.6	0.47

Table 7 [Continued]:

Stock	Year	Method	Number of fish	Total weight (t)	CV
HPB 3	2022–23	Panel survey	3 564	20.2	0.49
HPB 5	2022–23	Panel survey	369	2.5	0.69
HPB 7	2022–23	Panel survey	1 567	10.4	0.4
HPB 8	2022–23	Panel survey	428	2.9	0.72

Table 7a: Recreational harvest estimates for species specific groper stocks for 2022–23 (Heinemann & Gray, in prep). Mean weights from boat ramp survey (Davey et al in prep).

Stock	Year	Method	Number of fish	Total weight (t)	CV
BAS 1	2022–23	Panel survey	427	2.85	0.73
BAS 3	2022–23	Panel survey	708	4.02	0.72
BAS 7	2022–23	Panel survey	448	2.99	0.72
HAP 1	2022–23	Panel survey	1 982	13.20	0.39
HAP 2	2022–23	Panel survey	2 340	15.58	0.47
HAP 3	2022–23	Panel survey	2 856	16.20	0.55
HAP 5	2022–23	Panel survey	369	2.46	0.69
HAP 7	2022–23	Panel survey	1 118	7.44	0.40
HAP 8	2022–23	Panel survey	428	2.85	0.72

1.3 Customary non-commercial fisheries

Groper (hāpuku and bass) were certainly taken by early Maori, and would have been available in greater numbers at shallower depths than is the case at present. Traditional groper grounds are known in several regions. Quantitative information on the current level of customary non-commercial catch is not available.

1.4 Illegal catch

Quantitative information on the level of illegal catch is not available.

1.5 Other sources of mortality

None are apparent.

2. BIOLOGY

Both hāpuku and bass are widely distributed around New Zealand, generally over rough ground from the central shelf (about 100 m) to the shelf edge and down the upper slope. Their lower limits are ill-defined, but hāpuku extends to at least 300 m and bass to 500 m.

Hāpuku mature sexually between 10 and 13 years old and may live in excess of 60 years (Francis et al 1999). Cook Strait hāpuku mature over a wide size range, with the size at 50% maturity at 80–85 cm total length (TL) and 85–90 cm TL for males and females respectively (Paul 2002d). Spawning occurs during winter, anecdotally earlier in the north of New Zealand than in the south, but running ripe fish are seldom caught and spawning grounds are unknown. The smallest juveniles are virtually unknown, but are mottled, pelagic and epi-pelagic, perhaps schooling in association with drifting weed.

The size range of commercially caught hāpuku is 50–140 cm TL, with a broad mode between 70 and 100 cm TL. Bass are slightly larger at 60–150 cm TL, with a mode at 80–110 cm TL, but much bulkier and heavier at equivalent lengths.

There appear to be some regional differences in the size structure of populations. Trawl-caught hāpuku on the Stewart-Snares Shelf are mainly 50–80 cm, modal length 60 cm, and therefore juveniles. Trawl-caught hāpuku on the Chatham Rise are slightly larger, 50–100 cm, modal length 70 cm, with those on the shelf around the islands having their main mode at 60–75 cm; most of these fish are also juveniles. These offshore regions may be important nurseries.

Both groper species are assumed to be long-lived. Natural mortality in the past was assumed to be 0.2, however, a study of a South American (Juan Fernandez) population suggested that it may be lower (0.13–0.16) (Pavez & Oyarzun 1985). Furthermore, preliminary unvalidated ageing in New Zealand has indicated that maximum age may be greater than 40 years, and that M may be 0.1 or less (Francis et al 1999). This value of M will be retained until clearer information becomes available from ageing. Parker et al (2011) compared regional differences in the catch composition from observer collected data. This report noted that the proportion of age 10+ fish in the catch in the Kermadec and Northeastern regions (FMA 2) was greater than that of Southland.

Migration patterns are also little known, but are probably related to spawning. Tagging of mostly immature fish in Cook Strait has shown a high level of site fidelity, but about 5% of these fish have moved up to 160 km north and south. Other information is largely anecdotal and speculative. It is known that good fishing grounds, particularly pinnacles and reefs or ledges, can be quickly fished out and take some time to recover, suggesting a high level of residency (except, perhaps, for during the spawning season). On the other hand, trawlers sometimes catch groper on the flat and clear seafloor, and it is not known whether this represents their normal habitat, whether they are simply dispersing by travelling from one rough ground to another, or whether they are on a purposeful spawning migration.

Hāpuku and bass prey on a wide variety of fish and invertebrates, including red cod, tarakihi, blue cod, hoki and squid. In Cook Strait, they are preyed upon by sperm whales, although probably neither heavily nor selectively. Biological parameters relevant to stock assessment are shown in Table 8.

Table 8: Estimates of biological parameters of groper.

Fishstock	Estimate	Source
<u>1. Natural mortality (M)</u>		
All	$M = 0.1$	Francis et al (1999)
<u>2. Weight = a (length)^b (Weight in g, length in cm fork length)</u>		
	<u>Both sexes combined</u>	
BAS 1	$a = 0.2734$ $b = 2.382$	Johnston (1993)
HAP 1	$a = 0.0142$ $b = 3.003$	Johnston (1993)
HAP 2	$a = 0.0242$ $b = 2.867$	Johnston (1993)
HAP 7, 8	$a = 0.0142$ $b = 2.998$	Johnston (1983)
(HAP = hāpuku, BAS = bass groper)		

3. STOCKS AND AREAS

Tagging studies reveal considerable mixing of hāpuku between Otago, South Canterbury and Cook Strait. Fishstock boundaries in Cook Strait separate Cook Strait hāpuku into three separate "stocks" (HPB 2, HPB 7, and HPB 8), none of which include Otago-Canterbury fish (HPB 3). Current Fishstock boundaries appear inappropriate for the management of Cook Strait and South Island hāpuku. Current stock boundaries are based on QMAs and do not reflect biological stocks. Existing data cannot describe the stock structure of New Zealand groper (Paul 2002b). Electrophoretic studies suggest that separate stocks of hāpuku could occur. However, the genetic heterogeneity of Cook Strait hāpuku, seasonal movements of hāpuku through this area, moderately long-distance movements of some tagged hāpuku, the presence of both species on open ground and the eventual recovery of heavily exploited reefs, suggest that either each stock is moderately mobile or that there is essentially only one stock (of each species) with some small geographic or temporal genetic differences.

4. STOCK ASSESSMENT

Yield estimates for HPB 4 and HPB 5 have been removed because the previous method used is now considered obsolete. The yield estimates for the other Fishstocks have been revised based on a revision of the estimate of M .

4.1 Estimates of fishery parameters and abundance

Estimates of fishery parameters and abundance are not available. Paul (2002c) found that CPUE indices could not be developed for hāpuku and bass either separately or in combination.

4.2 Biomass estimates

Estimates of current and reference biomass are not available. Data for hāpuku from the East Coast South Island trawl surveys have moderate CVs (average over all years = 28.17; range 19–35) and although the survey does not extend to the entire habitat range, the survey may be monitoring settled juveniles (Figure 2).

4.3 Yield estimates and projections

Current biomass cannot be estimated, so *CAY* cannot be determined.




Figure 2: Biomass estimates $\pm 95\%$ CI (estimated from survey CVs assuming a lognormal distribution) and the time series mean (dotted line) from the East Coast South Island trawl survey.

4.4 Other factors

Although no distinct stocks of either groper species have been identified, results from trawl surveys suggest that there are reasonably large but dispersed populations over the Stewart-Snares Shelf and the Chatham Rise. The relationship between these "offshore" and the more traditionally fished "inshore" populations is not known due to the lack of information on groper movements. Little is known of the species composition and population structure of groper on the rough bottom shelf and ridges extending northwards from New Zealand.

The relative quantity of groper taken as target and non-target catch has not been investigated, but is likely to have varied both spatially and temporally. Groper have been taken by the foreign licensed, chartered and New Zealand-owned trawlers working offshore grounds; although being regarded as a small bycatch they were not accurately reported before 1986. The *MCY* may therefore be underestimated.

There are three regions where the groper catch has been substantially lower than the TACC.

HPB 1 - Three features of the fishery appear to explain the under-catch of the TACC. (i) A considerable part of the fishing effort which had generated the high catches in the early 1980s left the fishery. (ii) The allocated quota is widely distributed in small units among fishers who appear to use only a modest proportion of it to cover bycatch. (iii) The fishers who hold larger amounts of quota generally also use only a proportion of it to land high-quality fish (in contrast to the earlier bulk landings of lower-quality fish).

HPB 4 and 5 - The original yield estimates made before the introduction of the QMS and the original TAC were based on trawl surveys, not catch histories. The TACCs for these Fishstocks can only be economically targeted around the Chatham Islands in HPB 4, and a few localities in HPB 5. Elsewhere, it is used to cover a small bycatch from trawlers. A moderate quantity of quota is held, unused, by companies which would require it should they resume target fishing for ling and associated species.

5. STATUS OF THE STOCKS

No estimates of current biomass are available. An estimate of B_{AV} is available for HPB 5.

For all Fishstocks there is insufficient information to estimate current stock status.

6. FOR FURTHER INFORMATION

- Beentjes, M P; Francis, M P (1999) Movements of hapuku, *Polyprion oxygeneios* determined from tagging studies. *New Zealand Journal of Marine and Freshwater Research* 33(1): 1–12.
- Boyd, R O; Reilly, J L (2002) 1999/2000 national marine recreational fishing survey: harvest estimates. Draft New Zealand Fisheries Assessment Report. (Unpublished report held by Fisheries New Zealand.)
- Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16. 27 p. (Unpublished report held by Fisheries New Zealand.)
- Davey, N; Hartill, B; Carter, M (2019) Mean weight estimates for recreational fisheries in 2017–18. *New Zealand Fisheries Assessment Report 2019/25*. 32 p.
- Davey, N K; Johnson, K S; Maggs, J Q (in prep.) Mean weight estimates for recreational fisheries in 2022-23.
- Francis, M P; Mulligan, K P; Davies, N M; Beentjes, M P (1999) Age and growth estimates for New Zealand hapuku, *Polyprion oxygeneios*. *Fishery Bulletin*. 97(2): 227–242.
- Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*.
- Heinemann A; Gray, A. (in prep.) National Panel Survey of Recreational Marine Fishers 2022-23.
- Hurst, R J; Bagley, N W; Uozumi, Y (1990) New Zealand-Japan trawl survey of shelf and upper slope species off southern New Zealand, June 1986. *New Zealand Fisheries Technical Report No. 18*. 50 p.
- Johnston, A D (1983) The southern Cook Strait groper fishery. *Fisheries Technical Report No. 159*. 33 p.
- Johnston, R G (Ed.) (1993) Report from the Conversion Factors Working Group and Steering Committee 1992. MAF Fisheries, Greta Point Internal Report No. 201. 171 p. (Unpublished report held in NIWA library, Wellington.)
- Maggs, J Q; Parsons, D (2023) Design for a catch sampling programme survey to estimate the age structure of New Zealand hapuku (*Polyprion oxygeneios*). *New Zealand Fisheries Assessment Report 2023/34*. 30 p.
- McDougall, C R (1975) Age and growth of *Polyprion oxygeneios* (Pisces: Serranidae) in Cook Strait. (Unpublished B.Sc. (Hons) thesis (Zoology), Victoria University of Wellington.)
- Parker, S J; Paul, L J; Francis, M P (2011) Age structure characteristics of hapuku *Polyprion oxygeneios* stocks from existing samples of otoliths. *New Zealand Fisheries Assessment Report 2011/31*. 42 p.
- Paul, L (2002a) A description of the New Zealand fisheries for the two groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*). *New Zealand Fisheries Assessment Report 2002/13*. 47 p.
- Paul, L (2002b) Can existing data describe the stock structure of the two New Zealand groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*)? *New Zealand Fisheries Assessment Report 2002/14*. 24 p.
- Paul, L (2002c) Can separate CPUE indices be developed for the two groper species, hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*)? *New Zealand Fisheries Assessment Report 2002/15*. 24 p.
- Paul, L (2002d) Size structure of hapuku (*Polyprion oxygeneios*) and bass (*P. americanus*) populations in New Zealand. *New Zealand Fisheries Assessment Report 2002/16*. 17 p.
- Paul, L J (1985) The estimation of hapuku and bass yields for New Zealand fishing regions. Fisheries Research Division Internal Report No. 26. 31 p. (Unpublished report held in NIWA library, Wellington.)
- Paul, L J; Davies, N M (1988) Groper. New Zealand Fisheries Assessment Research Document 88/15. 27 p. (Unpublished report held by NIWA library, Wellington.)
- Pavez, P; Oyarzun, M E (1985) Determination of the relative efficiency of hooks, and growth parameters of the Juan Fernandez "cod" *Polyprion oxygeneios* Bloch and Schneider, 1801, in the Robinson Crusoe and Santa Clara Islands. In Arana, P (Ed.), "Investigaciones en el Archipelago de Juan Fernandez", pp. 341–345. Escuela de Ciencias del Mar, UCV, Valparaiso. [In Spanish, English summary.]
- Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991/92 to 1993/94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)
- Teirney, L; McKinnon, S; Kilner, A; Sylvester, T (1991) Marine Recreational Fisheries Working Group Report — November 1991. New Zealand Fisheries Working Group Report 91/1. 46 p.
- Teirney, L D; Olsen, D L (1992) Marine Recreational Fisheries Group Report — November 1992. New Zealand Fisheries Working Group Report 92/1. 13 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*.
- Wynne-Jones, J; Gray, A; Heinemann, A; Hill, L; Walton, L (2019) National Panel Survey of Marine Recreational Fishers 2017–2018. *New Zealand Fisheries Assessment Report 2019/24*. 104 p.