
RED GURNARD (GUR)

(*Chelidonichthys kumu*)
Kumukumu





1. FISHERY SUMMARY

1.1 Commercial fisheries

Red gurnard are a major bycatch of inshore trawl fisheries in most areas of New Zealand, including fisheries for red cod in the southern regions and flatfish off the west coast of the South Island (WCSI) and in Tasman Bay. They are also directly targeted in some areas, e.g., GUR 2. Some minor target fisheries for red gurnard are known in Pegasus Bay, off Mahia, and off the west coast South Island. Red gurnard is also a minor bycatch in the jack mackerel trawl fishery in the South Taranaki Bight. Up to 15% of the total red gurnard catch is taken by bottom longline and set net.

Red gurnard was introduced into the Quota Management System (QMS) in 1986. The 1986 TACCs were based on 1984 landings for Southland and 1983 landings for other regions. TACCs for GUR 2, GUR 8, and GUR 10 have remained unchanged since, and these fisheries remain TACC only fisheries. TACs and allowances have since been set for GUR 7, GUR 3, and GUR 1 (1997, 2000, and 2021, respectively), with GUR 3 and GUR 7 having numerous changes to TACC and allowances since their introduction to the QMS, discussed below. All current TACs, allowances, and TACCs are given in Table 1. Under the Adaptive Management Programme (AMP), which ended 30 September 2009, the TACCs for GUR 3 and 7 were increased to 600 t and 815 t, respectively, for the 1991–92 fishing year, and then the GUR 7 TACC was reduced to 678 t, in 1997–98. The TACC for GUR 3 was increased to 900 t for the 1996–97 fishing year under the AMP but was decreased to 800 t in 2002–03.

For the 2009–10 fishing season, the TACC in GUR 7 was increased to 715 t, including an allocation of 10 t for customary, 20 t for recreational use, and 14 t allocation for other sources of mortality. The GUR 7 TACC was further increased to 785 t in October 2012, 845 t in October 2015, 975 t in October 2017, 1073 t in October 2019, 1180 t in October 2020, 1298 in October 2021, and 1450 t in 2022–23 along with increased allowances. For the 2009–10 fishing season, the TACC for GUR 3 was increased from 800 t to 900 t, with allocations of 3 t, 5 t, and 45 t for customary, recreational, and other sources of mortality, respectively. The GUR 3 TACC was further increased to 1100 t in October 2012, 1220 t in October 2015, 1320 t in October 2018, 1500 t in 2020, and 1575 in 2022 along with increases in recreational and other sources of mortality allowances (now 6 t and 105 t, respectively). In October 2021, when a new TAC was set, the TACC for GUR 1 was decreased from 2288 t to 800 t and

allocations of 40 t for customary, 100 t for recreational, and 56 t for other sources of mortality were also set.

Reported landings since 1931 are shown in Tables 2 and 3. Historical landings and TACC values for the five main GUR stocks are shown in Figure 1.

Table 1: Current TACs, TACCs, and allowances (t) for red gurnard by Fishstock as of October 2023.

Fishstock	TAC	TACC	Customary allowance	Recreational allowance	Other mortality
GUR 1	996	800	40	100	56
GUR 2		725			
GUR 3	1 614	1 575	3	6	105
GUR 7	1 422	1 450	17	42	65
GUR 8		543			
GUR 10		10			

Table 2: Reported landings (t) for the main QMAs from 1931 to 1982.

Year	GUR 1	GUR 2	GUR 3	GUR 7	GUR 8
1931–32	67	0	1	16	0
1932–33	42	0	0	13	0
1933–34	67	84	1	20	0
1934–35	50	179	0	2	0
1935–36	75	147	18	2	0
1936–37	114	215	37	25	1
1937–38	205	193	83	21	0
1938–39	109	118	151	31	2
1939–40	121	149	147	25	1
1940–41	124	222	215	38	1
1941–42	107	200	267	38	0
1942–43	124	332	287	58	0
1943–44	128	244	294	53	0
1944	238	292	291	60	0
1945	360	338	222	94	3
1946	426	387	290	119	4
1947	376	297	243	162	10
1948	385	243	267	226	9
1949	371	264	316	323	13
1950	306	186	486	332	13
1951	221	231	750	202	10
1952	394	378	658	211	5
1953	490	494	614	334	3
1954	496	462	660	382	7
1955	495	283	652	490	25
1956	434	312	782	435	29
1957	494	402	737	409	46
1958	430	394	745	400	51
1959	460	320	806	212	44
1960	489	417	1 008	421	27
1961	559	419	1 180	419	27
1962	505	592	1 244	322	14
1963	576	562	1 364	367	8
1964	977	814	1 708	397	16
1965	1 020	668	1 459	400	34
1966	1 157	754	1 178	436	27
1967	1 051	836	745	522	45
1968	1 137	583	510	368	52
1969	1 345	632	487	256	33
1970	1 493	823	841	381	53
1971	1 225	570	940	379	37
1972	770	347	662	333	15
1973	1 278	406	1 393	491	21
1974	881	299	1 083	586	41
1975	691	199	655	365	28
1976	1 055	217	960	545	52
1977	1 288	381	975	579	45
1978	1 571	519	1 106	487	26
1979	1 936	382	690	349	18
1980	1 845	438	672	253	34
1981	2 349	603	438	318	16
1982	2 084	454	379	368	34

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. Data up to 1985 are from fishing returns; data from 1986 to 1990 are from Quota Management Reports.
3. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data include both foreign and domestic landings. Data were aggregated to FMA using methods and assumptions described by Francis & Paul (2013).

Table 3: Reported landings (t) of red gurnard by Fishstock from 1983–84 to the present and actual TACCs (t) from 1986–87 to the present. The QMS data are from 1986 to the present. [Continued on next page]

Fishstock	GUR 1		GUR 2		GUR 3		GUR 7	
QMA (s)	1 & 9		2		3, 4, 5 & 6		7	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	2 099	–	782	–	366	–	468	–
1984–85*	1 531	–	665	–	272	–	332	–
1985–86*	1 760	–	495	–	272	–	239	–
1986–87	1 021	2 010	592	610	210	480	421	610
1987–88	1 139	2 081	596	657	386	486	806	629
1988–89	1 039	2 198	536	698	528	489	479	669
1989–90	916	2 283	451	720	694	501	511	678
1990–91	1 123	2 284	490	723	661	524	442	678
1991–92	1 294	2 284	663	723	539	600	704	815
1992–93	1 629	2 284	618	725	484	601	761	815
1993–94	1 153	2 284	635	725	711	601	469	815
1994–95	1 054	2 287	559	725	685	601	455	815
1995–96	1 163	2 287	567	725	633	601	382	815
1996–97	1 055	2 287	503	725	641	900	378	815
1997–98	1 015	2 287	482	725	477	900	309	678
1998–99	927	2 287	469	725	395	900	323	678
1999–00	944	2 287	521	725	411	900	331	678
2000–01	1 294	2 287	623	725	569	900	571	678
2001–02	1 109	2 287	619	725	717	900	686	681
2002–03	1 256	2 287	552	725	888	800	793	681
2003–04	1 225	2 287	512	725	725	800	717	681
2004–05	1 354	2 287	708	725	854	800	688	681
2005–06	1 113	2 287	542	725	957	800	604	681
2006–07	1 180	2 287	575	725	1 004	800	714	681
2007–08	1 198	2 287	517	725	842	800	563	681
2008–09	1 060	2 287	621	725	939	800	595	681
2009–10	1 075	2 287	853	725	1 018	900	603	715
2010–11	1 046	2 288	587	725	929	900	545	715
2011–12	981	2 288	558	725	915	900	684	715
2012–13	1 103	2 288	603	725	1 168	1 100	763	785
2013–14	1 005	2 288	555	725	1 223	1 100	837	785
2014–15	1 020	2 288	695	725	1 150	1 100	852	785
2015–16	860	2 288	748	725	1 348	1 220	852	845
2016–17	856	2 288	669	725	1 279	1 220	905	845
2017–18	785	2 288	560	725	1 419	1 220	882	975
2018–19	710	2 288	587	725	1 467	1 320	998	975
2019–20	745	2 288	562	725	1 537	1 320	1 182	1 073
2020–21	847	2 288	412	725	1 646	1 500	1 153	1 180
2021–22	700	800	405	725	1 572	1 500	1 005	1 298
2022–23	615	800	338	725	1 546	1 575	774	1 450

Fishstock	GUR 8		GUR 10			
OMA (s)	8		10			Total
	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	251	–	0	–	3 966	–
1984–85*	247	–	0	–	3 047	–
1985–86*	163	–	0	–	2 929	–
1986–87	159	510	0	10	2 403	4 230
1987–88	194	518	0	10	3 121	4 381
1988–89	167	532	0	10	2 749	4 596
1989–90	173	538	0	10	2 745	4 730
1990–91	150	543	0	10	2 866	4 762
1991–92	189	543	0	10	3 390	4 975
1992–93	208	543	0	10	3 700	4 978
1993–94	174	543	0	10	3 142	4 978
1994–95	217	543	0	10	2 969	4 982
1995–96	182	543	0	10	2 927	4 982
1996–97	219	543	0	10	2 796	5 281
1997–98	249	543	0	10	2 532	5 143
1998–99	170	543	0	10	2 284	5 143
1999–00	222	543	0	10	2 429	5 143
2000–01	291	543	0	10	3 348	5 143
2001–02	302	543	0	10	3 429	5 143
2002–03	342	543	0	10	3 831	4 993
2003–04	329	543	0	10	3 508	4 993
2004–05	370	543	0	10	3 974	4 993
2005–06	373	543	0	10	3 589	4 993
2006–07	349	543	0	10	3 822	4 993
2007–08	223	543	0	10	3 344	4 993
2008–09	274	543	0	10	3 489	4 993
2009–10	239	543	0	10	3 789	5 181
2010–11	182	543	0	10	3 289	5 181
2011–12	213	543	0	10	3 351	5 181
2012–13	170	543	0	10	3 807	5 451
2013–14	151	543	0	10	3 769	5 451
2014–15	193	543	0	10	3 910	5 451
2015–16	145	543	0	10	3 953	5 631
2016–17	145	543	0	10	3 854	5 631
2017–18	209	543	0	10	3 855	5 761

Table 3 [Continued]:

Fishstock		GUR 8		GUR 10		
QMA (s)		8		10		Total
	Landings	TACC	Landings	TACC	Landings	TACC
2018–19	267	543	0	10	4 029	5 861
2019–20	386	543	0	10	4 412	5 959
2020–21	249	543	0	10	4 306	6 246
2021–22	221	543	0	10	3 903	4 876
2022–23	86	543	< 1	10	3 359	5 104

*FSU data.

Annual landings of GUR 1 were relatively stable from 1986–87 to 2014–15, generally ranging between 920 t and 1300 t; substantially lower than the 2288 t TACC. Since then, catches have declined slightly, with 745 t landed in 2019–20. About 60% of the GUR 1 total is taken from FMA 1, as a bycatch of a number of fisheries including inshore trawl fisheries for snapper, John dory, and tarakihi. The remaining 40% is taken from FMA 9, mainly as a bycatch of the snapper and trevally inshore trawl fisheries.

GUR 2 landings fluctuated within the range of 469–853 t from 1991–92 to 2019–20, typically well below the TACC. In addition to the target fishery, red gurnard are taken as a bycatch of the tarakihi, trevally, and snapper inshore trawl fisheries. A decreasing trend in effort in GUR 2 is evident from 2009–10 to 2022–23 and GUR 2 landings in 2022–23 of 338 t were the lowest since the mid 1970s.

GUR 3 landings regularly exceeded the TACC between 1988–89 and 1995–96 and this stock has been consistently over-caught since 2004–05. Landings increased steadily from 2010–11 to 2020–21 with multiple small increases in the TACC during that period.




Figure 1: Reported commercial landings and TACCs for the five main GUR stocks. From top to bottom: GUR 1 (Auckland East) and GUR 2 (Central East). [Continued on next page]




Figure 1 [Continued]: Reported commercial landings and TACCs for the five main GUR stocks. From top to bottom: GUR 3 (South East Coast), GUR 7 (Challenger), and GUR 8 (Central Egmont).

GUR 7 landings declined steadily from 761 t in 1992–93, to 309 t in 1997–98, but then increased to 793 t by 2002–03. Landings then generally declined to 2010–11. Landings increased steadily from 2010–11 to 2020–21, but then declined, with multiple small increases in the TACC during that period.

Landings in GUR 8 have remained well below the TACC since 1986–87, averaging about 225 t.

1.2 Recreational fisheries

Red gurnard is, by virtue of its wide distribution in harbours and shallow coastal waters, an important recreational species. It is often taken by fishers targeting snapper and tarakihi, particularly around the

North Island. The allowances within the TAC for each Fishstock are shown in Table 1, but they are currently only set for GUR 1, GUR 3, and GUR 7.

1.2.1 Management controls

The main methods used to manage recreational harvests of red gurnard are minimum legal size limits (MLS), method restrictions, and daily bag limits. Fishers can take up to 20 GUR as part of their combined daily bag limit and the MLS is 25 cm.

1.2.2 Estimates of recreational harvest

Recreational catch estimates are given in Table 4. There are two broad approaches to estimating recreational fisheries harvest: the use of onsite or access point methods where fishers are surveyed or counted at the point of fishing or access to their fishing activity; and offsite methods where some form of post-event interview and/or diary are used to collect data from fishers.

The first estimates of recreational harvest for red gurnard were calculated using an offsite approach: the offsite regional telephone and diary survey approach. Estimates for 1996 came from a national telephone and diary survey (Bradford 1998). Another national telephone and diary survey was carried out in 2000 (Boyd & Reilly 2004) and a rolling replacement of diarists in 2001 (Boyd et al 2004) allowed estimates for a further year (population scaling ratios and mean weights were not re-estimated in 2001).

Table 4: Recreational harvest estimates for red gurnard stocks. The telephone/diary surveys and earlier aerial-access surveys ran from December to November but are denoted by the January calendar year. The surveys since 2010 have run through the October to September fishing year but are denoted by the January calendar year. Mean fish weights were obtained from boat ramp surveys (for the telephone/diary and panel survey harvest estimates, Hartill & Davey 2015, and Davey et al 2019).

Stock	Year	Method	Number of fish	Total weight (t)	CV
GUR 1	1996	Telephone/diary	262 000	108	0.07
	2000	Telephone/diary	465000	223	0.16
FMA 1 only	2005	Aerial–access	–	127	0.14
FMA 1 only	2012	Aerial–access	–	24	0.09
FMA 1 only	2012	Panel survey	120 500	49	0.16
	2012	Panel survey	241 957	103	0.15
FMA 1 only	2018	Aerial–access	–	31	0.11
FMA 1 only	2018	Panel survey	85 000	36	0.14
	2018	Panel survey	168 798	86	0.15
	2023	Panel survey	59 705	30	0.14
GUR 2	1996	Telephone/diary	38 000	16	0.18
	2000	Telephone/diary	209 000	127	0.37
	2012	Panel survey	66 661	38	0.20
	2018	Panel survey	71 702	39	0.28
	2023	Panel survey	85 233	54	0.63
GUR 3	1996	Telephone/diary	1 000	–	–
	2000	Telephone/diary	11 000	5	0.70
	2012	Panel survey	4 605	2	0.62
	2018	Panel survey	3 486	2	0.39
	2023	Panel survey	3 475	2	0.34
GUR 7	1996	Telephone/diary	26 000	12	0.15
	2000	Telephone/diary	36 000	11	0.23
	2012	Panel survey	23 653	12	0.24
	2018	Panel survey	60 759	38	0.18
	2023	Panel survey	27 882	15	0.19
GUR 8	1996	Telephone/diary	67 000	28	0.15
	2000	Telephone/diary	99 000	40	0.36
	2012	Panel survey	93 656	47	0.23
	2018	Panel survey	55 314	31	0.19
	2023	Panel survey	24 805	14	0.21

The harvest estimates provided by these telephone diary surveys are no longer considered reliable for various reasons. With the early telephone/diary method, fishers were recruited to fill in diaries by way of a telephone survey that also estimates the proportion of the population that is eligible (likely to fish). A ‘soft refusal’ bias in the eligibility proportion arises if interviewees who do not wish to co-operate falsely state that they never fish. The proportion of eligible fishers in the population (and, hence, the harvest) is thereby under-estimated. Pilot studies for the 2000 telephone/diary survey suggested that this effect could occur when recreational fishing was established as the subject of the interview at the outset. Another equally serious cause of bias in telephone/diary surveys was that diarists who did not immediately record their day’s catch after a trip sometimes overstated their catch or the number of trips

made. There is some indirect evidence that this may have occurred in all the telephone/diary surveys (Wright et al 2004).

The recreational harvest estimates provided by the 2000 and 2001 telephone diary surveys are thought to be implausibly high for many species, which led to the development of an alternative maximum count aerial-access onsite method that provides a more direct means of estimating recreational harvests for suitable fisheries. The maximum count aerial-access approach combines data collected concurrently from two sources: a creel survey of recreational fishers returning to a subsample of ramps throughout the day; and an aerial survey count of vessels observed to be fishing at the approximate time of peak fishing effort on the same day. The ratio of the aerial count in a particular area to the number of interviewed parties who claimed to have fished in that area at the time of the overflight was used to scale up harvests observed at surveyed ramps, to estimate harvest taken by all fishers returning to all ramps. The methodology is further described by Hartill et al (2007).

This aerial-access method was first employed and optimised to estimate snapper harvests in the Hauraki Gulf in 2003–04. It was then extended to survey the wider SNA 1 fishery in 2004–05 and to provide estimates for other species, including red gurnard (FMA 1 only for GUR). In response to the cost and scale challenges associated with onsite methods, in particular the difficulties in sampling other than trailer boat fisheries, offsite approaches to estimating recreational fisheries harvest have been revisited. This led to the development and implementation of a national panel survey for the 2011–12 fishing year (Wynne-Jones et al 2014) and repeated for the 2017–18 and 2022–23 fishing years (Wynne-Jones et al 2019; Heinemann & Gray, in prep). The panel survey used face-to-face interviews of a random sample of New Zealand households to recruit a panel of fishers and non-fishers for a full year. The panel members were contacted regularly about their fishing activities and catch information in standardised phone interviews.

1.3 Customary non-commercial fisheries

Red gurnard is an important species for customary non-commercial fishing interests, by virtue of its wide distribution in shallow coastal waters. However, no quantitative estimates of customary non-commercial catch are currently available.

1.4 Illegal catch

No quantitative information is available.

1.5 Other sources of mortality

No quantitative information is available**.**

2. BIOLOGY

Gurnard growth rate varies with location, and females grow faster and are usually larger at age than males. Maximum age (A_{MAX}) is about 16 years and maximum size is 55+ cm. Red gurnard reach sexual maturity at an age of 2–3 years and a fork length (FL) of about 23 cm, after which the growth rate slows. An analysis of the age and growth of red gurnard in FMA 7 revealed that young fish 1–4 years old tend to be most common in Tasman Bay and Golden Bay. Three to six year old fish are found on the inshore areas off the west coast South Island and the older fish are predominantly found further offshore (Lyon & Horn 2011).

Biological parameters relevant to the stock assessment are shown in Table 5.

M was estimated using the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock. Samples from the east coast South Island (ECSI) suggested an A_{MAX} of about 16 years for males and 13 years for females, giving estimates for M of 0.29 and 0.35, respectively. Samples from the west coast South Island (WCSI) indicate an A_{MAX} of about 15 years for both sexes, giving an estimate of 0.31 for M. These samples were not from virgin populations, so M may be overestimated.

Table 5: Estimates of biological parameters for red gurnard.

Fishstock	Estimate					Source
1. Natural mortality (*M*)						
	Female			Males		
GUR 1W & 1E	0.30			0.35		Stevenson (2000)
GUR 3	0.29			0.35		Sutton (1997)
GUR 7	0.31			0.31		Sutton (1997)
2. Weight = a(length)^b (Weight in g, length in cm fork length)						
				Both Sexes		
	a			b		
GUR 1	0.00998			2.99		Elder (1976)
GUR 1W & 1 E	0.026			2.775		Stevenson (2000)
GUR 2	0.0053			3.19		Stevenson (2000)
3. von Bertalanffy growth parameters						
	Females			Males		
	L_∞	k	t_0	L_∞	k	t_0	
GUR 1	36.4	0.641	0.189	28.8	0.569	-0.552	Elder (1976)
GUR 1W	45.3	0.25	-0.88	36.5	0.45	-0.30	Stevenson (2000)
GUR 1E	44.5	0.28	-0.76	35.2	0.49	-0.24	Stevenson (2000)
GUR 3	48.2	0.44	0.1	42.2	0.49	-0.26	Sutton (1997)
GUR 7	45.7	0.40	-0.36	40.3	0.37	-0.96	Sutton (1997)

Red gurnard have a long spawning period which extends through spring and summer with a peak in early summer. In the Hauraki Gulf, ripe adults can be found throughout the year. Spawning grounds appear to be widespread, although perhaps localised over the inner and central shelf. Egg and larval development takes place in surface waters, and there is a period of at least eight days before feeding starts. Small juveniles (under 15 cm FL) are often caught in shallow harbours, but rarely in commercial trawls.

3. STOCKS AND AREAS

There are no data that would alter the current stock boundaries. No information is available on stock separation of red gurnard. For GUR 3 the Working Group noted that spatial information from the CPUE analyses indicated that separate stocks or sub-stocks may exist between the east and south coasts of the South Island.

4. STOCK ASSESSMENT

4.1 Biomass estimates

Relative abundance indices have been obtained from trawl surveys of the Bay of Plenty, west coast North Island, and Hauraki Gulf within the GUR 1 fish stock (Figure 2, Table 6); east coast South Island (GUR 3); and west coast South Island and Tasman Bay/Golden Bay combined (GUR 7) (Table 7).

East coast South Island (ECSI) inshore trawl survey

The ECSI winter surveys from 1991 to 1996 in 30–400 m were replaced by summer trawl surveys (1996–97 to 2000–01) which also included the 10–30 m depth range, but these were discontinued after the fifth in the annual time series because of the extreme fluctuations in catchability between surveys (Francis et al 2001). The winter surveys were reinstated in 2007 and this time included additional 10–30 m strata in an attempt to index elephantfish and red gurnard which were officially included in the list of target species in 2012. The 2007 survey and all surveys from 2012 provide full coverage of the 10–30 m depth range.

Red gurnard biomass in the 1990s averaged 422 t in the core strata, increasing more than three-fold in 2007, after which it has generally been increasing with the exception of 2016, when biomass more than halved (Beentjes et al 2023). The 2021 and 2022 biomass estimates were the two highest in the time series, with the 2022 estimate 39% higher than in 2021 (Figure 3, Table 7). Total biomass trends in the seven core plus shallow strata surveys (2007 and 2012 onward) were similar to those in the core strata with the same trend of increasing biomass from 2007 to 2022, except that biomass was higher (Figure 3, Table 7). The proportion of pre-recruit biomass in the core strata varied greatly among surveys, from 2 to 20%, and in 2022 it was 3% (Beentjes et al 2023). The proportion of juvenile biomass (based on the length-at-50% maturity) within the core strata was close to zero for all surveys including 2022 when it was 0.1%.

The proportion of the red gurnard biomass in the 10–30 m depth range has varied from 29% to 61% of the total biomass, indicating the importance of the inshore strata for this species.

The red gurnard time series of length frequency distributions in the seven core plus shallow strata (10–400 m) are reasonably consistent over time with clear adult modes, but variable 0+ and 1+ modes. The addition of the 10–30 m depth range had no significant effect on the length frequency distributions in some years (2007, 2014, and 2018), but in 2012 and 2016 there were abundant 1+ cohorts in 10–30 m that were poorly represented in the core strata, and a 0+ cohort (13–18 cm) is apparent in the 10–30 m depth range in 2021 and 2022 (Beentjes et al 2023). Based on the seven surveys that included the 10–30 m strata, there are generally more pre-recruit fish in the shallow strata, suggesting that the core plus shallow strata (10 to 400 m) survey is probably indexing red gurnard abundance, including juveniles. The spatial distribution of red gurnard aggregations varies over the time series, but overall this species is consistently well represented from about 10 to 75 m in the core plus shallow strata, along the entire survey coastline with many of the largest survey catches in less than 30 m They are almost absent deeper than 100 m. The inclusion of the 10–30 m depth range has thus been valuable for monitoring red gurnard.




Figure 2: Estimates of recruited (length ≥ 30 cm) red gurnard biomass (t) from *Kaharoa* trawl surveys within GUR 1. Error bars are ± two standard deviations.

Table 6: Estimates of red gurnard recruited biomass (t) from *Kaharoa* trawl surveys within GUR 1. Red gurnard is assumed to recruit at 30 cm TL. For the west coast North Island trawl survey, core strata are north of New Plymouth.

Survey area	Year	Trip code	Biomass (t)	CV(%)
Bay of Plenty	1983	KAH8303	229	31
	1985	KAH8506	27	17
	1990	KAH9004	234	11
	1992	KAH9202	166	10
	1996	KAH9601	185	16
	1999	KAH9902	180	17
	2020	KAH2001	130	14
	2021	KAH2101	85	10
Hauraki Gulf	1984	KAH8421	208	21
	1985	KAH8517	33	57
	1986	KAH8613	124	32
	1988	KAH8810	329	25
	1989	KAH8917	58	39
	1990	KAH9016	64	17
	1992	KAH9212	169	7
	1993	KAH9311	85	18
	1994	KAH9411	98	21
	1997	KAH9720	107	22
	2000	KAH0012	15	52
	2019	KAH1907	120	34
	2020	KAH2006	96	12
West coast North Island	1989	KAH8918	758	13
	1991	KAH9111	1435	18
	1994	KAH9410	1662	28
	1996	KAH9615	1090	11
	1999	KAH9915	610	13
	2018	KAH1806	551	14
	2019	KAH1906	758	13
	2020	KAH2005	824	13




Figure 3: Red gurnard total biomass for all ECSI winter surveys in core strata (30–400 m), and core plus shallow strata (10–400 m) in 2007, 2012, 2014, 2016, 2018, 2021 and 2022. Error bars are ± two standard deviations.

Table 7: Relative biomass indices (t) and coefficients of variation (CV) for red gurnard around the North Island and South Island*. Biomass estimates for ECSI in 1991 were adjusted to allow for non-sampled strata (7 & 9 equivalent to current strata 13, 16, and 17). The sum of pre-recruit and recruited biomass values do not always match the total biomass for the earlier surveys because at several stations length frequencies were not measured, affecting the biomass calculations for length intervals. – , not measured; NA, not applicable. Recruited is defined as the size-at-recruitment to the fishery (30 cm). Biomass estimates from current surveys with extreme catchability are denoted with a #.

Region	Fishstock	Year	Trip number	Total biomass estimate	CV (%)	Total biomass estimate	CV (%)	Pre-recruit	CV (%)	Pre-recruit	CV (%)	Recruited	CV (%)	Recruited	CV (%)
WCSI	GUR 7	1992	KAH9204	572	15	–	–	–	–	–	–	–	–	454.0	15.4
		1994	KAH9404	559	15	–	–	–	–	–	–	–	–	478.3	16.0
		1995	KAH9504	584	19	–	–	–	–	–	–	–	–	501.6	21.7
		1997	KAH9704	471	13	–	–	–	–	–	–	–	–	309.8	14.5
		2000	KAH0004	625	15	–	–	–	–	–	–	–	–	444.0	14.9
		2003	KAH0304	#270	20	–	–	–	–	–	–	–	–	253.7	20.9
		2005	KAH0503	442	17	–	–	–	–	–	–	–	–	374.7	16.2
		2007	KAH0704	553	17	–	–	–	–	–	–	–	–	431.6	17.9
		2009	KAH0904	651	18	–	–	–	–	–	–	–	–	400.4	19.1
		2011	KAH1104	1 070	17	–	–	–	–	–	–	–	–	798.6	18.6
		2013	KAH1305	754	12	–	–	–	–	–	–	–	–	546.5	13.4
		2015	KAH1503	1 774	16	–	–	–	–	–	–	–	–	1 335.2	18.6
		2017	KAH1703	1 708	12	–	–	–	–	–	–	–	–	1 352.0	12.0
		2019	KAH1902	1 642	16	–	–	–	–	–	–	–	–	1 079.0	16.0
		2021	KAH2103	2 022	18	–	–	–	–	–	–	–	–	1 663	18
		2023	KAH2302	1 498	15	–	–	–	–	–	–	–	–	1 057	15
North Island east coast	GUR 2	1993	KAH9304	439	44	–	–	–	–	–	–	–	–	–	–
		1994	KAH9402	871	16	–	–	–	–	–	–	–	–	–	–
		1995	KAH9502	178	26	–	–	–	–	–	–	–	–	–	–
		1996	KAH9602	708	29	–	–	–	–	–	–	–	–	–	–
ECSI (winter)	GUR 3			30–400 m		10–400 m		30–400 m		10–400 m		30–400 m		10–400 m	
		1991	KAH9105	763	33	–	–	NA	NA	–	–	NA	NA	–	–
		1992	KAH9205	142	30	–	–	21	58	–	–	121	30	–	–
		1993	KAH9306	576	31	–	–	26	45	–	–	551	31	–	–
		1994	KAH9406	123	34	–	–	2	42	–	–	121	34	–	–
		1996	KAH9606	505	27	–	–	8	44	–	–	496	26	–	–
		2007	KAH0705	1 453	35	2 048	27	298	40	494	32	1 155	35	1 554	27
		2008	KAH0806	1 309	34	–	–	100	59	–	–	1 210	33	–	–
		2009	KAH0905	1 725	30	–	–	62	34	–	–	1 663	30	–	–
		2012	KAH1207	1 680	28	3 515	17	193	40	742	31	1 487	27	2 773	16
		2014	KAH1402	2 063	25	3 215	17	409	45	585	32	1 654	23	2 630	16
		2016	KAH1605	941	30	2 420	15	63	41	306	19	877	30	2 114	15
		2018	KAH1803	2 043	19	3 831	17	308	24	610	21	1 735	20	3 221	18
		2021	KAH2104	2 068	32	3 724	19	215	50	422	28	1 854	14	3 302	18
		2022	KAH2204	3 410	30	5 471	20	110	49	273	32	3 299	30	5 199	20
ECSI (summer)	GUR 3	1996–97	KAH9618	765	13	–		–	–	–	–	–	–	–	–
		1997–98	KAH9704	317	16	–		–	–		–	–	–	–	–
		1998–99	KAH9809	493	13	–	–	–	–	–	–	–	–	–	–
		1999–00	KAH9917	202	20	–	–	–	–	–	–	–	–	–	–
		2000–01	KAH0014	146	34	–	–	–	–	–	–	–	–	–	–

*Assuming areal availability, vertical availability, and vulnerability equal 1.0. Biomass is only estimated outside 10 m. Note: because trawl survey biomass estimates are indices, comparisons between different seasons (e.g., summer and winter ECSI) are not strictly valid.

West coast South Island (WCSI) inshore trawl survey

Red gurnard biomass estimates were consistent from 1992 to 2000 but had declined by the 2003 survey to the time series low of 270 t (Table 7, Figure 4). Biomass increased from 2003 and, since 2015, has been at record highs, with 2021 being the highest estimate in the time series (2022 t). The 2023 estimate dropped by 26% but is still well above the time series mean (950 t). A large proportion of the biomass had always occurred in the Tasman Bay and Golden Bay region, but markedly more has been from the west coast South Island in most years since 2011 (MacGibbon et al 2024). Both regions saw a decline in biomass in 2023.

Throughout the time series, larger numbers of smaller fish have been in Tasman Bay and Golden Bay compared with the west coast (MacGibbon et al 2024). Conversely, the west coast had larger numbers of larger fish than Tasman Bay and Golden Bay. However, in 2023, larger numbers of smaller fish were seen off the west coast than in Tasman Bay and Golden Bay, including a strong 1+ cohort of about 13–18 cm. Although 1+ cohorts have been visible in a number of surveys (e.g., 2009, 2013, 2015, and 2021), the 1+ cohort in 2023 was the strongest in the time series (MacGibbon et al 2024).

Throughout the time series, recruit/adult (over 30 cm) biomass has been greater than pre-recruit/juvenile biomass, although juvenile biomass was often substantial (Figure 5). Although overall biomass declined in 2023 compared with 2021, this decrease came entirely from recruits/adults; pre-recruit/juvenile biomass had, in fact, increased slightly.




Figure 4: Red gurnard biomass trends from the west coast South Island inshore trawl survey time series. Error bars are ± two standard deviations.




Figure 5: Red gurnard pre-recruit (< 30 cm) and recruited biomass trends from the west coast South Island inshore trawl survey time series. Error bars are ± two standard deviations.

4.2 CPUE Analyses

GUR 1

In 2022, McKenzie & Moore (2023) updated CPUE analyses for GUR 1W (west coast, Figure 6), GUR 1E (east Northland and Hauraki Gulf, Figure 7), and GUR 1BP (Bay of Plenty, Figure 8).

The analyses were based on catch and effort data for individual tows reported on TCEPR, TCER, and ERS forms because adequate time series are available in the northern inshore trawl fisheries from 1995–96. Based on catch and effort data from single bottom trawls targeting gurnard, snapper, trevally, tarakihi, barracouta, or John dory, two GLM models were produced for each subarea: one based on the magnitude of positive catch (lognormal distribution), and the other a binomial model of the probability of capture (based on the proportion of tows capturing GUR). The two models were then combined to produce a single series for each sub-area, and the Working Group accepted the combined models as indices of abundance. The data used to generate the GLM models were restricted to core fleets of vessels having had at least three trips in each of three years.




Figure 6: Standardised probability of catch (binomial model), positive CPUE indices (lognormal model), and combined model for GUR 1W using bottom trawl tow data from TCEPR/ TCER/ERS forms (McKenzie & Moore 2023). Error bars are 95% confidence intervals.




Figure 7: Standardised probability of catch (binomial model), positive catch CPUE indices (lognormal model), and combined model for GUR 1E using bottom trawl tow data from TCEPR/ TCER/ERS forms (McKenzie & Moore 2023). Error bars are 95% confidence intervals.




Figure 8: Standardised probability of catch (binomial model), positive catch CPUE indices (gamma model), and combined model for GUR 1BP using bottom trawl tow data from TCEPR/ TCER forms (McKenzie & Moore 2023). Error bars are 95% confidence intervals.

All three series show strong cyclical fluctuations with a recovery from low levels between 1996 and 1999 to a peak in the early 2000s, followed by a subsequent decline to low levels again between 2009 and 2013, then an increase to levels near, or above the long-term average between 2015 and 2017. In subsequent years the levels have declined (GUR 1W, GUR 1E), or declined then increased again (GUR 1BP). Despite overall similarities, the series differ somewhat with respect to the magnitude of the fluctuations and the specific years for the nadir and the peak.

Establishing B_{MSY}-compatible reference points for GUR 1

In 2013, the Working Group accepted mean standardised bottom trawl CPUE for the period 1995–96 to 2011–12 as B_{MSY}-compatible proxies for each of the GUR 1 sub-stocks. All three series were based on combined positive catch and probability of capture models derived from event-scale fishing events (i.e., tow). GUR abundance tends to fluctuate in cycles, according to recruitment, and the period was chosen because it included at least one cycle of abundance and high catch. The Working Group accepted the default Harvest Strategy Standard definitions that the Soft Limit and Hard Limit would be one half and one quarter the target for each sub-stock, respectively.

Future Research Considerations

- Integrate the results of the reinstated west coast North Island (WCNI) trawl survey and CPUE series into the stock assessment.
- Integrate the results of the reinstated Hauraki Gulf trawl survey and CPUE series into the stock assessment.
- Integrate the results of the reinstated Bay of Plenty trawl survey and CPUE series into the stock assessment.
- Age otoliths from recent surveys for the stock assessments.

GUR 2

GUR 2 is monitored using standardised CPUE from the bottom trawl fishery targeting gurnard, snapper, or trevally.

In 2017, Schofield et al (2018a) updated CPUE analyses for GUR 2. Landings were allocated to daily aggregated effort using methods described by Langley (2014) to improve the consistency of the data collected from the different statutory reporting forms (CELR and TCER). A core fleet of vessels that had completed at least five trips per year in at least seven years was modelled using a Weibull distribution. A shorter time series based on TCEPR and TCER (available since 2007–08) data, analysed at tow by tow resolution, closely resembled the mixed-form series for the years in common.

The NINSWG noted that almost all of the records in the aggregated data had catches of gurnard and that a binomial index was flat. As a result, the positive catch index was retained as the key monitoring series.

The indices were updated in 2018 and 2019, and in 2020 a new fisheries characterisation was also carried out. This indicated that the fishery had been stable in the intervening period, and the accepted indices were updated with the addition of data from the ERS – Trawl reporting regime which was introduced for deepwater vessels from 2017–18, and for all other fisheries during 2019. The indices were further updated in 2022 and 2024, with data to 30 September 2023 (Middleton et al, in prep.).

In the longer CPUE series using aggregated data (i.e., PseudoCELR series) there are indications of cyclical variations in abundance with a 4- to 5-year period (Figure 9). There was an overall decreasing trend in CPUE from 1990 to 2007, after which CPUE stabilised and then increased to 2016, before decreasing to 2017. CPUE then increased to 2020, but dropped in 2021, remaining stable to 2023. The series using tow level data showed a similar pattern to the longer, daily aggregated, index for years after 2007–08 (Figure 9).

In 2022, a further tow level series was investigated, including tarakihi as a target species. Tarakihi-target effort has been excluded from previous GUR 2 CPUE standardisations due to the deeper distribution of tarakihi effort. Depth is available as a covariate in the event-level data and is retained as as an explanatory variable in a mixed-target CPUE series, with tarakihi target effort included. Gurnard are not caught universally in tarakihi-target trawling and a combined (binomial and positive catch) CPUE series is required if tarakihi effort is included. This series shows a generally similar pattern to both the existing series but shows more variation, with lower values during 2009–2013 and higher values during 2017–2023 (Figure 9). It is likely that these differences are attributable to changes in the spatial distribution of gurnard.

Schofield et al (2018a) demonstrated that during 2008 to 2016 there was evidence of a change in distribution of gurnard within Hawke Bay. During 2008–2010 gurnard were most abundant in the shallower coastal waters of Hawke Bay, but there was a shift to deeper water over the period 2011 to 2013, and during 2014 to 2016 gurnard were most abundant offshore, in the deeper parts of Hawke Bay. An updated analysis in 2024 found that during 2016 to 2018 gurnard were still found in greater abundance in the deeper parts of Hawke Bay, but in 2019 to 2021 were more uniformly distributed in Hawke Bay, and were also more abundant in the coastal area between the Māhia Peninsula and Poverty Bay. The impacts of Cyclone Gabrielle in February 2023 led to a reduction in nearer shore fishing effort in the northern parts of Hawke Bay; however, where data were available, they suggested that the gurnard distribution had shifted offshore and that this change was already underway in 2022, i.e. before the cyclone. The original series (without tarakihi effort) therefore remains more appropriate for monitoring, due to the dominance of gurnard target effort in that dataset which adapts to changes in the stock distribution.

Chapman-Robson estimates of total mortality (Z) for GUR 2, based on the age composition of bottom trawl landings in 2009–10, were 0.518 (SE = 0.0159, CV=3.1%) and 0.632 (0.0196, 3.1), depending on whether the age at full recruitment was at 2 or 3 years (Parker & Fu 2012). Assuming an instantaneous rate of natural mortality of 0.307, fishing mortality was estimated to be either 0.189 or 0.303.

Although it was not possible to produce reliable estimates of spawner biomass per recruit based targets of F (due to unreliable estimates of growth rate and size at maturity), estimates of F from this study were either lower than or approximately equal to the estimate of natural mortality (depending on the age at full recruitment assumed). Assuming that the fishery is sampling the age structure of the population, and given that catches and standardised CPUE have been reasonably constant over the last decade, these results suggest that GUR 2 was not over-exploited in 2010, and that the stock is likely to be at or above B_{MSY}.




Figure 9: Comparison of standardised catch per unit effort (CPUE) indices for GUR 2 from bottom trawling targeting gurnard, snapper, and trevally (BT.MIX pseudoCELR; Weibull) combined over all form types, and more recently from data based on TCEPR/ TCER/ER (tow) format data only (BT.MIX event; Weibull). A series using tow format data and including tarakihi as a target species is also illustrated (BT.MIX.TAR event; Weibull). Only a positive catch series is considered for the pseudoCELR data, and only a combined series is presented for the BT.MIX.TAR event series. Both the positive and combined indices are illustrated for the BT.MIX event series. The series are scaled relative to the geometric mean of the years they have in common.

Establishing B_{MSY}-compatible reference points

In 2014, the NINSWG adopted mean CPUE from the (BT(MIX)) model for the period 1990–91 to 2009–10 as a B_{MSY}-compatible proxy for GUR 2. In 2020 the reference period was extended from 1991 to 2018, on the grounds that the new period included two peaks in abundance. The Working Group adopted the default Harvest Strategy Standard definitions for the Soft Limit and Hard Limit of one half and one quarter the target, respectively.

Future research considerations

GUR 3

Prior to 2022, GUR 3 was monitored based on trends in standardised CPUE indices from the main trawl fisheries: the flatfish target fishery and a mixed trawl fishery (targeting red cod, giant stargazer, barracouta, tarakihi, and red gurnard). These indices were derived from data from the main inshore General Statistical Areas of GUR 3 (018, 020, 022, 024, 026, 025, and 030).

In 2022, the CPUE analyses were revised and restricted to Pegasus Bay (Statistical Area 020) and Canterbury Bight (022) to enable a more direct comparison with the ECSI trawl survey biomass indices. The CPUE analyses were also partitioned by depth (proxied by target species) to be comparable with the depth partition of the trawl survey indices. The BT-MIX CPUE series was derived from trawls targeting red cod, tarakihi, and barracouta and is comparable with the longer time series of trawl surveys encompassing the 30–400 m depth range, while the BT-FLA CPUE indices are comparable with the shallower area (10–30 m) sampled by the more recent trawl surveys.

The catch and effort data from the two fisheries were aggregated ('rolled up') to a standard vessel-day format for the period 1989–90 to 2020–21. A core fleet was defined for each data set and CPUE indices were derived using a delta-lognormal modelling approach, including the explanatory variables fishing year, vessel, month, target species, statistical area, and fishing duration to model the probability of red gurnard catch (binomial model) and the magnitude of positive catches (lognormal model).

The two sets of CPUE indices increased considerably over the time series, although the scale of the increase was substantially higher for the BT-FLA (8-fold) compared with the BT-MIX CPUE indices

(3 fold). The BT-MIX indices were the preferred set of indices because they encompassed data from a broader area compared with the BT-FLA fishery. There has also been a marked reduction in the scale of the BT-FLA fishery in recent years due to the recent retirement of several vessels and considerably higher variability in the associated CPUE indices.

The scale of the increase in the BT-MIX CPUE indices is comparable with the overall increase in the biomass indices from the 30–400 m core area of the ECSI trawl survey, although there is some variability in the trends in the two sets of indices particularly during the 2010s (Figure 10).

A separate CPUE analysis was conducted for the FLA trawl fishery in the Otago and Southland area (Statistical Areas 024, 025, 026, and 030) (BT-SouthFLA). From 2000, the indices increased 3–4 fold, with generally similar annual trends in the indices as the BT-FLA and BT-MIX CPUE indices (Figure 11).




Figure 10: A comparison of the standardised BT-MIX CPUE indices and the trawl survey biomass estimates for red gurnard from the winter ECSI inshore trawl survey for the 30–400 m depth strata. Error bars show ±95% confidence intervals. Both sets of indices have been normalised to the average for the years with a survey biomass index.




Figure 11: A comparison of the standardised BT-MIX, BT-FLA, and Southland/Otago FLA (BTSouthFLA) red gurnard CPUE indices.

GUR 7
Previously, CPUE indices were derived for the main trawl fisheries in GUR 7. The most recent analysis was completed in 2017 and included CPUE indices for the west coast South Island flatfish (WCSI-FLA) and mixed (WCSI-MIX) trawl fisheries. The trends in the CPUE indices deviated from the time series of red gurnard biomass indices from the WCSI trawl survey. In 2017, the Plenary adopted the trawl survey biomass indices as the main tool for monitoring GUR 7 and the CPUE indices have not been subsequently updated.

4.3 Stock assessments

GUR 3
The first fully quantitative stock assessment of GUR 3 was conducted in 2022. Previously, GUR 3 was assessed using partial quantitative stock assessments based on standardised CPUE indices.

The stock assessment was conducted using an age-structured population model implemented in Stock Synthesis. The model incorporated data to the 2020–21 fishing year (2021 model year). The input data were limited to commercial catches for 1985–2021, trawl CPUE indices (MIX), and *Kaharoa* ECSI trawl survey biomass indices and length/age compositions (1991–2021).

Catches
Commercial catch data are available for the GUR 3 fishery from 1931. However, the catch data are considered less reliable from the period prior to the introduction of the Quota Management System in 1986. The model was initialised in 1985 assuming equilibrium, exploited conditions. Commercial catch included in the stock assessment is shown in Figure 12. An initial model including the full catch series from 1931 did not perform as well due to implausibly high estimates of exploitation rates.




Figure 12: Commercial catch for GUR 3 included in the stock assessment models.

The model data set was configured to include the red gurnard catch from two commercial trawl fisheries: the flatfish trawl fishery and the deeper mixed trawl fishery. Reported annual catches were increased by 10% to account for an assumed level of under-reporting of catches from the commercial fishery. No length or age composition data are available from the commercial fishery.

Estimates of annual recreational catches of red gurnard from GUR 3 are small (< 5 t) and were not included in the assessment model.

There are no estimates of customary catch available for GUR 3. Recent customary catches are likely to have been a minor component of the total catch and are not explicitly included in the model catch history.

Trawl survey
The *Kaharoa* east coast South Island inshore trawl survey commenced in 1991. Red gurnard is one of the target species for the trawl survey and there is comprehensive biological sampling of the red gurnard catch (including the collection of otoliths). The survey was conducted in winter (April-May) during 1991–1996 and 2007–2021 (N=13). A separate set of trawl surveys was conducted in summer (December) during the intervening years (1997–2001, N= 6).

Initially, the trawl survey area encompassed the 30–400 m depth range. From 2007, the survey area was extended to include the 10–30 m depth range, primarily to improve the monitoring of red gurnard; approximately 50% of the red gurnard trawl survey biomass is within the shallower area of the survey although the proportion varies between surveys. Catches of red gurnard in the deeper portion (> 30 m) of the survey area generally comprised a higher proportion of larger male fish compared with the shallower area of the survey.

Age compositions were available from three early winter surveys (1992, 1993, and 1994) and length compositions were used for the remainder of the surveys.

CPUE indices
CPUE indices were derived from the daily catch and effort data from the mixed inshore bottom trawl fishery in Pegasus Bay/Canterbury Bight. The fishery included trawl records from targeting red cod, barracouta, and tarakihi (BT-MIX). The CPUE indices from the FLA trawl fishery were not included in the assessment modelling due to the recent retirement of several vessels and considerably higher variability in the associated CPUE indices.

Model structure and assumptions
A statistical age-structured population model for GUR 3 was implemented using Stock Synthesis (Methot & Wetzell 2013). Input data are listed in Table 8. The main model structural assumptions for the base model are as follows:

- The initial population (1985) was assumed to be an exploited, equilibrium state. The population is partitioned by sex and consists of 10 age classes, including a plus group. The model data period is 1985–2021 (the 2021 model year represents the 2020–21 fishing year) with no seasonal structure.
- Biological parameters for growth and natural mortality are available for GUR 3 and were fixed (Table 9). Estimated parameters are listed in Table 10.
- Recruitment was parameterised using a Beverton-Holt stock-recruitment relationship (SRR) with a steepness parameter (h) of 0.85. Recruitment deviates were estimated for 1985–2020. Recruitment for 2021 was assumed based on the average level of recruitment from the stock-recruitment relationship.
- For each *Kaharoa* trawl survey series, age based selectivities were parameterised using a sex-specific logistic function. Separate selectivities were estimated for the three sets of surveys: winter 30–400 m, summer 30–400 m, and winter 10–400 m. The trawl survey catchability coefficients (q) were freely estimated.
- A single commercial fishery selectivity was parameterised as an age-based logistic function, equivalent for both sexes. The selectivity parameters were fixed at values to approximate full selectivity of fish at a length of about 30 cm (TL), the minimum length of red gurnard in the landed catch.
- The initial (1985) fishery mortality rates for the two fisheries were estimated, informed by prior distributions that assumed relatively high initial levels of fishing mortality.
- The BT-MIX CPUE indices were assigned a CV of 0.20 based on the RMSE of the fit to the indices from preliminary model runs.

- The *Kaharoa* trawl survey length compositions were assigned an Effective Sample Size (ESS) of 10, while the age compositions were assigned a higher weighting (ESS 20). These ESS values generally approximate the relative weighting of the compositional data recommended following the approach of Francis (2011). Further evaluation of the relative weighting of the compositional data was conducted as sensitivities to the base assessment model.

The base model provided a good fit to the time series of CPUE indices. The model fitted the general increase in the 30–400 m winter trawl survey biomass indices, although it under-estimated the overall magnitude of the increase between the two time blocks (1991–1996 and 2007–2021) (Figure 13). The model also fitted the two shorter series of trawl survey biomass indices (summer 30–400 m and winter 10–400 m) and provided a reasonable fit to associated age and length composition data from the trawl surveys.

Table 8: Summary of input data sets for the Base Case assessment model. The relative weighting includes the Effective Sample Size (ESS) of age/size composition data and the coefficient of variation (CV) associated with the abundance data. Note that model year 2021, is fishing year 2020–21, and includes the trawl survey conducted in March 2021. Nobs is number of observations.

Data set	Model years	Nobs	Error structure	CVs/ESS
CPUE indices BT-MIX	1990–2021	32	Lognormal	0.20
Trawl survey indices				
Winter 30–400m	1991, 1992, 1993, 1994, 1996, 2007, 2008, 2009, 2012, 2014, 2016, 2018, 2021	13	Lognormal	0.19–0.35
Summer 30–400m	1997, 1998, 1999, 2000, 2001	5	Lognormal	0.13–0.34
Winter 10–400m	2007, 2012, 2014, 2016, 2018, 2021	6	Lognormal	0.15–0.27
Trawl survey age comp				
Winter 30–400m	1992, 1993, 1994	3	Multinomial	ESS 20
Trawl survey length comp				
Winter 30–400m	1991, 1996, 2007, 2008, 2009, 2012, 2014, 2016, 2018, 2021	10	Multinomial	ESS 10
Summer 30–400m	1997, 1998, 1999, 2000, 2001	5	Multinomial	ESS 10
Winter 10–400m	2007, 2012, 2014, 2016, 2018, 2021	6	Multinomial	ESS 10

Table 9: Details of parameters that were fixed in the base model.

Natural mortality	female	0.29 y^{-1}
	male	0.35 y^{-1}
Stock-recruit steepness (Beverton & Holt)		0.85
Std deviation of rec devs (sigmaR)		0.6
Proportion mature		0 for age 1, 0.5 for age 2, 1 for ages > 2
Length-weight [mean weight (kg) = a (length (cm))^b]		$a = 5.3 \times 10^{-6}$, $b = 3.19$
Growth parameters - female		$L_\infty = 48.2$, $k = 0.44$, Length1=19.4
	male	$L_\infty = 42.2$, $k = 0.49$, Length1=19.4
Coefficients of variation for length-at-age		0.10
Selectivity BT commercial		2

Table 10: Estimated parameters for the base model.

Parameter	Number of parameters	Parameterisation, priors, constraints
LnR_0	1	Uniform, uninformative
Rec devs (1980–2020)	41	SigmaR 0.6
Selectivity trawl surveys (3)	8	Logistic, female offset
Trawl Survey ln q	3	Free
Initial F	2	Normal (0.4, 0.2)

The inclusion of data from the separate 10–400 m and 30–400 m trawl survey series duplicates the data from the deeper area of the survey from overlapping years (2007–2021) rather than configuring the recent data as a separate time series of 10–30 m trawl survey indices. The inclusion of the entire depth range of the recent trawl surveys enables the whole population of red gurnard to be monitored as a single index (with the assumption of full selectivity of older fish). A sensitivity run with the 10–30 m and 30–400 m trawl surveys gave similar results.

The CPUE and trawl survey indices indicate that stock abundance has been higher since the late 2000s. The model estimates that the increase in abundance is attributable to higher recruitment in the mid-2000s and from 2010 to 2019, with exceptionally high recruitments in 2016 and 2019 (Figure 14). There are no age composition data available to inform the estimates of recruitment, although the increase in recruitment is consistent with the substantial increase in the trawl survey biomass indices for smaller (less than 30 cm) red gurnard from 2007 (the pre-recruit index was not explicitly fitted in the model).




Figure 13: The fit (MPD) to the trawl survey biomass indices (points) for the base model option. W=winter and S=summer.




Figure 14: Annual recruitment for the base model (MCMC results). Recruitment deviates were estimated for 1985–2020. The line represents the median and the shaded area represents the 95% credible interval.

During the model development a wide range of model options were evaluated to investigate key model assumptions. A set of model options were retained as key sensitivities to capture plausible uncertainty in assessment results, including: a lower (0.25) value of natural mortality (compared with females = 0.29 and males = 0.35), excluding the CPUE indices from 2007 onwards, and limiting the model catch to the FMA 3 (ECSI) portion of GUR 3 (Table 11). FMA 5 has accounted for approximately 25% of the GUR 3 catch since the late 1990s; however, this area is not monitored by the CPUE and trawl survey abundance indices. The sensitivities were treated as single changes from the base model.

Table 11: Description of model sensitivities.

Sensitivity run	Description
ExCPUE2007	Exclude 2007–2021 CPUE indices
FMA3catch	Exclude non FMA3 catch
LowM	Natural mortality *M* 0.25

Stock status (current 2021 = 2020–21 fishing year and forecast to 2025–26) for the GUR 3 spawning biomass was reported relative to the default hard limit of 10% SB_0 and the default soft limit of 20% SB_0 and interim target biomass level of 35% SB_0. Fishing mortality (2021) was reported relative to $F_{SB40\%}$ reference level. The reference points represent the default values for a medium productivity stock as described by the Harvest Strategy Standard Operational Guidelines.

For the base model and the range of model sensitivities, biomass is estimated to have increased considerably from 2007 and current (2021) stock status is estimated to be well above the target biomass level although there is considerable uncertainty associated with the estimate of current stock status (Table 12, see Figure 15).

For all model options, current rates of fishing mortality are near the fishing mortality threshold ($F_{SB40\%}$) (Table 12, see Figure 16).

For all model options, estimates of current and equilibrium yield were derived for the stock based on the fishing mortality rate that corresponds to the interim target biomass level (Table 13). Equilibrium yields at the interim target biomass level are estimated to be about 800–1000 t per annum (for GUR 3). $F_{SB40\%}$ yields at 2020–21 biomass levels are substantially higher than the equilibrium yields (about 1400–1800 t).

Table 12: Estimates of current (2020–21) and virgin spawning biomass (t) (median and the 95% confidence interval from the MCMCs) and probabilities of current biomass being above specified levels and probability of fishing mortality being below the level of fishing mortality associated with the interim target biomass level.

Model option	SB_0	SB_{2021}	SB_{2021}/SB_0	Pr(SB_{2021} > X% SB_0)	Pr(SB_{2021} < X% SB_0)	
				35%	20%	10%
Base	6 307	4 061	0.644	0.999	0	0
	(5 552–7 749)	(2 500–6 808)	(0.435–0.924)			
CatchFMA3	4 804	3 083	0.638	0.995	0	0
	(4 250–5 814)	(1 903–5 096)	(0.434–0.906)			
exCPUE2007	6 475	4 397	0.684	0.997	0.001	0.001
	(5 658–7 761)	(2 542–7 531)	(0.428–0.993)			
LowM	6 164	3 123	0.506	0.960	0.007	0.004
	(5 592–6 969)	(1 941–4 784)	(0.325–0.722)			

	$F_{SB40\%}$	$F_{2021}/F_{SB40\%}$	Pr(F_{2021} < $F_{SB40\%}$)
Base	0.2028	0.996	0.506
		(0.627–1.513)	
CatchFMA3	0.2028	1.004	0.492
		(0.645–1.517)	
exCPUE2007	0.2028	0.925	0.632
		(0.569–1.499)	
LowM	0.1857	1.297	0.087
		(0.884–1.987)	

Table 13: Estimates of equilibrium yield (t) at $F_{SB40\%}$ and yield at the 2020–21 biomass levels for the base model and the model sensitivities. The values represent the median and the 95% confidence interval from the MCMCs.

Model option	$F_{SB40\%}$	
	Equilibrium Yield	Yield at current biomass
Base	986	1 617
	(867–1211)	(1 411–1 855)
CatchFMA3	751	1 229
	(664–908)	(1 069–1 397)
exCPUE2007	1 012	1 655
	(884–1 213)	(1 451–1 888)
LowM	932	1 632
	(846–1 054)	(1 390–1 865)




Figure 15: Annual trend in spawning biomass relative to the 35% SB_0 interim target biomass level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The projection period (2022–2026) is in red. The dashed line represents the interim target level.




Figure 16: Annual trend in fishing mortality relative to the $F_{SB40\%}$ interim threshold exploitation level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The projection period (2022–2026) is in red. The dashed line represents the interim threshold level.

Projections

Projections were conducted for the base model. Stock projections were conducted for the 5-year period following the terminal year of the model (i.e., 2022–2026). Projections assumed that future recruitments were resampled from the average level of recent recruitment (2011–2020) and variation equivalent to sigmaR. Commercial catches in the projection period were held constant at the current TACC of 1500 t with an additional 10% allowance for unreported catch.

For the base case, stock abundance is predicted to remain at about the current level during the projection period, well above the target biomass level (Table 14). However, there is considerable uncertainty in the stock projections due to the uncertainty associated with the estimates of recent recruitment (especially the strong 2016 and 2019 year classes).

Table 14: Estimates of projected (2025–26) spawning biomass (t) (median and the 95% confidence interval from the MCMCs) and probability of the spawning biomass being above default biomass limits and the interim target level in 2026 from the base model projections.

Model option	SB_{2026}/SB_0	$Pr(SB\ 2026 < X\%\ SB_0)$		$Pr(SB\ 2026 > X\%$
		10%	20%	35%
Base	0.609 (0.337–0.970)	0	0	0.970

Qualifying comments

The base model includes the catch from the entire GUR 3 QMA, whereas the trawl survey and CPUE indices are limited to FMA 3. Since 2000, approximately 25% of the GUR 3 catch has been taken from outside FMA 3 (principally in Southland FMA 5). Limiting the model catch to FMA 3 did not appreciably change the estimate of current stock status.

There are no direct observations to determine the selectivity of the commercial fishery and the commercial catch is assumed to comprise fish larger than 30 cm.

Reference points (SB_0 based) are derived from the average level of recruitment during 1985–2020. There was an apparent shift in the productivity of GUR 3 from the mid-2000s. The increasing catches since the mid-2000s were sustained by the higher level of recruitment. Recent strong recruitments (in 2016 and 2019) are supported by higher abundance of pre-recruit red gurnard from recent trawl surveys but are yet to be confirmed from age sampling.

The model does not include GUR 3 catches prior to 1987 and the initial level of fishing mortality (in 1985) is estimated.

There are large differences between MPD and MCMC results for R_0 that require further investigation.

Future research considerations

- Determine the age compositions of red gurnard sampled by the ECSI trawl surveys, particularly for one or two of the most recent surveys. These data will improve estimates of recent recruitments, provide for more recent estimates of growth rate and the CV of length at age, and potentially resolve the differences between MPD and MCMC results. Also, consider re-examining the earlier 1992–1994 age estimates.
- Determine the length/age composition of the east coast commercial catch of GUR 3 from primarily the BT-MIX but also the BT-FLA trawl fisheries to inform the model regarding the selectivity of the two fisheries.
- Consider standardising the length frequency and age survey data outside the model.
- Consider spatio-temporal modelling of the trawl survey biomass data potentially including environmental covariates (INLA or VAST).
- Re-examine summer trawl survey data for maturity and the winter survey data for changes in the length-weight relationships.
- Investigate stock relationships between FMAs 3 and 5.

- Explore changes in commercial mesh sizes over time.

GUR 7

The first fully quantitative stock assessment of GUR 7 was conducted in 2022; previous assessments were partial quantitative assessments based on the WCSI trawl survey series. The stock assessment was conducted using an age-structured population model implemented in Stock Synthesis. The model incorporated data to the 2020–21 fishing year (2021 model year). The input data were limited to commercial catches (1987–2021) and *Kaharoa* WCSI trawl survey biomass indices and length/age compositions (1992–2021, N = 15).

Catches

Commercial catch data are available for the GUR 7 fishery from 1931. However, the catch data are considered less reliable from the period prior to the introduction of the Quota Management System in 1986. There are no other composition or abundance data available prior to 1987. The model was initialised in 1987 assuming equilibrium, exploited conditions. Catch data used in the stock assessment are presented in Figure 17.

The model data set was configured to include a single commercial trawl fishery. There was no allowance for the under-reporting of catches from the commercial fishery. No catch composition (length or age) data are available from the commercial fishery.

Estimates of annual recreational catches of red gurnard from GUR 7 are relatively small (10–48 t) and were not included in the assessment model.

There are no estimates of customary catch available for GUR 7. Recent customary catches are likely to have been a minor component of the total catch and are not explicitly included in the model catch history.




Figure 17: Commercial catch for GUR 7 included in the stock assessment models.

Trawl survey

The west coast South Island inshore trawl survey, including the Tasman Bay/Golden Bay area, commenced in 1992 and has been conducted biennially since 2002. Red gurnard is one of the target species for the trawl survey; red gurnard is one of the main species caught and there is comprehensive sampling of the red gurnard catch (including the collection of otoliths).

Age compositions were available from five surveys (1994, 1995, 2003, 2005, and 2007) and length compositions were available for the remainder of the surveys.

Model structure and assumptions

A statistical age-structured population model for GUR 7 was implemented using Stock Synthesis (Methot & Wetzell 2013). Input data are listed in Table 15. The main model structural assumptions for the base model were as follows:

- The initial population (1987) was assumed to be in an exploited, equilibrium state. The population was partitioned by sex and consists of 10 age classes, including a plus group. The model data period was 1987–2021 (the 2021 model year represents the 2020–21 fishing year) with no seasonal structure.
- Biological parameters for growth and natural mortality are available for GUR 7 and were fixed (Table 16). Estimated parameters are listed in Table 17.
- Recruitment was parameterised using a Beverton-Holt stock-recruitment relationship (SRR) with a beta prior assumed for the steepness parameter. Recruitment deviates were estimated for 1987–2020. Recruitment for 2021 was assumed based on the average level of recruitment from the stock-recruitment relationship.
- Age based selectivity for the *Kaharoa* trawl survey was parameterised using a sex-specific logistic selectivity function. The trawl survey catchability coefficient (q) was parameterised with a relatively uninformative prior.
- Commercial fishery selectivity was an age-based logistic function with the two parameters informed by priors that approximate full selectivity of fish at a length of about 30 cm (TL), the minimum size of red gurnard in the landed catch. No length or age data were available from the commercial fishery.
- The initial (1987) fishing mortality rate was estimated, informed by a prior distribution that assumed a relatively high initial fishing mortality rate.
- The trawl survey age and length compositions were assigned a relative high weighting (Effective Sample Size (ESS) of 50). The high weighting reflected the comprehensive sampling conducted for each survey and ensured that the model was informed by these data.

Table 15: Summary of input data sets for the Base Case assessment model. The relative weighting includes the Effective Sample Size (ESS) of age/size composition data and the coefficient of variation (CV) associated with the abundance data. Note that model year 2021, is fishing year 2020–21. Nobs is number of observations.

Data set	Model years	Nobs	Error structure	CV/ESS
Trawl survey indices	1992, 1994, 1995, 1997, 2000, 2003, 2005, 2007, 2009, 2011, 2013, 2015, 2017, 2019, 2021	15	Lognormal	0.12–0.19
Trawl survey age comp	1994, 1995, 2003, 2005, 2007	5	Multinomial	ESS 50
Trawl survey length comp	1992, 1997, 2000, 2009, 2011, 2013, 2015, 2017, 2019, 2021	10	Multinomial	ESS 50

Table 16: Details of parameters that were fixed in the base model.

Natural mortality (male and female)	0.31 y^{-1}
Std deviation of rec devs (sigmaR)	0.6
Proportion mature	0 for age 1, 0.5 for age 2, 1 for ages > 2
Length-weight [mean weight (kg) = a (length (cm))^b]	$a = 5.3 \times 10^{-6}$, $b = 3.19$
Growth parameters - female	$L_\infty = 45.7$, $k = 0.40$, Length1=19.4
male	$L_\infty = 40.3$, $k = 0.37$, Length1=19.4
Coefficients of variation for length-at-age	0.10

Table 17: Estimated parameters for the base model.

Parameter	Number of parameters	Parameterisation, priors, constraints
LnR_0	1	Uniform, uninformative
Stock-recruit steepness (Beverton & Holt)	1	Beta(0.80, 0.125)
Rec devs (1987–2020)	34	SigmaR 0.6
Selectivity trawl survey	3	Logistic, female offset
Selectivity BT commercial	2	Logistic
Trawl Survey ln q	1	Normal (-1.1, 1)
Initial F	1	Normal (0.4, 0.2)

For the base model option, the model also provides a good fit to the time series of trawl survey biomass indices, with the exception of the 2003 trawl survey index (Figure 18). The model also provides an acceptable fit to associated age and length composition data.

The trawl survey indices increased considerably from 2009 to 2021. The model estimates that the increase in abundance was due to a considerably higher level of recruitment from 2008 onwards. There are age data from 5 surveys (1994, 1995, 2003, 2005, 2007) but all remaining surveys have length data. These are used by the model to inform recruitment (Figure 19). Recruitments since 2007 are estimated with lower precision. The increase in recruitment is, however, consistent with the substantial (3–4 fold) increase in the trawl survey biomass indices for smaller (less than 30 cm) red gurnard from 2007.




Figure 18: The fit (MPD) to the trawl survey biomass indices (points) for the base model option.




Figure 19: Annual recruitment for the base model (MCMC results). Recruitment deviates were estimated for 1987–2020. The line represents the median and the shaded area represents the 95% credible interval.

A range of model sensitivities was undertaken to investigate the key model assumptions; these included: lower (0.25) and higher (0.35) values of natural mortality (compared with 0.31), the prior on the initial (1987) level of fishing mortality, and the precision of the trawl survey biomass indices (Table 18). The sensitivities were implemented as single changes from the base model.

Additional model trials were conducted relating to the selectivity of the commercial fishery, the relative weighting of the length/age composition data from the trawl survey, and relaxing the prior on trawl survey q. These changes did not appreciably change the model results.

Table 18: Description of model sensitivities.

Sensitivity run	Description
InitialFlow	Initial *F* prior Norm(0.2,0.2)
Mhigh	Natural mortality *M* 0.35
Mlow	Natural mortality *M* 0.25
Trawl Survey Indices Process Error	Process Error 0.1

Stock status (current 2021 = 2020–21 fishing year and forecast to 2025–26) for the GUR 7 spawning biomass was reported relative to the default hard limit of 10% SB_0 and the default soft limit of 20% SB_0 and interim target biomass level of 35% SB_0, and fishing mortality (2021) was reported relative to $F_{SB40\%}$. The interim reference points represent the default values for a medium productivity stock as described by the Harvest Strategy Standard Operational Guidelines.

For the base model and the range of model sensitivities, biomass is estimated to have increased considerably from 2010 and current (2021) stock status is estimated to be at about the equilibrium, unexploited level (i.e. SB_0) (Table 19, Figure 20).

For all model options, current rates of fishing mortality are well below the fishing mortality threshold ($F_{SB40\%}$) (Table 19, Figure 21).

Table 19: Estimates of current (2020–21) and virgin spawning biomass (t) (median and the 95% confidence interval from the MCMCs) and probabilities of current biomass being above specified levels and probability of fishing mortality being below the level of the fishing mortality threshold.

Model option	SB_0	SB_{2021}	SB_{2021}/SB_0	Pr(SB_{2021} > X% SB_0)	Pr(SB_{2021} < X% SB_0)	
				35%	20%	10%
Base	4 990	5 528	1.108	1.00	0	0
	(4 226–6 247)	(3 779–7 919)	(0.799–1.44)			
InitialFlow	4 933	5 528	1.125	1.00	0	0
	(4 217–6 131)	(3 832–7 914)	(0.804–1.457)			
Mhigh	5 365	6 830	1.262	1.00	0	0
	(4 318–7 346)	(4 369–10 504)	(0.921–1.605)			
Mlow	4 939	4 369	0.885	1.00	0	0
	(4 313–5 981)	(3 045–6 070)	(0.63–1.147)			
TSurveyProcessError	4 421	4 034	0.901	1.00	0	0
	(3 772–5 411)	(2 499–6 482)	(0.616–1.297)			

	$F_{SB40\%}$	$F_{2021}/F_{SB40\%}$	Pr(F_{2021} < $F_{SB40\%}$)
Base		0.44	
	0.2248	(0.296–0.686)	1.00
InitialFlow		0.44	
	0.2267	(0.294–0.691)	1.00
Mhigh		0.342	
	0.2342	(0.219–0.551)	1.00
Mlow		0.624	
	0.2016	(0.447–0.908)	0.99
TSurveyProcessError		0.593	
	0.2298	(0.359–0.957)	0.98




Figure 20: Annual trend in spawning biomass relative to the 35% SB_0 interim target biomass level for the base model. The line represents the median and the shaded area represents the 95% confidence interval. The projection period (2022–2026) is in red. The dashed line represents the interim target level.




Figure 21: Annual trend in fishing mortality relative to the $F_{SB40\%}$ threshold level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The projection period (2022–2026) is in red. The dashed line represents the interim threshold level.

For all model options, estimates of current and equilibrium yield were derived for the stock based on the fishing mortality rate that corresponds to the interim target biomass level (Table 20). Equilibrium yields at the interim target biomass level are estimated to be about 800–1000 t per annum. $F_{SB40\%}$ yields at 2020–21 biomass levels are substantially higher than the equilibrium yields (about 2000–3000 t), although the magnitude of the current $F_{SB40\%}$ yields is highly uncertain (1100–6500 t).

Table 20: Estimates of equilibrium yield (t) at $F_{SB40\%}$ and at the 2020–21 biomass levels for the base model and the model sensitivities. The values represent the median and the 95% confidence interval from the MCMCs.

Model option	$F_{SB40\%}$	
	Equilibrium Yield	Yield at current biomass
Base	953 (811–1 149)	2 748 (1 609–4 681)
InitialFlow	952 (808–1 160)	2 750 (1 583–4 957)
Mhigh	1 105 (891–1 453)	3 621 (1 964–6 547)
Mlow	807 (726–906)	1 875 (1 163–2 819)
TSurveyProcessError	861 (749–1 043)	2 020 (1 094–3 934)

Projections

Projections were conducted for the base model. Stock projections were conducted for the 5-year period following the terminal year of the model (i.e., 2022–2026). Projections assumed future recruitments were resampled from the average level of recent recruitment (2011–2020) and variation equivalent to sigmaR. Commercial catches in the projection period were held constant at the current TACC of 1298 t. There was no explicit allowance for unreported catch, recreational catch, or customary catch.

For the base case, stock abundance is predicted to decline during the projection period, although the biomass remains at about the SB_0 reference level throughout the period (Table 21).

Table 21: Estimates of projected (2025–26) spawning biomass (t) (median and the 95% confidence interval from the MCMCs) and probability of the spawning biomass being above default biomass limits and the interim target level in 2026 from the base model projections.

Model option	SB_{2026}/SB_0	$Pr(SB\ 2026 > X\%\ SB_0)$		$Pr(SB\ 2026 < X\%$
		10%	20%	35%
Base	0.969 (0.667–1.376)	0	0	1.00

Qualifying comments

There are no direct observations to determine the selectivity of the commercial fishery and the commercial catch is assumed to be composed of fish larger than 30 cm.

Reference points (SB_0 based) are derived from the average level of recruitment during 1987–2020. There was an apparent shift in the productivity of GUR 7 in the late 2000s. The increasing catches since 2012 were sustained by the higher level of recruitment.

The model does not include GUR 7 catches prior to 1987 and the initial level of fishing mortality (in 1987) is estimated. Prior to 1987, average annual catches were about 400 t from the early 1950s.

Because no allowance was made for unreported catch or for non-commercial harvest, the predicted yields relate only to the reported commercial catch.

Future research considerations

- Determine the age compositions of red gurnard sampled by the WCSI trawl surveys, particularly for one or two of the most recent surveys. Also consider re-examining age compositions for the earlier surveys.
- Determine length/age composition of the catch from the GUR 7 trawl fishery to inform the model regarding the selectivity of the fishery.
- Re-examine development of a CPUE series, potentially starting in 2008. This may help to calibrate *Kaharoa* replacement.

Management procedure (MP)

A range of management procedures were developed for GUR 7 based on the empirical linear relationship derived between the WCSI trawl survey biomass indices and the corresponding annual reported commercial catch (Figure 22). The relationship was forced through the origin to ensure that catches were aligned with low survey biomass levels. The relationship provides a moderate fit to the observed data although there are a subset of records with a higher catch than predicted from the linear model.




Figure 22: Comparison of red gurnard WCSI trawl survey biomass indices and GUR 7 commercial catch from the corresponding fishing year. The grey line represents the linear relationship constrained to the origin (zero intercept). The slope of the line is denoted F_{MSY} proxy.

Conceptually, the relationship represents a constant level of fishing mortality (F_{MSY} proxy). The simplest management procedures scaled the base fishing mortality level using a scalar applied to the slope coefficient (base) (0.8, 1.0, 1.2….2.0, 2.5). More complex management procedures incorporated declining levels of fishing mortality below threshold levels of the observed trawl survey biomass indices, with inflection points at the lower quartile, median or upper quartile (Figure 23).

Simulation-testing of the management procedures was carried out using the MCMC output from the 2022 GUR 7 assessment model. The individual (1000) MCMC samples were used to initialise the population age structure and projections were conducted over 100 years with recruitment resampled from the estimated recruitment deviates. Trawl surveys were conducted biennially during the projection period and the specific management procedure was applied to set the TACC for the next two years following each trawl survey (conducted in March-April).

$$TACC_{(year + 1, year + 2)} = Scalar \times F_{MSY} \text{ proxy} \times TrawlSurveyBiomass_{year}$$

Constraints were applied to investigate the effect of limiting the scale of the changes in the TACC (maximum change of ±20% or 50%). Two alternative periods were applied to define the projected recruitments: the entire recruitment period (1987–2020) and the lower recruitment period 1987–2007. The MPs were also tested with different levels of autocorrelation between annual recruitments and additional process error associated with the trawl survey biomass estimates.

The results of the MP evaluation indicated that a Scalar of 1.6–2.0 would yield results that were consistent with fishing the stock at the $F_{SB40\%}$ fishing mortality level, maintaining the stock at or above the HSS default target biomass level of 35% SB_0 (moderate productivity) and well above the 20% SB_0 soft limit. There was no appreciable improvement in the performance of the MP with the inclusion of more complex relationships between trawl survey biomass and F_{MSY} proxy (i.e., inflection points at different biomass levels).

If recruitment was to revert to the lower (1987–2007) level, then the stock would fluctuate about the 30% SB_0 level with an increased probability of the stock declining below the soft limit. Nonetheless, since 2008 recruitment has been maintained at an appreciably higher level (approximately doubled) and future recruitment levels would continue to be monitored via the trawl survey. The MP could be modified if a sustained period of lower recruitment is observed (via a breakout rule).

This MP should be evaluated about every 5 years (after 2 surveys), which should include a new stock assessment.




Figure 23: The range of scalars applied to the F_{MSY} proxy value in the suite of management procedures. The different functional forms of the scalars have inflection points at different levels of trawl survey biomass from the reference data set (15 surveys).

Breakout rules

- Review of the biennial trawl survey results by the Inshore Working Group, including determination of the magnitude of recent recruitment (pre-recruit biomass), survey precision and investigation of extreme catchability.

4.4 Other factors

Red gurnard is a major bycatch of target fisheries for several different species, such as snapper and flatfish. The target species may differ between areas and seasons. The recorded landings are influenced directly by changes in the fishing patterns of fisheries for these target species and indirectly by the abundance of these target species. Some target fishing for gurnard also occurs.

5. STATUS OF THE STOCKS


Stock Structure Assumptions

For the purpose of this summary GUR 1 is considered to be a single stock with three sub-stocks.


- **GUR 1W**

Stock Status		
Most Recent Assessment Plenary Publication	2022	
Catch in most recent year of assessment	Year: 2020–21	Catch: 544 t
Assessment Runs Presented	Standardised CPUE and WCNI trawl survey series	
Reference Points	Target: B_{MSY}-compatible proxy based on the mean CPUE from 1995–96 to 2011–12 of the bottom trawl GUR 1 west (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY}-compatible proxy based on the mean relative exploitation rate for the period: 1995–96 to 2011–12	
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target	
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below	
Status in relation to Overfishing	Overfishing is About as Likely as Not (40–60%) to be occurring	

Historical Stock Status Trajectory and Current Status



Landings (dashed brown line), standardised CPUE combined model using tow by tow data (blue line), and trawl survey recruited biomass indices (purple with error bars ± two standard deviations). The green, orange, and red horizontal lines represent the target, soft, and hard limits, respectively. Trawl survey recruited biomass indices are scaled to have the same arithmetic mean value as the standardised CPUE indices, over the years they have in common.



Annual relative exploitation rate (landings divided by standardised CPUE and normalised to an arithmetic mean of one over the reference period 1996 to 2012) for red gurnard in the GUR 1 west coast sub-stock. The horizontal grey dashed line at one represents the average relative exploitation rate during the period used to define the reference points (depicted by vertical dotted green lines).

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The CPUE has increased since 2008–09 and in 2020–21 was well above the long-term mean. The WCNI trawl survey series shows that abundance in 2018–2020 was lower than it was during the early 1990s.
Recent Trend in Fishing Intensity or Proxy	Relative exploitation rate declined from 1995–96 to 2018–19, and then increased to 2020–21, but remained below the threshold.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Current Catch Soft Limit: Unlikely (< 40%) Hard Limit: Unlikely (< 40%) TACC Unknown for both the Soft and Hard Limits
Probability of Current Catch or TACC causing Overfishing to continue or to commence	About as Likely as Not (40–60%) if the catch remains at current levels Unknown if the catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from bottom trawl	
Assessment Dates	Latest assessment Plenary publication year: 2022	Next assessment: 2025
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	Catch and effort data WCNI trawl survey series	1 – High Quality 1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	- Inclusion of WCSI trawl survey series	
Major Sources of Uncertainty	-	


Qualifying Comments
As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation. This makes it difficult to predict future trends without recruitment information. The WCNI trawl survey series shows a substantial decline in abundance between the 1990s (prior to the commencement of the CPUE series) and 2018–2020, leading to the Working Group moderating assessment results based on CPUE alone. As the TACC for GUR 1 is substantially higher than the current catch, it is not possible to evaluate potential impacts if catches increased to the level of the TACC.

Fishery Interactions
Red gurnard is taken off the west coast by bottom trawl targeted at snapper and trevally. A Danish seine summer fishery for red gurnard and John dory also occurs off the west coast.


- **GUR 1E**

Stock Status		
Most Recent Assessment Plenary Publication Year	2022	
Catch in most recent year of assessment	Year: 2020–21	Catch: 85 t
Assessment Runs Presented	- Standardised CPUE - HG trawl survey series	
Reference Points	Target: B_{MSY}-compatible proxy based on the mean CPUE from 1995–96 to 2011–12 for the bottom trawl GUR 1 East (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period: 1995–96 to 2011–12	
Status in relation to Target	Unknown	
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Unlikely (< 40%) to be below	
Status in relation to Overfishing	Unknown	

Historical Stock Status Trajectory and Current Status



Landings (dashed brown line), standardised CPUE combined model using tow by tow data (blue line), and trawl survey recruited biomass indices (purple with error bars ± two standard deviations). The green, orange, and red horizontal lines represent the target, soft, and hard limits, respectively. Trawl survey recruited biomass indices are scaled to have the same arithmetic mean value as the standardised CPUE indices, over the years they have in common.



Annual relative exploitation rate (landings divided by standardised CPUE and normalised to an arithmetic mean of one over the reference period 1996 to 2012) for red gurnard in the GUR 1 east coast sub-stock. The horizontal grey dashed line at one represents the average relative exploitation rate during the period used to define the reference points (depicted by vertical dotted green lines). For fishing years 2018 to 2021, standardised CPUE and the annual relative exploitation rate are plotted with a cross (see Qualifying Comment below).

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	CPUE declined strongly from 2017–18 to 2020–21, but the trawl survey series showed that abundance in 2020 and 2021 was close to the series mean for the period 1984–85 to 2021. The amount of data available for CPUE declined substantially in the last four years of the series.
Recent Trend in Fishing Intensity or Proxy	Relative exploitation rate declined from 1995–96 to 2002–03, then fluctuated without trend below the long-term average, until rising from 2017–18.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown if the catch remains at current levels

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from bottom trawl	
Assessment Dates	Latest assessment Plenary publication year: 2022	Next assessment: 2025
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	• Inclusion of HG trawl survey series	

Major Sources of Uncertainty	-


Qualifying Comments
As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation. This makes it difficult to predict future trends without recruitment information. Largely due to vessels switching to PSH gear, bottom trawl catch and effort declined substantially after 2016–17, and the standardised CPUE may not track abundance.

Fishery Interactions
Red gurnard is taken as a bycatch on the east coast mainly by bottom longlines targeted at snapper, with the balance taken almost equally by bottom trawl and Danish seine targeting snapper and John dory.


- **GUR 1 Bay of Plenty**

Stock Status		
Most Recent Assessment Plenary Publication Year	2022	
Catch in most recent year of assessment	Year: 2020–21	Catch: 152 t
Assessment Runs Presented	- Standardised CPUE - BoP trawl survey series	
Reference Points	Target: B_{MSY}-compatible proxy based on the mean CPUE from 1995–96 to 2011–12 for the bottom trawl GUR 1 BoP (tow) series Soft Limit: 50% of target Hard Limit: 25% of target Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period: 1995–96 to 2011–12	
Status in relation to Target	Unknown	
Status in relation to Limits	Soft Limit: Unlikely (< 40%) to be below Hard Limit: Very Unlikely (< 10%) to be below	
Status in relation to Overfishing	Unknown	

Historical Stock Status Trajectory and Current Status



Landings (dashed brown line), standardised CPUE combined model using tow by tow data (blue line), and trawl survey recruited biomass indices (purple with error bars ± two standard deviations). The green, orange and red horizontal lines represent the target, soft and hard limits, respectively. Trawl survey recruited biomass indices are scaled to have the same arithmetic mean value as the standardised CPUE indices, over the years they have in common.



Annual relative exploitation rate (landings divided by standardised CPUE and normalised to an arithmetic mean of one over the reference period 1996 to 2012) for red gurnard in the Bay of Plenty. The horizontal grey dashed line at one represents the average relative exploitation rate during the period used to define the reference points (depicted by vertical dotted green lines).

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	The CPUE index has fluctuated without trend, but the trawl survey series showed that abundance in 2019–20 and 2020–21 was below the mean for the period 1989–90 to 1998–99.
Recent Trend in Fishing Intensity or Proxy	Unknown
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unknown Hard Limit: Unknown
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unknown if the catch remains at current levels Unknown if the catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE based on positive catches from bottom trawl	
Assessment Dates	Latest assessment Plenary publication year: 2022	Next assessment: 2025
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	- Catch and effort data	1 – High Quality
Data not used (rank)	-	
Changes to Model Structure and Assumptions	- Inclusion of the re-instated BoP trawl survey series.	
Major Sources of Uncertainty	-	

Qualifying Comments
- As the red gurnard fishery in FMAs 1 and 9 has a long history, it is difficult to infer stock status from recent abundance trends. The abundance of all three sub-stocks appears to be cyclical, probably in response to recruitment variation. This makes it difficult to predict future trends without recruitment information. - CPUE fluctuated without trend, whereas trawl survey biomass in 2020–21 was lower than the average for the period 1995–96 to 2020–21.

Fishery Interactions
Red gurnard is taken as a bycatch in the Bay of Plenty mainly by bottom longline targeted at snapper, with the balance taken almost equally by bottom trawl and Danish seine targeting snapper and John dory.


- **GUR 2**

Stock Structure Assumptions
For the purpose of this summary GUR 2 is considered to be a single stock.

Stock Status		
Most Recent Assessment Plenary Publication Year	2024	
Catch in most recent year of assessment	Year: 2022–23	Catch: 338 t
Assessment Runs Presented	Standardised CPUE for daily BT.MIX series up to 2022–23	
Reference Points	Target: B_{MSY}-compatible proxy based on the geometric mean CPUE (BT.MIX pseudoCELR) for period 1990–91 to 2017–18 Soft Limit: 50% of target Hard Limit: 25% of target	

	Overfishing threshold: F_{MSY} compatible proxy based on the mean relative exploitation rate for the period 1990–91 to 2017–18
Status in relation to Target	About as Likely as Not (40–60%) to be at or above the target
Status in relation to Limits	Soft Limit: Very Unlikely (< 10%) to be below Hard Limit: Very Unlikely (< 10%) to be below
Status in relation to Overfishing	Overfishing is Unlikely (< 40%) to be occurring

Historical Stock Status Trajectory and Current Status



Fishing year

a) Annual removals for GUR 2; (b) the standardised catch per unit effort (CPUE) index, relative to the agreed reference points, for GUR 2 from bottom trawling targeting gurnard, snapper and trevally (BT.MIX) and combining data from all form types at a daily aggregation. (c) annual relative exploitation rate (catch/CPUE) gurnard in GUR 2.

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Biomass has fluctuated around the target since 2017
Recent Trend in Fishing Intensity or Proxy	Relative exploitation has been below the threshold and declining since 2018
Other Abundance Indices	Tow based series using data from 2007–08 closely resembles the daily series.
Trends in Other Relevant Indicators or Variables	

Projections and Prognosis	
Stock Projections or Prognosis	Without information on recruitment, it is not possible to predict how the stock is going to respond in the next few years.

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Unlikely (< 40%) for current catch Hard Limit: Very Unlikely (< 10%) for current catch Unknown if the catch were to increase to the level of the TACC
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Unlikely (<40%) for current catch Unknown if the catch were to increase to the level of the TACC

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Standardised CPUE	
Assessment Dates	Latest assessment Plenary publication year: 2024	Next assessment: 2027
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	BT.Daily CPUE series	1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
Most of the GUR 2 commercial catch is taken in Hawke Bay, and the index of abundance is naturally weighted to abundance of GUR in this area.


Fishery Interactions
Red gurnard is taken in FMA 2 by the bottom trawl fishery targeting gurnard and tarakihi.

- **GUR 3**

Stock Structure Assumptions
No information is available on the stock separation of red gurnard. The Fishstock GUR 3 is treated in this summary as a unit stock.


Stock Status		
Most Recent Assessment Plenary Publication Year	2022	
Catch in most recent year of assessment	Year: 2020–21	Catch: 1 646 t
Assessment runs presented	Base case model	
Reference Points	Interim Target: 35% B_0 (HSS default) Soft Limit: 20% B_0 (HSS default) Hard Limit: 10% B_0 (HSS default) Overfishing threshold: $F_{SB40\%}$ (HSS default)	
Status in relation to Target	$B_{2020–21}$ was estimated to be 64% B_0; Very Likely (> 90%) to be at or above the target	
Status in relation to Limits	Soft limit: Very Unlikely (< 10%) to be below Hard Limit: Exceptionally Unlikely (< 1%) to be below	
Status in relation to Overfishing	$F_{2020–21}$ was estimated to be at about $F_{SB40\%}$. Overfishing is About as Likely as Not (40–60%) to be occurring	

Historical Abundance and Catch Trajectories




Annual trend in spawning biomass relative to the 35% SB_0 interim target biomass level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The dashed line represents the interim target level. The red and orange dashed lines represent the hard and soft biomass limits, respectively.

Fishing Intensity Trajectory



Annual fishing mortality compared to the $FSB_{40\%}$ interim target fishing mortality level (dashed line) for the base case model (median values from MCMCs).

Combined status



Annual spawning biomass and fishing mortality compared to the $SB_{35\%}$ interim target biomass level and corresponding fishing mortality reference for the base case model (median values from MCMCs). The green dashed lines represent the biomass target and fishing mortality threshold levels. The red and orange dashed lines represent the hard and soft biomass limits, respectively.

Fishery and Stock Trends	
Recent trend in Biomass or Proxy	Spawning biomass was estimated to have increased from the late 2000s following above average recruitment since the mid-2000s.
Recent trend in Fishing Intensity or Proxy	Fishing mortality has been at about the target level since 2010.
Other Abundance Indices	FLA BT CPUE indices have increased considerably from the early 1990s. These CPUE indices were not included in the 2022 assessment. A Southland CPUE series also had generally similar levels of increase.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Abundance is Very Likely (> 90%) to remain above the target biomass level over the next five years at the current TACC or catch.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Very Unlikely (< 10%) Hard Limit: Exceptionally Unlikely (< 1%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	About as Likely as Not (40–60%)

Assessment Methodology and Evaluation		
Assessment Type	Level 1 - Full Quantitative Stock Assessment	
Assessment Method	Age-structured Bayesian stock assessment implemented with Stock Synthesis software and uncertainty estimated by MCMC	
Assessment Dates	Latest assessment Plenary publication year: 2022	Next assessment: 2025
Overall assessment quality rank	1 – High Quality	
Main data inputs	- Commercial catch (from 1985 onwards) - Survey biomass indices. - Survey age/length frequencies. - BT-MIX CPUE indices - Estimates of biological parameters (e.g., growth, natural mortality, age-at-maturity, and length/ weight)	1 – High Quality 1 – High Quality 1 – High Quality 1 – High Quality 1 – High Quality
Data not used (rank)	N/A	
Changes to Model Structure and Assumptions	- First implementation of full quantitative assessment, which includes trawl survey data and a new CPUE series excluding FLA target; previous (2015) assessment was a partial quantitative assessment based on the CPUE series, BT(MIX+FLA)	
Major Sources of Uncertainty	- Choice of the period used to derive reference points, as it is unclear whether recent increases in recruitment represent an increase in productivity - Lack of recent age composition data from trawl survey and fishery - The trawl survey and CPUE indices are for the FMA 3 (ECSI) area only and do not cover the entire GUR 3 QMA; approximately 25% of the GUR 3 catch is taken outside FMA 3	

Qualifying Comments
-

Fishery Interactions
Red gurnard in GUR 3 are taken almost entirely by bottom trawl in fisheries targeted at red cod, barracouta, and flatfish. Some red gurnard are also taken in the target tarakihi and stargazer bottom trawl fisheries. The level of targeting of this species was historically low, averaging less than 10%, but has increased to approximately 25% since 2017–18.


- **GUR 7**

Stock Structure Assumptions
Stock boundaries are unknown, but for the purpose of this summary, GUR 7 is considered to be a single management unit.

Stock Status		
Most Recent Assessment Plenary Publication Year	2022	
Catch in most recent year of assessment	Year: 2020–21	Catch: 1 153 t
Assessment runs presented	Base case	
Reference Points	Interim Target: 35% B_0 (HSS default) Soft Limit: 20% B_0 (HSS default)	


	Hard Limit: 10% B_0 (HSS default) Overfishing threshold: $F_{SB40\%}$ (HSS default)
Status in relation to Target	$B_{2020-21}$ was estimated to be 111% B_0. Virtually certain (> 99%) to be at or above the target
Status in relation to Limits	Soft limit: Exceptionally Unlikely (< 1%) to be below Hard Limit: Exceptionally Unlikely (< 1%) to be below
Status in relation to Overfishing	$F_{2020-21}$ was estimated to be 44% $F_{SB40\%}$. Overfishing is Very Unlikely (< 10%) to be occurring

Historical Abundance and Catch Trajectories




Annual trend in spawning biomass relative to the 35% SB_0 interim target biomass level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The dashed line represents the interim target level. The red and orange dashed lines represent the hard and soft biomass limits, respectively.

Fishing Intensity Trajectories



Annual fishing mortality compared to the $F_{SB40\%}$ interim threshold fishing mortality level (dashed line) for the interim base case model (median values from MCMCs). Grey shaded area represents 95% credible intervals.

Combined trajectories



Annual spawning biomass and fishing mortality compared to the $SB_{35\%}$ interim target biomass level and corresponding fishing mortality threshold for the base case model (median values from MCMCs). The green dashed lines represent the biomass target and fishing mortality threshold levels. The red and orange dashed lines represent the hard and soft biomass limits, respectively. The 95% credible intervals are shown for the final year.

Fishery and Stock Trends	
Recent trend in Biomass or Proxy	Spawning biomass was estimated to have increased from 2010 following above average recruitment since the late 2000s, which has continued to be high over the last 5 years.
Recent trend in Fishing Intensity or Proxy	Fishing mortality has been well below the threshold level since 2010.
Other Abundance Indices	The BT FLA and BT mixed CPUE indices increased from 2009–10 to 2015–16. The CPUE indices were not updated for the 2022 assessment.
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	Abundance is Very Likely (> 90%) to remain above the target biomass level over the next five years at the current catch or TACC.
Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Soft Limit: Exceptionally Unlikely (< 1%) Hard Limit: Exceptionally Unlikely (< 1%)
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (< 10%)

Assessment Methodology and Evaluation		
Assessment Type	Level 1 - Full Quantitative Stock Assessment	
Assessment Method	Age-structured Bayesian stock assessment implemented with Stock Synthesis software and uncertainty estimated by MCMC	
Assessment Dates	Latest assessment Plenary publication year: 2022	Next assessment: 2025
Overall assessment quality rank	1 – High Quality	
Main data inputs	- Commercial catch (from 1987 onwards) - Survey biomass indices - Survey age/length frequencies - Estimates of biological parameters (e.g., growth, natural mortality, age-at-maturity, and length/ weight)	 1 – High Quality 1 – High Quality 1 – High Quality 1 – High Quality
Data not used (rank)	- CPUE indices (to 2015–16)	2 – Medium or Mixed quality: doesn't appear to be indexing abundance throughout the series
Changes to Model Structure and Assumptions	- First implementation of a full quantitative assessment; the previous partial quantitative assessment was based on WCSI trawl survey biomass indices relative to reference period	
Major Sources of Uncertainty	- Lack of recent age composition data from the trawl survey and age or length composition data from the fishery	

Qualifying Comments
- Reference points (SB_0 based) are derived from the average level of recruitment during 1987–2020. There was an apparent shift in the productivity of GUR 7 in the late 2000s. The increasing catches since 2012 were sustained by the higher level of recruitment. - Stock assessment and yield estimates apply only to the reported catch from the commercial fishery.

Fishery Interactions
Red gurnard are primarily taken in conjunction with the following QMS species: flatfish, barracouta, stargazer, red cod, tarakihi, and other species in the west coast South Island target bottom trawl fishery.

6. FOR FURTHER INFORMATION

Beentjes, M P; MacGibbon, D; Lyon, W S (2015) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2014 (KAH1402). *New Zealand Fisheries Assessment Report 2015/14.*

Beentjes, M P; MacGibbon, D J; Escobar-Flores, P (2023) Inshore trawl survey of Canterbury Bight and Pegasus Bay, May–June 2022 (KAH2204). *New Zealand Fisheries Assessment Report 2023/55.* 147 p.

Beentjes, M P; MacGibbon, D J; Ladroit, Y (2022) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2021 (KAH2104). *New Zealand Fisheries Assessment Report 2022/23.* 147 p.

Beentjes, M P; MacGibbon, D J; Parkinson, D (2016) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2016 (KAH1605). *New Zealand Fisheries Assessment Report 2016/61.* 135 p.

Blackwell, R (1988) Red gurnard. New Zealand Fisheries Assessment Research Document 1988/23: 18 p. (Unpublished report held by NIWA library, Wellington.)

Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. (Unpublished New Zealand Fisheries Assessment Research Report for the Ministry of Fisheries Project REC2000-01 held by Fisheries New Zealand, Wellington.) 92 p.

Boyd, R O; Reilly, J L (2004) 1999/2000 National Marine Recreational Fishing Survey: harvest estimates. (Unpublished New Zealand Fisheries Assessment Research Report for the Ministry of Fisheries Project REC9803 held by Fisheries New Zealand, Wellington.) 28 p.

Bradford, E (1998) Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished report held by NIWA library, Wellington.)

Challenger Finfisheries Management Company (2003) Report to the Adaptive Management Programme Fishery Assessment Working Group. GUR 7 Adaptive Management Proposal for the 2004–05 fishing year. (Unpublished report held by Fisheries New Zealand, Wellington.)

Cordue, P L (1998) Designing optimal estimators for fish stock assessment. *Canadian Journal of Fisheries and Aquatic Sciences 55*: 376–386.

Davey, N; Hartill, B; Carter, M (2019) Mean weight estimates for recreational fisheries in 2017–18. *New Zealand Fisheries Assessment Report 2019/25*. 32 p.
Elder, R D (1976) Studies on age and growth, reproduction and population dynamics of red gurnard, *Chelidonichthys kumu* (Lesson and Garnot), in the Hauraki Gulf, New Zealand. *Fisheries Research Bulletin No: 12*. 62 p.
Francis, R I C C (1992) Recommendations concerning the calculation of maximum constant yield (*MCY*) and current annual yield (*CAY*). New Zealand Fisheries Assessment Research Document 1992/8. 26 p. (Unpublished report held by NIWA library, Wellington.)
Francis, R I C C (2011) Data weighting in statistical fisheries stock assessment models. *Canadian Journal of Fisheries and Aquatic Sciences 68*: 1124–1138.
Francis, R I C C; Hurst, R J; Renwick, J A (2001) An evaluation of catchability assumptions in New Zealand stock assessments. *New Zealand Fisheries Assessment Report 2001/1*. 37 p.
Francis, M P; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report 2013/55*. 136 p.
Hartill, B; Bian, R; Armiger, H; Vaughan, M; Rush, N (2007) Recreational marine harvest estimates of snapper, kahawai, and kingfish in QMA 1 in 2004–05. *New Zealand Fisheries Assessment Report 2007/26*. 44 p.
Hartill, B; Bian, R; Rush, N; Armiger, H (2013) Aerial-access recreational harvest estimates for snapper, kahawai, red gurnard, tarakihi and trevally in FMA 1 in 2011–12. *New Zealand Fisheries Assessment Report 2013/70.* 44 p.
Hartill, B; Davey, N (2015) Mean weight estimates for recreational fisheries in 2011–12. *New Zealand Fisheries Assessment Report 2015/25*. 37 p.
Heinemann A; Gray, A. (in prep.) National Panel Survey of Recreational Marine Fishers 2022-23.
Kendrick, T H (2009a) Fishery characterisation and catch-per-unit-effort indices for three sub-stocks of red gurnard in GUR 1; 1989–90 to 2004–05. *New Zealand Fisheries Assessment Report 2009/10*.
Kendrick, T H (2009b) Updated Catch-per-Unit effort indices for red gurnard in GUR 2; 1989–90 to 2004–05 *New Zealand Fisheries Assessment Report 2009/11*.
Kendrick, T H; Bentley, N (2011) Fishery characterisations and catch-per-unit-effort indices for three sub-stocks of red gurnard in GUR 1, 1989–90 to 2008–09. *New Zealand Fisheries Assessment Report 2011/4.*
Kendrick, T H; Bentley, N; Langley, A (2011) Report to the Challenger Fishfish Company: CPUE analyses for FMA 7 Fishstocks of gurnard, tarakihi, blue warehou, and ghost shark. (Unpublished client report held by Trophia Limited, Kaikōura.)
Kendrick, T H; Walker, N (2004) Characterisation of the GUR 2 red gurnard (*Chelidonichthys kumu)* and associated inshore trawl fisheries, 1989–90 to 2000–01. *New Zealand Fisheries Assessment Report 2004/21*. 83 p.
Langley, A (2011) Characterisation of the Inshore Finfish fisheries of Challenger and South East coast regions (FMAs 3, 5, 7 & 8). (Unpublished client report available from http://www.seafoodindustry.co.nz/SIFisheries.)
Langley, A D (2014) Updated CPUE analyses for selected South Island inshore finfish stocks. *New Zealand Fisheries Assessment Report 2014/40.* 116 p.
Langley, A D (2015) Fishery characterisation and Catch-Per-Unit-Effort indices for John dory in JDO 1. *New Zealand Fisheries Assessment Report 2015/47*. 76 p.
Langley, A D (2022a) A stock assessment of red gurnard in GUR 3 for 2022. *New Zealand Fisheries Assessment Report 2022/21*. 56 p.
Langley, A D (2022b) A stock assessment of red gurnard in GUR 7 for 2022. *New Zealand Fisheries Assessment Report 2022/22*. 36 p.
Lydon, G J; Middleton, D A J; Starr, P J (2006) Performance of the GUR 3 Logbook Programme. AMP-WG-06/22. (Unpublished manuscript available from the New Zealand Seafood Industry Council, Wellington.)
Lyon, W S; Horn, P L (2011) Length and age of red gurnard (*Chelidonichthys kumu*) from trawl surveys off west coast South Island in 2003, 2005, and 2007, with comparisons to earlier surveys in the time series. *New Zealand Fisheries Assessment Report 2011/46.*
MacGibbon, D J (2019) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March-April 2019 (KAH1902) *New Zealand Fisheries Assessment Report 2019/64*. 87 p.
MacGibbon, D J; Beentjes, M P; Lyon, W; Ladroit, Y (2019) Inshore trawl survey of Canterbury Bight and Pegasus Bay, April–June 2018 (KAH1803). *New Zealand Fisheries Assessment Report 2019/03.* 136 p.
MacGibbon, D J; Stevenson, M L (2013) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March-April 2013 (KAH1305). *New Zealand Fisheries Assessment Report 2013/66.* 115 p.
MacGibbon, D J; Walsh, C; Buckthought, D; Bian, R (2022) Inshore trawl survey off the west coast South Island and in Tasman Bay and Golden Bay, March–April 2021 (KAH2103). *New Zealand Fisheries Assessment Report 2022/11*. 97 p.
MacGibbon, D J; Walsh, C; Buckthought, D; Bian, R (2024) Inshore trawl survey off the west coast South Island and in Tasman Bay and Golden Bay, March–April 2023 (KAH2302). *New Zealand Fisheries Assessment Report 2024/06*. 91 p.
McKenzie, A; Moore, B R (2023) Fishery characterisation and catch per unit effort for red gurnard in GUR 1 to 2020/21, with an exploration of stock structure in New Zealand waters. *New Zealand Fisheries Assessment Report 2023/16*. 75 p.
Methot, R D; Wetzel, C R (2013) Stock synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research 142 (2013)*: 86–99.
Middleton, D A J; Neubauer, P; Thompson, F N (in prep.) Characterisation and CPUE for the red gurnard fishery in GUR 2 from 1989 to 2023. New Zealand Fisheries Assessment Report 2024/xx.
Morrison, M A; Francis, M P; Parkinson, D M (2002) Trawl survey of the Hauraki Gulf, 2000 (KAH0012). *New Zealand Fisheries Assessment Report 2002/46*. 48 p.
Parker, S; Fu, D (2012) Length and age structure of commercial landings of red gurnard (*Chelidonichthys kumu*) in GUR 2 in 2009–10. *New Zealand Fisheries Assessment Report 2012/35*. 36 p.
Schofield, M I; Langley, A D; Middleton, D A J (2018a) Fisheries characterisation and catch-per unit-effort analyses GUR 2. *New Zealand Fisheries Assessment Report 2018/10*. 53 p.
Schofield, M I; Langley, A D; Middleton, D A J (2018b) Catch-per unit-effort (CPUE) update for FMA 2 gurnard (GUR 2). (Report for Fisheries Inshore New Zealand.) https://www.inshore.co.nz/fileadmin/Documents/Science/GUR2_rapidCPUEupdate_2018.pdf
Schofield, M I; Langley, A D; Middleton, D A J (2018c) Catch-per unit-effort (CPUE) update for FMA 3 gurnard (GUR 3). Report for Southern Inshore Fisheries Management Company.
Starr, P J; Kendrick, T H (2013) GUR 3 Fishery Characterisation and CPUE. *New Zealand Fisheries Assessment Report 2013/37*. 71 p.
Starr, P J; Kendrick, T H (2017) GUR 7 Fishery Characterisation and CPUE Report. *New Zealand Fisheries Assessment Report 2017/49*. 144 p.
Starr, P J; Kendrick, T H; Lydon, G J; Bentley, N (2007) Report to the Adaptive Management Fishery Assessment Working Group: Full term review of the GUR 3 Adaptive Management Programme. AMP-WG-07/11v2. (Unpublished manuscript available from the New Zealand Seafood Industry Council, Wellington.)
Stevenson, M L (2000) Assessment of red gurnard (*Chelidonichthys kumu*) stocks GUR 1 and GUR 2. *New Zealand Fisheries Assessment Report 2000/40*. 51 p.
Stevenson, M L (2004) Trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2003 (KAH0304). *New Zealand Fisheries Assessment Report 2004/4*.
Stevenson, M L (2006) Trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2005 (KAH0503). *New Zealand Fisheries Assessment Report 2006/4*.

Stevenson, M L (2007) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2007 KAH0704. *New Zealand Fisheries Assessment Report 2007/41*.

Stevenson, M L (2009) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2009. *New Zealand Fisheries Assessment Report 2010/11*.

Stevenson, M L (2012) Inshore trawl survey of the west coast of the South Island and Tasman and Golden Bays, March–April 2011. *New Zealand Fisheries Assessment Report 2012/50*. 77 p.

Stevenson, M L; MacGibbon, D J (2015) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March-April 2015 (KAH1503). *New Zealand Fisheries Assessment Report 2015/67*. 94 p.

Sutton, C P (1997) Growth parameters, and estimates of mortality for red gurnard (Chelidonichthys kumu) from off the east and west coasts of the South Island, New Zealand. New Zealand Fisheries Assessment Research Document 1997/1. 15 p. (Unpublished report held by NIWA library, Wellington.)

Teirney, L D; Kilner, A R; Millar, R E; Bradford, E; Bell, J D (1997) Estimation of recreational catch from 1991–92 to 1993–94 New Zealand Fisheries Assessment Research Document 1997/15. 43 p. (Unpublished report held by NIWA library, Wellington.)

Vignaux, M (1997) CPUE analyses for fishstocks in the adaptive management programme. New Zealand Fisheries Assessment Research Document 1997/24. 68 p. (Unpublished report held by NIWA library, Wellington.)

Wright, P; McClary, D; Boyd, R O (2004) 2000/2001 National Marine Recreational Fishing Survey: direct questioning of fishers compared with reported diary data. (Final Research Report for Ministry of Fisheries Project REC2000–01: Objective 2 held by Fisheries New Zealand.)

Wynne-Jones, J; Gray, A; Heinemann, A; Hill, L; Walton, L (2019) National Panel Survey of Marine Recreational Fishers 2017–2018. *New Zealand Fisheries Assessment Report 2019/24*. 104 p.

Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.