


SPRAT (SPR)

(*Sprattus antipodum*, *S. muelleri*)
Kupae



1. FISHERY SUMMARY

There are two species of sprats in New Zealand, *Sprattus antipodum* (slender sprat) and *S. muelleri* (stout sprat). They can be distinguished by body shape, colour, and some morphological features, but since they are very similar it is impractical to separate them in large catches.

Sprats were introduced into the QMS on 1 October 2002, with the allowances, TACCs and TACs shown in Table 1, which have not been changed since.

Table 1: Recreational and customary non-commercial allowances, TACCs and TACs for sprats by Fishstock.

Fishstock	Recreational Allowance	Customary non-commercial		Other mortality	TACC	TAC
		Allowance				
SPR 1	20	10		0	70	100
SPR 3	10	5		0	285	300
SPR 4	3	2		0	10	15
SPR 7	10	5		0	85	100
SPR 10	0	0		0	0	0

1.1 Commercial fisheries

The sprat “fishery” is minor and intermittent. There is no information on catches or landings of sprats prior to 1990, although occasional catches were made during exploratory fishing projects on small pelagic species, mainly in the 1960s and 1970s. Sprats have undoubtedly been caught in most years, but were either not reported, reported as “bait”, or included in the category “mixed species”. The name “sprat” is used in a general sense for several unrelated small fishes, and the juveniles of some larger species. This may have introduced errors into catch records. Reported landings since 1990 have ranged from less than 1 t to 7 t (Table 2); no landings were reported in 2017–18 and 2019–20, and < 1 t has been landed annually since 2004–05. The most consistent (but small) catches have been by bottom trawl. Reported landings by setnet and beach seine could be of true sprats, but may also be of yellow-eyed mullet (*Aldrichetta forsteri*), known colloquially as sprats. This is particularly likely in the upper North Island where the presence of sprats is considerably reduced or non-existent.

Table 2: Reported landings (t) of sprat by Fishstock and fishing year. No catches reported for SPR 10, which has a TACC of 0.

FMA	SPR 1		SPR 3		SPR 4		SPR 7		Total	
	1, 2, 8 & 9		3, 5 & 6		4		7		Landings	TACC
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1990–91†	3	-	<1	-	0	-	<1	-	3	-
1991–92†	1	-	0	-	0	-	0	-	1	-
1992–93†	<1	-	<1	-	0	-	0	-	<1	-
1993–94†	<1	-	<1	-	0	-	<1	-	1	-
1994–95†	<1	-	<1	-	0	-	<1	-	1	-
1995–96†	<1	-	6	-	0	-	<1	-	7	-
1996–97†	<1	-	1	-	0	-	<1	-	1	-
1997–98†	<1	-	<1	-	0	-	<1	-	<1	-
1998–99†	2	-	<1	-	0	-	<1	-	4	-
1999–00†	<1	-	<1	-	0	-	1	-	2	-
2000–01†	<1	-	<1	-	0	-	<1	-	<1	-
2001–02	<1	-	<1	-	0	-	<1	-	<1	-
2002–03	<1	70	<1	285	0	10	0	85	<1	450
2003–04	<1	70	3	285	0	10	0	85	3	450
2004–05	<1	70	0	285	0	10	0	85	<1	450
2005–06	<1	70	0	285	0	10	0	85	<1	450
2006–07	<1	70	<1	285	0	10	0	85	<1	450
2007–08	<1	70	0	285	0	10	0	85	<1	450
2008–09	<1	70	<1	285	0	10	<1	85	1	450
2009–10	<1	70	0	285	0	10	0	85	0	450
2010–11	<1	70	0	285	0	10	0	85	<1	450
2011–12	<1	70	0	285	0	10	0	85	<1	450
2012–13	<1	70	<1	285	0	10	<1	85	<1	450
2013–14	<1	70	0	285	<1	10	0	85	<1	450
2014–15	<1	70	<1	285	0	10	<1	85	<1	450
2015–16	<1	70	0	285	0	10	0	85	<1	450
2016–17	0	70	<1	285	0	10	<1	85	<1	450
2017–18	0	70	0	285	0	10	0	85	0	450
2018–19	<1	70	0	285	0	10	0	85	<1	450
2019–20	0	70	0	285	0	10	0	85	0	450
2020–21	0	70	<1	285	0	10	0	85	<1	450
2021–22	<1	70	0	285	0	10	0	85	<1	450
2022–23	0	70	0	285	<1	10	<1	85	<1	450

† CELR

1.2 Recreational fisheries

There is no known recreational fishery, but small numbers are caught in small-mesh setnets and beach seines.

1.3 Customary non-commercial fisheries

Quantitative information on the current level of customary non-commercial take is not available.

1.4 Illegal catch

Estimates of illegal catch are not available, but are probably insignificant or nil.

1.5 Other sources of mortality

Some accidental captures of sprats by vessels purse seining for other small pelagic species may be discarded if no market is available.

2. BIOLOGY

Sprats occur in coastal waters from the Bay of Islands to Stewart Island, and are present at the Auckland Islands. It is not known whether the two species have different distributions. Sprats appear to be most abundant off the southeastern coast of the South Island, where anchovies are absent. Their vertical distribution within the water column is not known.

Spawning occurs in areas of reduced salinity when water temperatures are coolest 9–10.5 °C; there are consequently regional differences in spawning season with spawning peaks occurring between June and November (Taylor & Marriott 2004). The eggs are pelagic.

No reliable ageing work has been undertaken. Sprats are assumed to feed on zooplankton, and are preyed upon by larger fishes, seabirds, and marine mammals.

There have been no biological studies that are directly relevant to the recognition of separate stocks, or to yield estimates. Consequently no estimates of biological parameters are available. There is an extensive international literature base on sprats, mainly *Sprattus sprattus*, but the relevance of this to the New Zealand species is unknown.

3. STOCKS AND AREAS

There is no biological information on which to make an assessment on whether separate stocks exist. However, there are two species, and their relative distributions are unknown. As presently understood, both species are more common around southern New Zealand. If their distributions do differ, and the biomass of each species fluctuates independently, there are unknown implications for localised stock depletion.

4. STOCK ASSESSMENT

There have been no previous stock assessments of sprats. There have been two very general estimates of biomass in the Canterbury Bight region: 50 000 t (Robertson 1978), and 60 000 t (Colman 1979), with a possible yield of 10 000 t. No information on biomass variability is available.

4.1 Estimates of fishery parameters and abundance

No fishery parameters are available.

4.2 Biomass estimates

No estimates of biomass (B_0 , B_{MSY} , or $B_{CURRENT}$) are available.

4.3 Yield estimates and projections

Estimation of Maximum Constant Yield (MCY)

MCY cannot be determined.

Estimation of Current Annual Yield (CAY)

Current biomass cannot be estimated, so CAY cannot be determined.

Other yield estimates and stock assessment results

No information is available.

4.4 Other factors

Data from some ichthyoplankton surveys show one or both sprat species to be locally abundant. However, it is unlikely that the biomass is comparable to the very large stocks in the northern hemisphere where there are large sprat fisheries.

It is not known whether the biomass of sprats is stable or variable, but the latter is considered more likely.

In some localities around the South Island, sprats are a major food source for many fishes, seabirds, and marine mammals. Excessive localised harvesting may disrupt ecosystems.

5. STATUS OF THE STOCKS

No estimates of current biomass are available. At the present level of minimal catches, stocks are at or close to their natural level. This is nominally a virgin biomass, but not necessarily a stable one.

6. FOR FURTHER INFORMATION

- Baker, A N (1973) Spawning and development of the New Zealand sprat, *Sprattus antipodum* (Hector). Zoology Publications from Victoria University of Wellington No. 62. 12 p.
- Colman, J A (1979) Spawning of the sprat, *Sprattus antipodum* (Hector), round the South Island of New Zealand. *New Zealand Journal of Marine and Freshwater Research* 13(2): 263–272.
- Fenaughty, J M; Bagley, N W (1981) *W.J. Scott* New Zealand trawling survey: South Island east coast. *Fisheries Technical Report* No. 157. 224 p.
- Morgans, J F C (1966) Possibilities raised by a study of the size distribution in a sample of a shoal of sprats, *Sprattus antipodum* (Hector). *Transactions of the Royal Society of New Zealand, Zoology* 8(13): 141–147.
- Robertson, D A (1978) Blue mackerel, pilchard, anchovy, sprat, saury, and lanternfish. In Habib, G; Roberts, P E (Comps.) Proceedings of the Pelagic Fisheries Conference July 1977. pp. 85–89. Fisheries Research Division Occasional Publication No. 15.
- Smith, P J; Robertson, D A (1981) Genetic evidence for two species of sprat (*Sprattus*) in New Zealand waters. *Marine Biology* 62(4): 227–233.
- Taylor, P.R., Marriot, P.M. (2004) A summary of information on spawning of the small inshore pelagic species, anchovy (*Engraulis australis*), garfish (*Hyporhamphus ihi*), pilchard (*Sardinops sagax*), and sprat (*Sprattus antipodum* and *S. muelleri*), with a series of stock boundaries proposed for future testing. Draft New Zealand Fisheries Assessment Report 2004/xx. 33 p.
- Whitehead, P J P; Smith, P J; Robertson, D A (1985) The two species of sprat in New Zealand waters (*Sprattus antipodum* and *S. muelleri*). *New Zealand Journal of Marine and Freshwater Research* 19(2): 261–271.