


LARGE TROUGH SHELL (MMI)*(Mactra murchisoni)***1. FISHERY SUMMARY**

This species is part of the surf clam fishery and the reader is guided to the Introduction – surf clams chapter for information common to all relevant species.

1.1 Commercial fisheries

Large trough shells (*Mactra murchisoni*) were introduced into the Quota Management System on 1 April 2004 with a total TACC of 162 t. No allowances were initially made for customary, recreational, or other sources of mortality; some allowances were introduced for MMI 8 and 7 in 2013 and 2016, respectively. Biomass surveys in QMA 3 supported a TACC increase from April 2010. This increased the TACC for MMI 3 from 3 t to 62 t. A subsequent biomass survey in 2012 supported a TAC increase in MMI 8 from 25 t to 631 t in April 2013. Another biomass survey supported a TAC increase in MMI 7 from 61 t to 144 t in April 2016. The current total TAC is 872 t (Table 1).

Table 1: Recreational and Customary non-commercial allowances, other mortality, TACCs, and TACs (t) for large trough shell (*Mactra murchisoni*) by Fishstock.

Fishstock	Description	Recreational allowance	Customary non-commercial allowance	Other sources of mortality	TACC	TAC
MMI 1	Auckland	–	–	–	2	2
MMI 2	Central (East)	–	–	–	3	3
MMI 3	South East (Coast)	0	0	3	62	65
MMI 4	South East (Chatham Rise)	–	–	–	1	1
MMI 5	Southland	–	–	–	1	1
MMI 7	Challenger	1	5	7	131	144
MMI 8	Central (West)	0	10	32	589	631
MMI 9	Auckland (West)	–	–	–	25	25

All reported landings have been from MMI 3, MMI 7 and MMI 8. Between the 1991–92 and 1995–96 fishing years landings were small and confined to MMI 7. No further landings were reported until 2002–03. Since then the reported total landings have ranged between about 23 t and 99.6 t (Table 2).

MMI 3 landings reached the TACC in 2013–14, and again since 2019–20, but decreased to levels well below the TACC in the intervening years. MMI 7 landings were close to the TACC from 2004–05 to 2006–07 but have fluctuated around a lower level since this time; the TACC was increased in 2015. Figure 1 shows the historical landings and TACCs for the two main MMI stocks.

Table 2: TACCs and reported landings (t) of large trough shell by Fishstock from 1991–92 to present from CELR and CLR data. Fishstocks where no catch has been reported are not tabulated, but Total Landings and TACC correspond to all SAE stocks. See Table 1 for TACC of stocks not landed. The fishing year is from 1 April to 31 March.

Year	MMI 3		MMI 7		MMI 8		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1991–92	0	0	0.35	–	–	–	0.35	–
1992–93	0	0	1.54	–	–	–	1.54	–
1993–94	0	0	8.33	–	–	–	8.33	–
1994–95	0	0	10.43	–	–	–	10.43	–
1995–96	0	0	0.14	–	–	–	0.14	–
1996–97	0	0	0	–	–	–	0	–
1997–98	0	0	0	–	–	–	0	–
1998–99	0	0	0	–	–	–	0	–
1999–00	0	0	0	–	–	–	0	–
2000–01	0	0	0	–	–	–	0	–
2001–02	0	0	0	–	–	–	0	–
2002–03	0	0	22.62	–	–	–	22.62	–
2003–04	0	44	29.68	61	–	–	29.68	162
2004–05	0	44	60.02	61	0	25	60.86*	162
2005–06	0	44	53.96	61	0	25	57.92*	162
2006–07	7.48	44	54.09	61	0	25	61.57	162
2007–08	36.90	44	15.04	61	0	25	51.94	162
2008–09	32.15	44	6.66	61	0	25	38.81	162
2009–10	25.76	44	3.42	61	0	25	29.18	162
2010–11	12.60	62	17.43	61	0	25	30.03	180
2011–12	0	62	47.34	61	0	25	47.34	180
2012–13	44.45	62	32.81	61	0	25	77.27	180
2013–14	63.87	62	4.89	61	0	589	68.75	744
2014–15	59.00	62	9.69	61	0	589	68.64	744
2015–16	46.72	62	23.98	131	1.08	589	71.77	814
2016–17	35.79	62	25.62	131	1.18	589	62.59	814
2017–18	40.39	62	29.43	131	2.05	589	71.87	814
2018–19	29.92	62	32.43	131	0.57	589	62.92	814
2019–20	62.91	62	36.12	131	0.59	589	99.62	814
2020–21	63.10	62	16.53	131	0.33	589	79.96	814
2021–22	62.07	62	17.98	131	5.02	589	85.07	814
2022–23	59.14	62	9.40	131	2.36	589	70.90	814
2023–24	57.44	62	1.29	131	0.60	589	59.32	814

*In 2004–05 and 2005–06, 0.84 and 3.9554 t respectively were reportedly landed, but the QMA was not recorded. These amounts are included in the total landings for these years.

1.2 Recreational fisheries

Offshore clams such as *M. murchisoni* are likely to have been harvested for recreational use only when washed ashore after storms. There are no estimates of recreational take for this surf clam.

1.3 Customary fisheries

Offshore clams such as *M. murchisoni* are likely to have been harvested for customary use only when washed ashore after storms. Shells of this clam have been found irregularly, and in small numbers, in a few middens (Conroy et al 1993). There are no estimates of current customary catch of this clam.

1.4 Illegal catch

There is no documented illegal catch of this clam.

1.5 Other sources of mortality

There is no quantitative information on other sources of mortality, although this clam is subject to localised catastrophic mortality from erosion during storms, high temperatures and low oxygen levels during calm summer periods, blooms of toxic algae, and excessive freshwater outflow (Cranfield & Michael 2001).




Figure 1: Reported commercial landings and TACC for MMI 3 (South East Coast) (top) and MMI 7 (Challenger) (bottom). Note that these figures do not show data prior to entry into the QMS.

2. BIOLOGY

M. murchisoni is most abundant around the lower half of the North Island and the South Island. It is found most commonly between about 4 m and 8 m in depth. Maximum length is variable between areas, ranging from 63 mm to 102 mm (Cranfield et al 1993). The sexes are separate, they are broadcast spawners, and the larvae are thought to be planktonic for between 20 and 30 days (Cranfield & Michael 2001). Recruitment of spat is to the same depth zone that adults occur in, although recruitment between years is highly variable (Conroy et al 1993).

3. STOCKS AND AREAS

For management purposes stock boundaries are based on FMAs, however the boundaries of stocks of surf clams are likely to be the continuous lengths of exposed sandy beaches between geographical features (rivers, headlands, etc). Circulation patterns may isolate surf clams genetically as well as ecologically.

4. ENVIRONMENTAL AND ECOSYSTEM CONSIDERATIONS

See the Introduction – surf clams chapter.

5. STOCK ASSESSMENT

5.1 Estimates of fishery parameters and abundance

No estimates of fisheries parameters or abundance are available for this species.

5.2 Biomass estimates

Biomass has been estimated from MMI 2, 3, 7, and 8 at various times between 1994 and 2015 with stratified random surveying using a hydraulic dredge. Survey size has been expressed either as length of beach (Table 3), or as area (Table 4), which makes comparisons difficult.

In both 2012 (FMA 8) and 2015 (Cloudy Bay, FMA 7), White et al (2012, 2015) have conducted a 2-phase stratified random sampling survey. The survey area was stratified by 4 depth strata (0–2 m, 2–4 m, 4–6 m, and 6–8 m, each with respect to Chart Datum). Each station comprised a ~50 m tow, sampling ~80 m² of seabed. All commercial species of subtidal surf clams caught were sorted by species. The total weight of each of these species was measured on board. Individuals from each species were collected and measured for shell length along the anterior-posterior axis (to the nearest millimetre). For tows with less than ~500 individuals, the maximum of either 20 individuals or 20% of the total was measured. For tows with higher than ~500 individuals, 10% with an upper limit of ~200 individuals per tow were measured. To subsample large catches and to avoid issues of size sorting inside the dredge, each of the bins was subsampled by tipping one bin into two bins and repeating until the requisite sub sample size was reached. The number and weight of the main bycatch species was also recorded. Both the biomass densities and biomass estimates were calculated for all the commercial species of subtidal surf clams caught.

Table 3: A summary of biomass estimates in tonnes greenweight (with standard deviation in parentheses) from exploratory surveys of Cloudy Bay (Cranfield et al 1994a) and Clifford Bay in Marlborough (Michael et al 1994), and Foxton beach on the Manawatu coast (White et al 2012).

Area	Cloudy Bay (MMI 7)	Clifford Bay (MMI 7)	Foxton Beach (MMI 8)
Length of beach (km)	11	21	46 [#]
Biomass (t)	248 (96)	192 (79)	3 603 (342) [#]

[#] Biomass was estimated at Foxton Beach from a mix of a systematic survey to the north and a stratified survey to the south of this location.

Table 4: A summary of biomass estimates in greenweight (t) from the surveys in MMI 2 (Triantifillos 2008b), MMI 3 (Triantifillos 2008a), and MMI 7 (White et al 2015). Note: unless otherwise stated the CV is less than 20%.

Location	Five sites (MMI 2)	Ashley River to 6 nm south of the Waimakariri River (MMI 3)	Cloudy Bay (MMI 7)
Area surveyed (km ²)	28.0	13.4	5.7
Biomass (t)	33.8	444.1	1 008.8

5.3 Yield estimates and projections

Growth and mortality data from Cloudy Bay in Marlborough and the Kapiti Coast in Manawatu (Cranfield et al 1993) have been used in a yield per recruit model to estimate the reference fishing mortality $F_{0.1}$ (Cranfield et al 1994a, Triantifillos 2008a, 2008b). The Shellfish Working Group (SFWG) did not accept these estimates of $F_{0.1}$ because there was considerable uncertainty in both the estimates and the method used to generate them. The MCY estimates of Triantifillos (2008a, 2008b) and White et al (2012) using the full range of $F_{0.1}$ estimates from Cranfield et al (1993) are shown in Table 5. The SFWG recommended that MCY estimates are adequate to use to inform management decisions relevant to all surf clam fisheries, with the following caveats: 1) due to the uncertainty in $F_{0.1}$ values, for all species other than SAE, the MCY estimates should use the $F_{0.1}$ values toward the higher end of the range, and 2) there is a need to account for any substantial catch that has already come out of any surf clam fishery when estimating MCY ; however there was no consensus on the best way to do this.

Estimates of MCY are available from numerous locations (Table 5) and were calculated using Method 1 for a virgin fishery (MPI 2015) with an estimate of virgin biomass B_0 , where:

$$MCY = 0.25 * F_{0.1} B_0$$

Table 5: *MCY* estimates (t) for *M. murchisoni* from virgin biomass at locations sampled around New Zealand (Triantifillos 2008a, 2008b, White et al 2012). The two $F_{0.1}$ values, which are subsequently used to estimate *MCY*, are the minimum and maximum estimates from Cranfield et al (1993).

Location	$F_{0.1}$	<i>MCY</i>
Five sites (MMI 2)	0.43/0.57	47.7/63.3
Ashley River to 6 nm south of the Waimakariri River (MMI 3)	0.70/0.89	5.9/7.5
Cloudy Bay (MMI 7)	0.43/0.57	108.4/143.7
46 km of coast north and south of the Manawatu River (MMI 8)	0.70/0.89	630.6/801.7

Estimation of Current Annual Yield (CAY)

CAY has not been estimated for *M. murchisoni*.

The SFWG recommended moving all surf clam fisheries away from an *MCY* management strategy and towards an exploitation rate management strategy. The SFWG recognised that an exploitation rate approach is more survey intensive, but better allows for the variable nature of biomass for surf clams because it allows greater flexibility in catch (to take greater landings from available biomass) whilst keeping catches sustainable.

6. STATUS OF THE STOCKS

• MMI 3- South East (Coast)

The most recent survey conducted for MMI 3 was in 2008. There is no longer adequate information to inform current stock status which is therefore Unknown.

• MMI 7 – Marlborough Sounds

Stock Status		
Most Recent Assessment Plenary Publication Year	2015 – now considered out of date	
Intrinsic Productivity Level	Medium	
Catch in most recent year of assessment	Year: 2013–14	Catch: 5 t
Assessment Runs Presented	Survey biomass	
Reference Points	Target: Not defined, but B_{MSY} assumed Soft Limit: 20% B_0 Hard Limit: 10% B_0 Overfishing threshold: -	
Status in relation to Target	Unknown	
Status in relation to Limits	Unknown	
Status in relation to Overfishing	Unknown	

Historical Stock Status Trajectory and Current Status
Unknown

Fishery and Stock Trends	
Recent Trend in Biomass or Proxy	Unknown
Recent Trend in Fishing Mortality or Proxy	Landings have been variable but averaged 26.1 t since 2002.
Other Abundance Indices	-
Trends in Other Relevant Indicators or Variables	-

Projections and Prognosis	
Stock Projections or Prognosis	-

Probability of Current Catch or TACC causing Biomass to remain below or to decline below Limits	Current catches are Very Unlikely (< 10%) to cause declines below soft or hard limits in the short to medium term.
Probability of Current Catch or TACC causing Overfishing to continue or to commence	Very Unlikely (< 10%)

Assessment Methodology and Evaluation		
Assessment Type	Level 2 - Partial Quantitative Stock Assessment	
Assessment Method	Absolute biomass estimates from quadrat surveys	
Assessment Dates	Last assessment: 2015	Next assessment: Unknown
Overall assessment quality rank		
Main data inputs (rank)	Abundance and length frequency information	
Data not used (rank)	-	
Changes to Model Structure and Assumptions	-	
Major Sources of Uncertainty	-	

Qualifying Comments
Stock size could fluctuate markedly as a result of catastrophic mortality from a number of causes. There is a need to review fishery parameters for this species.

Fishery Interactions
MMI can be caught together with other surf clam species and non-QMS bivalves.

• MMI 8 – Manawatu coast

The most recent survey conducted for MMI 8 was in 2008. There is no longer adequate information to inform current stock status which is therefore Unknown.

For all other MMI stocks there is insufficient information to estimate current stock status.

7. FOR FURTHER INFORMATION

- Beentjes, M P; Baird, S J (2004) Review of dredge fishing technologies and practice for application in New Zealand. *New Zealand Fisheries Assessment Report 2004/37*. 40 p.
- Brierley, P (Convenor) (1990) Management and development of the New Zealand sub-tidal clam fishery. Report of the surf clam working group, MAF Fisheries. (Unpublished report held in NIWA library, Wellington). 57 p.
- Conroy, A; Smith, P; Michael, K; Stotter, D (1993) Identification and recruitment patterns of juvenile surf clams, *Macra discors* and *M. murchisoni* from central New Zealand. *New Zealand Journal of Marine and Freshwater Research* 27: 279–285.
- Cranfield, H J; Doonan, I J; Michael, K P (1994a) Dredge survey of surf clams in Cloudy Bay, Marlborough. *New Zealand Fisheries Technical Report* 39. 18 p.
- Cranfield, H J; Michael, K P (2001) The surf clam fishery in New Zealand: description of the fishery, its management, and the biology of surf clams. *New Zealand Fisheries Assessment Report 2001/62*. 24 p.
- Cranfield, H J; Michael, K P; Stotter, D R (1993) Estimates of growth, mortality, and yield per recruit for New Zealand surf clams. New Zealand Fisheries Assessment Research Document 1993/20. 26 p. (Unpublished report held by NIWA library, Wellington).
- Cranfield, H J; Michael, K P; Stotter, D R; Doonan, I J (1994b) Distribution, biomass and yield estimates of surf clams off New Zealand beaches. New Zealand Fisheries Assessment Research Document 1994/1. 17 p. (Unpublished document held by NIWA library, Wellington).
- Haddon, M; Willis, T J; Wear, R G; Anderlini, V C (1996) Biomass and distribution of five species of surf clam off an exposed west coast North Island beach, New Zealand. *Journal of Shellfish Research* 15: 331–339.
- Michael, K; Cranfield, H; Doonan, I; Hadfield, J (1994) Dredge survey of surf clams in Clifford Bay, Marlborough. *New Zealand Fisheries Data Report No. 54*.
- Ministry for Primary Industries (2015) Fisheries Assessment Plenary, May 2015: stock assessments and stock status. Compiled by the Fisheries Science Group, Ministry for Primary Industries, Wellington, New Zealand. 1475 p.
- Triantifillos, L (2008a) Survey of subtidal surf clams in Pegasus Bay, November–December 2007. Prepared by NIWA for Seafood Innovations Limited and SurfCo. Limited. 43 p.
- Triantifillos, L (2008b) Survey of subtidal surf clams in Quota Management Area 2, June–August 2008. Prepared by NIWA for Seafood Innovations Limited and SurfCo. Limited. 40 p.
- White, W; Millar, R; Breen, B; Farrington, G (2012) Survey of subtidal surf clams from the Manawatu Coast (FMA 8), October–November 2012. (Unpublished Report held by Fisheries New Zealand Wellington.) 35 p. + Addendum.

White, W; Millar, R; Farrington, G; Breen, D; Selveraj, S (2015) Stock assessment of surf clams from Cloudy Bay, NZ. *Institute for Applied Ecology New Zealand Report 15/01*. Published by Applied Ecology New Zealand, an Institute of Auckland University of Technology. 34 p.