

**SNAPPER (SNA 7)***(Chrysophrys auratus)*

Tamure, Kouarea


**1. FISHERIES SUMMARY****1.1 Commercial fisheries**

Table 1 and Table 2 provide a summary by fishing year of the reported commercial catches, TACCs, and TACs for SNA 7. Landings and TACC are plotted in Figure 1.

**Table 1: Reported landings (t) of snapper from SNA 7 from 1931 to 1990.**

Year	SNA 7	Year	SNA 7
1931–32	69	1961	583
1932–33	36	1962	582
1933–34	65	1963	569
1934–35	7	1964	574
1935–36	10	1965	780
1936–37	194	1966	1 356
1937–38	188	1967	1 613
1938–39	149	1968	1 037
1939–40	158	1969	549
1940–41	174	1970	626
1941–42	128	1971	640
1942–43	65	1972	767
1943–44	29	1973	1 258
1944	96	1974	1 026
1945	118	1975	789
1946	232	1976	1 040
1947	475	1977	714
1948	544	1978	2 720
1949	477	1979	1 776
1950	514	1980	732
1951	574	1981	592
1952	563	1982	591
1953	474	1983	544
1954	391	1984	340
1955	504	1985	270
1956	822	1986	253
1957	1 055	1987	210
1958	721	1988	193
1959	650	1989	292
1960	573	1990	200

Notes:

1. The 1931–1943 years are April–March but from 1944 onwards are calendar years.
2. The ‘QMA totals’ are approximations derived from port landing subtotals, as follows: SNA 7, Marlborough Sounds ports to Greymouth
3. Before 1946 the ‘QMA’ subtotals sum to less than the New Zealand total because data from the complete set of ports are not available.
4. Data up to 1985 are from fishing returns: data from 1986 to 1990 are from Quota Management Reports.
5. Data for the period 1931 to 1982 are based on reported landings by harbour and are likely to be underestimated as a result of under-reporting and discarding practices. Data include both foreign and domestic landings.

**Table 2: Reported landings (t) of snapper from SNA 7 from 1983–84 to present and gazetted and actual TACCs (t) for 1986–87 to present. QMS data from 1986–present.**

Fishstock FMAs	SNA 7	
	Landings	TACC
1983–84†	375	–
1984–85†	255	–
1985–86†	188	–
1986–87	257	330
1987–88	256	363
1988–89	176	372
1989–90	294	151
1990–91	160	160
1991–92	148	160
1992–93	165	160
1993–94	147	160
1994–95	150	160
1995–96	146	160
1996–97	162	160
1997–98	182	200
1998–99	142	200
1999–00	174	200
2000–01	156	200
2001–02	141	200
2002–03	187	200
2003–04	215	200
2004–05	178	200
2005–06	166	200
2006–07	248	200
2007–08	187	200
2008–09	205	200
2009–10	188	200
2010–11	206	200
2011–12	216	200
2012–13	211	200
2013–14	210	200
2014–15	210	200
2015–16	189	200
2016–17	263	250
2017–18	263	250
2018–19	257	250
2019–20	289	250
2020–21	337	350
2021–22	361	350
2022–23	518	450
2023–24	475	450
2024–25		720

† FSU data. SNA 7 = Statistical Areas 017, 033–036, 038

The SNA 7 TACC was increased to 350 t in 2020–21, to 450 t in 2022–23 and to 720 t in 2025–25 (Table 2). All commercial fisheries have a minimum legal size (MLS) for snapper of 25 cm.

**Table 3: Recreational and Customary non-commercial allowances, other mortality, TACCs, and TACs (t) for SNA 7 from October 2024.**

Fishstock	Recreational allowance	Customary non-commercial allowance	Other sources of mortality	TACC	TAC
SNA 7	275	60	61	720	1 116

### Foreign fishing

Japanese catch records and observations made by New Zealand naval vessels indicate that significant quantities of snapper were taken from New Zealand waters by Japanese vessels from the late 1950s until 1977. There are insufficient data to quantify historical Japanese catch tonnages for the respective snapper stocks. However, trawl catches have been reported by area from 1967 to 1977, and longline catches from 1975 to 1977 (Table 4). These data were supplied to the Fisheries Research Division of MAF in the late 1970s; however, the data series is incomplete, particularly for longline catches.


A substantial snapper catch was taken by the Japanese trawl fleet operating in the southern area of SNA 8 (South Taranaki Bight) adjacent to the northern SNA 7 boundary.

**Table 4: Reported landings (t) of snapper and harvest within SNA 7 from 1967 to 1977 by Japanese trawl and longline fisheries.**

Year	(a) Trawl	Trawl catch (all species)	Total snapper trawl catch	SNA 7
1967		3092	30	NA
1968		19 721	562	17
1969		25 997	1 289	251
1970		31 789	676	131
1971		42 212	522	115
1972		49 133	1 444	225
1973		45 601	616	117
1974		52 275	472	98
1975		55 288	922	85
1976		133 400	970	NA
1977		214 900	856	NA

Year	(b) Longline	Total Snapper	SNA 7
1975		1 510	–
1976		2 057	–
1977		2 208	–

**Figure 1: Total reported landings and TACCs for SNA 7.**

## 1.2 Recreational fisheries

The snapper fishery is the largest recreational fishery in New Zealand. It is the major target species on the northeast and northwest coasts of the North Island and is targeted seasonally around the rest of the North Island and the top of the South Island. The current allowance within the SNA 7 TAC is shown in Table 3.

### 1.2.1 Management controls

The two main methods used to manage recreational harvests of snapper are minimum legal size limits (MLS) and daily bag limits. Both have changed over time (Table 5). The number of hooks permitted on a recreational longline was reduced from 50 to 25 in 1995.

**Table 5: Changes to minimum legal size limits (MLS) and daily bag limits used to manage recreational harvesting levels in SNA 7.**

Stock	MLS	Bag limit	Introduced
SNA 7	25	30	1/01/1985
SNA 7 (excl Marlborough Sounds)	25	10	1/10/2005
SNA 7 (Marlborough Sounds)	25	3	1/10/2005

### 1.2.2 Estimates of recreational harvest

A background to the estimation on recreational harvest of snapper is provided in the Introduction – Snapper chapter. Recreational harvest estimates for SNA 7 are provided in Table 6.

Plausible estimates for recreational catches from SNA 7 are available from the 1987 tagging programme, the aerial access surveys (in 2005–06 and 2015–16) and the national panel surveys (2011–12, 2017–18 and 2022–23). The estimates of recreational catch increased considerably from 2005–06 to 2017–18. The 2022–23 harvest estimate was similar to the previous (2017–18) estimate.

Most of the recreational catch has been recorded from Tasman Bay and Golden Bay. The catch is predominantly taken by rod-and-line, although a significant proportion of the catch was taken by longline during the mid 2010s. A small proportion of the total SNA 7 recreational catch was recorded from the Marlborough Sounds.

Boat ramp indices of snapper recreation catch have been developed from web cam monitoring at the main Nelson boat ramp. However, the indices are not considered to represent reliable indices of the snapper harvest for the entirety of SNA 7.

**Table 6: Recreational catch estimates for SNA 7. Totals for a stock are given in bold. The telephone/diary surveys ran from December to November but are denoted by the January calendar year. Mean fish weights were obtained from boat ramp surveys (for the telephone/diary and panel survey catch estimates). Numbers and mean weights are not calculated in the tag ratio method. Amateur charter vessel (ACV) and recreational take from commercial vessels under s111 general approvals as reported, with Total the sum of NPS, ACV and s111. ACVs have only been required to report harvest for SNA since 2020–21.**

Stock	Year	Method	Harvest survey			ACV (t)	s111 (t)	Total (t)
			Number of fish (0000)	Harvest estimate (t)	CV			
<u>SNA 7</u>								
Tasman Bay / Golden Bay	1987	Tag ratio	–	15	–			
Total	1993	Telephone/diary	77	184	–			
Total	1996	Telephone/diary	74	177	–			
Total	2000	Telephone/diary	63	134	–			
Total	2001	Telephone/diary	58	125	–			
Total	2005–06	Aerial-access	–	43	0.17			
Total	2011–12	Panel survey	110	88	0.17	0.4	1.7	90.3
Total	2015–16	Aerial-access	–	83	0.18			
Total	2017–18	Panel survey	96	144	0.16	1.0	13.4	158.5
Total	2022–23	Panel survey	88	130	0.14	7.6	1.6	139.0

### 1.3 Customary non-commercial fisheries

There are no estimates of customary catch available for SNA 7. Current levels of customary catch in SNA 7 are considered to be small and are assumed to be included into recreational catch estimates.

### 1.4 Illegal catch

No new information is available to estimate illegal catch. For modelling in SNA 7 an assumption was made that non-reporting of catch was 20% of reported domestic commercial catch prior to 1986 and 10% of reported domestic commercial catch since the QMS was introduced. This was to account for all forms of under-reporting. These proportions were based on the black-market trade in snapper and higher levels of under-reporting (to avoid tax) that existed prior to the introduction of the QMS. The 10% under-reporting post-QMS accounts for the practice of ‘weighing light’ and the discarding of legal-size snapper.

### 1.5 Other sources of mortality

No estimates are available regarding the amount of other sources of mortality on snapper stocks; although high-grading of longline fish and discarding of under-sized fish by all methods occurs. An at-sea study of SNA 1 commercial longline fisheries in 1997 (McKenzie 2000) found that 6–10% of snapper caught by number were under 25 cm (MLS). Results from a holding net study indicate that mortality levels amongst lip-hooked snapper caught shallower than 35 m were low.

Estimates for incidental mortality were based on other catch-at-sea data using an age-length structure model for longline, trawl, seine, and recreational fisheries. In SNA 1, estimates of incidental mortality for the year 2000 from longlines were less than 3% and for trawl, seine, and recreational fisheries between 7% and 11% (Millar et al 2001). In SNA 8, estimates of trawl and recreational incidental mortality were lower, mainly because of low numbers of 2- and 3-year old fish estimated in 2000.

In SNA 1, recreational fishers release a high proportion of their snapper catch, most of which was less than 27 cm (recreational MLS). An at-sea study in 2006–07 recorded snapper release rates of 54.2% of the catch by trailer boat fishers and 60.1% of the catch on charter boats (Holdsworth & Boyd 2008). Incidental mortality estimated from condition at release was 2.7% to 8.2% of total catch by weight depending on assumptions used.

## 2. BIOLOGY

For further information on snapper biology refer to the Introduction – Snapper chapter. A summary of published estimates of biological parameters for SNA 7 is presented in Table 7.

There is evidence of changes in snapper growth rates in SNA 7 over the history of the fishery. Growth rates were estimated to be higher in the 1990s and 2000s than the earlier and more recent periods.

A length-based maturity ogive was derived from ovarian staging data collected from west coast North Island inshore trawl surveys. Female snapper mature from about 25 cm in length and reach full maturity at about 40 cm with 50% maturity at 35 cm.

**Table 7: Estimates of biological parameters.**

Fishstock	Estimate			Source
<u>1. Instantaneous rate of natural mortality (<math>M</math>)</u>				
SNA 1, 2, 7, & 8	0.075			Hilborn & Starr (unpub. analysis)
<u>2. Weight = <math>a(\text{length})^b</math> (Weight in g, length in cm fork length)</u>				
All	$a = 0.04467$	$b = 2.793$		Paul (1976)
<u>3. von Bertalanffy growth parameters</u>				
	<u>Both sexes combined</u>			
	$K$	$t_0$	$L_\infty$	
SNA 7 (1990s)	0.122	-0.71	69.6	MPI (unpub. data)
<u>4. Age at recruitment (years)</u>				
SNA 7	3*			MPI (unpub. data)

\* Age at maturity in 2024 assessment based on ovarian sampling from WCNI trawl survey

A loss in genetic diversity has been associated with historical overfishing in SNA 7 (Bernal-Ramírez et al 2003).

## 3. STOCKS AND AREAS

New Zealand snapper are thought to comprise either seven or eight biological stocks based on: the location of spawning and nursery grounds; differences in growth rates, age structure, and recruitment strength; and the results of tagging studies. These stocks comprise three in SNA 1 (East Northland, Hauraki Gulf, and Bay of Plenty (BoP)), two in SNA 2 (one of which may be associated with the BoP stock), two in SNA 7 (Marlborough Sounds and Tasman Bay/Golden Bay) and one in SNA 8.

Tagging studies in SNA 7 (1986/87) and SNA 8 (1990) revealed reciprocal movements of snapper between Tasman Bay/Golden Bay and South Taranaki Bight (STB), although the scale of the movement was relatively low during that period.

Location-based snapper catch data from the trawl fisheries in SNA 7 and southern SNA 8 has revealed an overlap of the distribution of snapper catches in western approaches to Cook Strait between Durville Island and Kapiti Island, particularly since 2014/15. Snapper age compositions are available from recent (2018–2020 and 2022) *Kaharoa* trawl surveys of the South Taranaki Bight and the Tasman Bay/Golden Bay area of the WCSI trawl survey. There are strong differences in the relative strength of individual year classes from the 2019 South Taranaki Bight age composition compared to the 2018 and 2020 surveys, while the 2019 STB age composition was very similar to the age structures from the 2019 Tasman Bay/Golden Bay trawl survey and the commercial fishery in the TBGB area. These observations indicate a degree of mixing of the snapper populations between SNA 7 and the STB area (SNA 8), although the extent of mixing may vary between years, potentially related to variation in the timing of the main spawning period in each area.

The 2022 STB survey age composition was dominated by a very strong 5 yr age class, representing the 2017 year class. The 2017 year class was not present as a strong 3 yr age class in the previous (2020) survey, suggesting an immigration of snapper into the STB region. The year class appeared to be moderately strong in the 2018–2020 survey age compositions from the northern area of SNA 8 (at ages 1, 2 and 3 yr, respectively) but was not particularly strong in the 2022 survey age composition. The year class was observed to be very strong in Tasman Bay and Golden Bay when surveyed at 1, 3 and 5 years of age and represented the dominant age class in the 2022–23 age composition from the SNA 7 commercial fishery.

During 2022–23, catch sampling was conducted from the SNA 8 trawl fishery, partitioned between the northern and southern areas. The age composition of the commercial fishery in SBT was very similar to the age composition from the 2022 trawl survey with the dominance of the 5 yr age class. For the three fisheries, there were broad similarities in the relative proportion of fish in the older (greater than 9 years) age classes. A comparison of the average length at age from the three areas revealed that initial growth rates were faster for fish sampled from Tasman Bay and Golden Bay, while growth rates were similar between STB and northern SNA 8 up to age 5 years. For older age classes, the average length of age diverged between STB and northern SNA 8, with average length at age for STB approximating Tasman Bay and Golden Bay from about 7 years of age.

#### 4. STOCK ASSESSMENT

An assessment for SNA 7 was conducted in 2015 and updated in 2018, 2020, 2021 and 2024.

##### **SNA 7 (Challenger)**

The SNA 7 fishery is concentrated within Tasman Bay and Golden Bay and this area is considered to represent the main spawning area and nursery area for the stock. Most of the main data sets included in the stock assessment were derived from the Tasman/Golden Bay area. However, since the mid 2010s there has been an increase in the spatial domain of the stock, particularly for older fish, with the distribution extending into deeper areas beyond TBGB (i.e., the western approaches to Cook Strait) and southward along the west coast of the South Island. It is currently assumed that fish in these areas migrate to spawn in the shallower areas of TBGB during late October-early December and disperse during late summer.

A stock assessment of SNA 7 was conducted in 2024. The assessment updated and refined the previous stock assessments conducted during 2015–2021 (see Langley 2021a). Those assessments were primarily based on a time series of CPUE indices from the SNA 7 trawl fishery, in addition to age compositional data from the trawl fishery and a tagging biomass estimate from 1987. The current stock assessment also incorporated the snapper biomass estimates, and the associated length and age compositions, from the time-series of *Kaharoa* inshore trawl surveys of west coast South Island and Tasman Bay/Golden Bay.

The 2024 stock assessment of SNA 7 was conducted using an age-structured population model implemented in Stock Synthesis. The model incorporated data to the 2023–24 fishing year (2023 model year) including:

- Commercial catches by method, 1931–2023;
- Recreational catches, 1931–2023;
- Tag biomass estimate 1987;
- Seasonal (Oct-Dec, Jan-Apr) single trawl CPUE indices 1989–2022;
- *Kaharoa* trawl survey biomass indices (1991–2022) and length/age compositions;
- Single trawl catch age compositions 1992–2022;
- Pair trawl catch age compositions 1975–1983; and
- Recreational catch length compositions 2005–2021.

### Commercial catches

Commercial catch data are available for the SNA 7 fishery from 1931 to the 2022–23 fishing year. The model data set was configured to include three commercial fisheries: two seasonal single trawl fisheries (BT) in October–December (BT1) and January–September (BT2) and a pair trawl fishery (BPT). The SNA 7 catch taken by the purse-seine method during the late 1970s and early 1980s was assigned to the pair trawl fishery, as both methods are considered to harvest the full range of adult age classes in the population.

The seasonal division of the BT catch followed the derivation of separate seasonal CPUE indices and enabled the evaluation of different assumptions regarding the seasonal availability (selectivity) of snapper to the BT fisheries.

The reported commercial catches from 1931–1986 were increased by 20% to account for an assumed level of under-reporting. Since the introduction of the Quota Management System (QMS), the accuracy of the reporting of commercial catches has improved considerably, although a degree of under-reporting may persist. For 1987–2023, reported catches were increased by 10% to account for the assumed level of under-reporting in the more recent period. These assumptions are consistent with the formulation of the commercial catch histories incorporated in other inshore finfish stock assessments (based on assumptions for SNA 1 and SNA 8 made according to quota appeals when the QMS was first introduced).


The base assessment model initialised the model in 1975 under exploited conditions. This removed the influence of the earlier catches which are considered to be much less reliable than the more recent catches. The full catch history was retained in a model sensitivity that was initialised in 1931 assuming equilibrium, unexploited conditions.

### Non-Commercial catches

The recreational catch history was constructed based on estimates of recreational catch from 1987, 2005–06, 2011–12, 2015–16, 2017–18 and 2022–23 (Figure 2). The point estimates were used to determine estimates of recreational exploitation rates in each year based on the annual estimates of biomass from preliminary model runs. Exploitation rates were interpolated between successive recreational catch estimates to determine annual estimates of recreational catch from 1987 to 2021. For the period prior to 1987, the exploitation rate was extrapolated, declining by 10% per annum, to the early 1960s when a lower threshold of 10 t per annum was attained. Length compositions from the recreational fishery (2005, 2011, 2015–2021) were derived from sampling conducted during boat ramp interviews.

Two options were assumed for the 2023–24 recreational catch: the catch was set equal to the 2022–23 harvest estimate (139.1 t) or was derived based on the average of the recreational fishery exploitation rates corresponding to the last three NPS harvest estimates (213 t).

There are no estimates of customary catch available for SNA 7. Recent customary catches are likely to have been a minor component of the total catch and are not explicitly included in the model catch history.



**Figure 2: Commercial (top) and recreational (bottom) catch histories for SNA 7 included in the stock assessment models. The commercial catch history attributes all the single trawl catch prior to 1989 to the BT1 fishery. Commercial catches include an allowance for 20% unreported catch prior to the QMS and 10% allowance in the subsequent years. The red points represent the survey estimates of recreational catch. Only catches from 1975 onwards were included in the base assessment model.**

### Tagging biomass estimate


An estimate of 1987 stock biomass was derived from a tag release-recovery programme (Kirk et al 1988). A subsequent reanalysis of the tagging data yielded a very similar estimate of snapper biomass (1549 t) Harley & Gilbert (2000). Harley & Gilbert (2000) expressed concerns regarding the reliability of the 1987 tag biomass estimate due to spatial heterogeneity of tagged fish and the lack of tag releases in deeper water. Consequently, the tag biomass estimate was assigned a moderate level of precision (CV 30%).



### CPUE indices

The previous stock assessments of SNA 7 incorporated a time series of CPUE indices as a primary index of stock abundance. The CPUE indices were based on catch and effort data from the Tasman Bay/Golden Bay trawl fishery targeting snapper, flatfish, red gurnard, and, to a lesser extent, barracouta during October–April. A detailed analysis of catch and effort data from the fishery indicated that since 2010–11 the operation of the trawl fishery had changed to increasingly avoid snapper, particularly during October–December. There was also some indication that the age composition of the snapper catch may vary between October–December and January–April. On that basis, separate sets of trawl CPUE indices were derived for the two seasons (BT1 and BT2). The analyses included catch and effort data from the 1989–90 to 2022–23 fishing years, aggregated by vessel fishing day. For each seasonal data set, a GLM approach was applied to separately model the probability of catching snapper (binomial model) and the magnitude of positive (non-zero) snapper catch (lognormal model) and the combined CPUE indices (delta-lognormal) were derived from the annual coefficients of the two models.

Due to the increase in snapper avoidance, the more recent (2010–2022) October–December (BT1) CPUE indices were not included in the assessment modelling.



**Figure 3:** Relative CPUE indices derived from the delta lognormal (all years) model for the single trawl fishery during October–December (left) and January–April (right). The vertical lines represent the 95% confidence intervals. The confidence intervals were derived using a bootstrapping procedure.

The BT1 CPUE indices decline during the early 1990s and then remain at the lower level until 2010–11. The time series of BT2 CPUE indices are relatively constant during 1989–90 to 2009–10, increase initially in 2010–11 and then increase substantially in 2011–12. The indices fluctuate considerably about the higher level during the subsequent years (Figure 3). The scale of the variation in the BT2 CPUE indices may indicate a high degree of inter-annual availability of snapper within the TBGB area during summer/autumn.

### Trawl survey

The West Coast South Island inshore trawl survey, including the Tasman Bay/Golden Bay area, commenced in 1992 and has been conducted biennially since 2002. The survey occurs in March–April coinciding with the period when larger, mature snapper are dispersing from TBGB following the spawning season. The survey area does not extend out into the deeper waters of the western approaches to Cook Strait but does include the west coast of the South Island extending from Farewell Spit in the north, as far as Haast in the south, in waters ranging from 20–400 metres for the core strata.

Prior to 2009, so few snapper were caught that the length frequency distribution was sparse and uninformative (MacGibbon et al 2024). Large numbers of 1+ snapper (around 14–19 cm) were caught on the 2009 survey (Stevenson & Hanchet 2010). This indicated that a strong year class of fish was spawned over the summer of 2007–08. This year class was dominant in the length frequency distribution from 2013 until about 2019 (MacGibbon 2019). Also visible were fish from the 2011 and 2013 year classes, which were also relatively strong (Parker et al 2015, Parsons et al 2018) and could be tracked through subsequent surveys. These strong year classes and the apparent increasing abundance of snapper in the Tasman Bay and Golden Bay region were the impetus for expanding the trawl survey area to include two new strata in 10–20 m, one in each of Tasman Bay and Golden Bay, beginning with the 2017 survey (Stevenson & MacGibbon 2018).

There was a conspicuous absence of 2+ fish between 22 and 29 cm in 2019 (Walsh et al 2019). Even with a 60 mm codend these fish would not likely have been caught as 0+ fish in the 2017 survey, and there was no survey in 2018 where they might have been caught as 1+ fish. Oddly however, the 2021 survey caught them as 4+ fish, though not in great numbers (MacGibbon et al 2022), and again as 6+ fish in 2023. This suggests that there may be some variability in the catchability of snapper to the survey. Since being introduced, the 10–20 m strata are where the majority of juvenile fish (under 25 cm) have been caught each year.

This expansion is now in its fourth year and the 2023 snapper biomass estimate for the expanded survey was the highest in the time series at 4404 t, almost three times the estimate from 2021 (the previous highest estimate) (MacGibbon et al 2024). At 3633 t, biomass in the core survey was more than four times the core estimate in 2021. Biomass increased in all strata in Tasman Bay and Golden Bay in 2023, but these were only modest increases for the 10–20 m strata. Stratum 17 (Golden Bay) saw a nearly twofold increase from 230 to 454 t, stratum 18 (Tasman Bay) increased more than 7-fold from 125 to 885 t, and stratum 19 increased more than 4-fold from 399 to 1606 t. Over the time series, core snapper biomass was low and showed a relatively flat trend until 2013, from which time there was a step increase to around 1000 t, before the massive increase in 2023. Biomass has increased every year in the core plus 10–20 m strata, especially in 2023.

More snapper biomass has come from Tasman Bay and Golden Bay than from the west coast, throughout the time series, including the period before the stock began to rebuild, when biomass was low in all areas (MacGibbon et al 2024). The west coast biomass increased in 2019, slightly decreased in 2021, and substantially increased in 2023 (by a factor of more than 5). As a proportion of the total, the biomass from the west coast in 2023 (16%) was more than in 2021 (8%), but similar to 2019 (14%).

The length frequency distribution for snapper from the west coast has typically been sparse compared with the distribution from Tasman Bay and Golden Bay and numbers were much lower (MacGibbon et al 2024). Fewer smaller fish were caught here and, apart from a very small number in 2009, no juveniles were caught off the west coast. Fish as large as those seen in Tasman Bay and Golden Bay were caught, albeit in much lower numbers. While still only a fraction of the number seen in Tasman Bay and Golden Bay, record numbers of fish were seen off the west coast in 2023.

Throughout the time series, juvenile fish have comprised only a small minority of the total biomass (MacGibbon et al 2024). The introduction of the 10–20 m strata in 2017 indicated that while these strata were important for juvenile snapper, most of the total biomass was adult fish.

### **Commercial age compositions**

Commercial age frequency data are available from the TBGB BPT fishery from the pre QMS era (N=4) and BT from the QMS era (N=11). The annual BPT age compositions were derived from a small number of sampled landings. These data provide information regarding the age composition during the period of the highest catches from the fishery.

The more recent BT age compositions (2006, 2013, 2016, 2019 and 2022) were partitioned between the two seasons. In some years, a higher proportion of older fish were sampled from October–December (BT1) compared to January–April (BT2). This may indicate that older snapper are more available in

TBGB during the main spawning period and subsequently disperse from the TBGB area over the following summer. The BT age compositions from the earlier years (1992, 1997–2000 and 2003) were assumed to represent the age composition of snapper from October–December (BT1), because most of the sampling took place during that period.

### Model structure and assumptions

A statistical age-structured population model for SNA 7 was implemented using Stock Synthesis (Methot & Wetzell 2013). A summary of input data, fixed and estimated parameters are provided in Tables 8, 9 and 10. The main model structural assumptions for the base model are as follows:

- The initial population (1975) was initialised under exploited conditions and was configured as a single sex model comprised of 30 age classes, including a plus group. The model data period is 1975–2023 (the 2023 model year represents the 2023–24 fishing year) and includes two seasons (October–December and January–September). The initial age structure was derived by estimating an equilibrium fishing mortality rate and recruitment deviations for the initial age structure (1960–1974 year classes).
- The estimation of the initial equilibrium fishing mortality rate was informed by a prior (normal, mean 0.2, sd=0.1). The prior was required to stabilize the estimation of the initial  $F$  and recruitment parameters. The prior did not influence the estimate of absolute biomass for the period after 1980.
- Annual recruitments for 1975–2021 were estimated as unconstrained deviates from a Beverton-Holt stock-recruitment relationship (SRR) with steepness of 0.95. Recruitment for 2022–23 was assumed based on the average level of recruitment from the recent period (2010–2019).
- Commercial fisheries (BPT, BT1 and BT2) selectivities are age-based and temporally invariant. The three fisheries have full selection for all recruited age classes (parameterised using a logistic selectivity function).
- Age based selectivity for the *Kaharoa* trawl survey (core area) was parameterised using a logistic selectivity function. Temporal variation in the selectivity ( $a_{50}$ ) was estimated for the trawl surveys from 2012–13 to 2022–23 to account for an apparent increase in the availability of younger fish.
- The age compositions from the 2018, 2020, 2022 core + SNA survey area were considered to represent estimates of the full population age structure (i.e., selectivity 1.0 for all age classes). This assumption was required to adequately inform the model regarding the relative abundance of the youngest age classes (1–4 y) in conjunction with the longer time series of data from the *Kaharoa* trawl survey (core area). Model trials that estimated the selectivity function for the core + SNA survey resulted in a comparable selectivity function.
- The selectivity of the recreational fishery is length-based and parameterised using a double normal function. Selectivity is configured with three time blocks (pre-2013, 2013–2015, and 2016 onwards) to account for the increase in the catch of larger fish by the longline method in the intermediate period and increased targeting of larger fish in more recent years.
- The two sets of CPUE indices (BT1 and BT2) were assigned additional process error of 30% and 20%, respectively, based on RMSE from preliminary model runs. Similarly, the *Kaharoa* trawl survey (core area) biomass indices were assigned process error of 20%.
- The tag biomass estimate was assumed to represent the proportion of the stock biomass that had recruited to the commercial BPT fishery in 1987. The tag biomass estimate was assigned a CV of 30% following Harley & Gilbert (2000). The moderate CV was adopted to reflect concerns regarding the reliability of the tag biomass estimate.
- The relative weighting of the individual commercial age compositions were assigned based on the MWCV of the observations. The final relative weightings (ESS) of the BT1, BT2 and BPT age compositions were determined following the approach of Francis (2011). The two sets of trawl survey age compositions were each assigned an ESS of 50 to ensure that these data were informative in the model estimation of recent recruitments.
- The recreational length compositions were also assigned an ESS of 1.0, as they may not fully represent the fishery, and the selectivity of the recreational fishery appears to have changed over time.

- Growth rates of snapper in TBGB were variable over the model period with higher growth of younger fish occurring during 1990–2009. The model was divided into three periods (pre-1990, 1990–2009 and post-2009), based on these observed differences in growth rates. The length-at-age data from each period was used to estimate time specific values of the  $k$  VB growth parameter. The time-specific growth functions were applied to the three time periods of the model.

**Table 8: Summary of input data sets for the Base Case assessment model. The relative weighting includes the Effective Sample Size (ESS) of age/size composition data and the coefficient of variation (CV) associated with the abundance data. Note that model year 2022, is fishing year 2022–23, and includes the trawl survey conducted in March 2023.**

Data set	Model years	Nobs	Error structure	Observation error/ESS	Process error
Tag biomass	1987	1	Lognormal	0.30	–
BT1 CPUE indices	1989–2009	21	Lognormal	0.09–0.16	0.3
BT2 CPUE indices	1989–2022	34	Lognormal	0.14–0.20	0.2
Trawl survey Core indices	1991, 1993, 1994, 1996, 1999, 2002, 2004, 2006, 2008, 2010, 2012, 2014, 2016, 2018, 2020, 2022	16	Lognormal	0.13–0.94	0.2
Trawl survey Core age comp	2016, 2018, 2020, 2022	4	Multinomial	ESS 50	
Trawl survey Core length comp	2008, 2010, 2012, 2014	4	Multinomial	ESS 10	
Trawl survey Core+SNA age comp	2018, 2020, 2022	3	Multinomial	ESS 50	
BT1 age comp	1992, 1997–2000, 2003, 2006, 2013, 2016, 2019, 2022	11	Multinomial	ESS 20	
BT2 age comp	2006, 2013, 2016, 2019, 2022	5	Multinomial	ESS 10	
BPT age comp	1978–1980, 1983	4	Multinomial	ESS 18	
Recreational length comp	2005, 2011, 2015–2021,	9	Multinomial	ESS 1	

**Table 9: Details of parameters that were fixed in the base model.**

Natural mortality	0.075 $y^{-1}$
Stock-recruit steepness (Beverton & Holt)	0.95
Std deviation of rec devs (sigmaR)	1.5
Proportion mature (female, length-based logistic)	L50% = 35 cm, L5% 26 cm, L95% 44 cm
Length-weight [mean weight (kg) = $a$ (length (cm)) <sup><math>b</math></sup> ]	$a = 3.61 \times 10^{-5}$ , $b = 2.8644$
Growth parameters	$L_{\infty} = 69.6$ , Length1=13.1
pre-1990 (1), 1990-2009 (2), post- 2009 (3)	$k1=0.098$ , $k2=0.122$ , $k3=0.103$ ,
Coefficients of variation for length-at-age	0.075

**Table 10: Estimated parameters for the base model.**


Parameter	Number of parameters	Parameterisation, priors, constraints
$\ln R_0$	1	Uniform, uninformative
Initial equilibrium F	1	Normal(0.2,0.1)
Initial Rec devs (1960–1974)	15	SigmaR 1.5
Rec devs (1975–2021)	47	SigmaR 1.5
Selectivity BPT commercial	2	Logistic
Selectivity BT1 commercial	2	Logistic
Selectivity BT2 commercial	5	Logistic
Selectivity trawl survey core	8	Logistic
Selectivity trawl survey core+SNA	–	Constant
Selectivity tag	–	Equivalent to commercial BPT
Selectivity Recreational	8	Double normal
CPUE BT1 $q$	1	Uniform, uninformative
CPUE BT2 $q$	1	Uniform, uninformative
Trawl Survey $q$	1	Uniform, uninformative

For the base model option, the model biomass approximates the point estimate of the 1987 recruited biomass from the tagging programme (Figure 4). The model also provides a good fit to the time series


of BT2 CPUE indices to 2010. Stock biomass is predicted to have increased considerably from 2010 (2010–11 fishing year) following the overall magnitude of the increase in trawl survey biomass indices and BT2 CPUE indices. However, the fits to the individual BT2 CPUE indices from 2011–12 to 2022–23 are relatively poor (Figure 4). The fit to the BT1 CPUE indices (1989–2009) is poor (Figure 4).

The model provides a good fit to the time-series of trawl survey biomass indices, with the exception of the 2022 index. The model estimate of the vulnerable trawl survey biomass is approximately half of the observed biomass index. A range of model options were investigated to endeavour to fit the high recent biomass index. None of those options provided credible results as they substantially degraded the fit to the other recent data sets.

The initial increases in the CPUE and trawl survey biomass indices are consistent with the recruitment of the very strong 2007 year class (Figure 5). This year class dominated the age compositions from the trawl fishery and (core) trawl survey during 2013–14 to 2018–19. More recent age compositions were augmented by the recruitment of subsequent year classes, most notably the 2010 year class. The 2018–19 and 2020–21 trawl surveys (core + SNA) yielded relatively high catch rates of juvenile snapper in the shallower TBGB strata (10–20 m), which dominated the associated age compositions. Correspondingly, the model estimated exceptionally strong 2017 and 2018 year classes. These year classes also dominated the recent (2022–23) age compositions sampled from the commercial fishery. The most recent (2022–23) trawl survey indicated that recent (2019–2021) recruitments were low.



**Figure 4: Biomass trajectories (MPD) for the base model option presenting the fit to the tag biomass estimate (top left panel), trawl survey (core) biomass indices (top right panel) and the CPUE indices (lower panels).**



**Figure 5: Annual recruitment for the base model (MCMC results). The main recruitment deviates were estimated for 1975–2021. The line represents the median and the shaded area represents the 95% credible interval.**

A range of model trials was conducted to investigate the relative influence of the individual data sets. These trials revealed that estimates of recent biomass were relatively insensitive to the relative weighting of the CPUE indices and trawl survey biomass indices and age compositions. The increasing trend in stock biomass was driven by the recent BT2 CPUE indices; the trawl survey biomass indices were less informative due to the higher variability (and lower frequency) of the series.

The base model provides estimates of current stock status that are quite uncertain, primarily due to the uncertainty associated with the estimates of the strength of recent recruitment (2017 and 2018 year classes). There is also uncertainty associated with the scale of the increase in stock abundance due to differential trends in the increase of the two principal abundance indices and the associated assumptions regarding fishery/survey selectivities. A range of model sensitivities was undertaken to investigate model assumptions, these included a lower value of natural mortality (0.06 compared with 0.075), a lower value of variation in the recruitment deviates (sigmaR 1.1 compared with 1.5), alternative parameterisation of the recruitment deviates, a lower value of steepness (0.85) and the removal of time varying selectivity of the trawl survey (Table 11). In addition, a model option incorporated additional snapper catch from South Taranaki Bight (FMA 8) to consider a wider spatial definition of the stock. The sensitivities were generally treated as single changes from the base model. A full catch history model, initialising the unexploited equilibrium population in 1931, was retained for comparison with the structure of the previous (2022) assessment.

**Table 11: Description of model sensitivities.**

Sensitivity run	Description
NatMort sensitivity	$M = 0.06$
RecDev variation sensitivity	$\text{sigmaR} = 1.1$
RecDev vector	Constrained rec dev vector
Steepness sensitivity	$h = 0.85$
Init 1975 (Pseudo Fishery) sensitivity	Alternative initialisation of model in 1975
Trawl Survey selectivity sensitivity	Temporally invariant trawl survey selectivity
Stock structure sensitivity	Include FMA 8 catches
Full catch history sensitivity	Initialise model in 1931, unexploited equilibrium conditions, rec devs 1960–2021.

Stock status was determined for 2023 (= 2023–24 fishing year) relative to the interim target fishing mortality level, i.e.,  $U_{SB40\%}$ . the default value for a low productivity stock as described by the Harvest Strategy Standard. Spawning biomass was assessed relative to hard and soft biomass limits proposed by the INS WG (see Derivation of Reference Points – below).

B, biomass is estimated to have increased considerably from 2010. Current (2023) biomass is well above (%) the soft limit (Figure 6, Table 12). For all model options, current rates of fishing mortality are well below the corresponding exploitation rate based target ( $U_{SB40\%}$ ) (Figure 7, Table 12).





Figure 6: Annual trend in spawning biomass for the base model. The line represents the median and the shaded area represents the 95% confidence interval. The projection period (2024–2028) is in red (*RecF* option).

Table 12: Estimates of current (2023–24) spawning biomass (t) and fishing mortality relative to  $U_{SB40\%}$  (median and the 95% confidence interval from the MCMCs) and the probability of being below the target  $U_{SB40\%}$ . Estimates of current yield (t) at  $U_{SB40\%}$  at the 2023–24 biomass levels are also provided.

Model	$SB_{2023}$	$U_{SB40\%}$	$U_{2023}/U_{SB40\%}$	$Pr(U_{2023} < U_{SB40\%})$	Yield $U_{SB40\%}$
<i>Base_RecCatch</i>	13 816 (9 810–18 530)	5.1%	0.376 (0.278–0.532)	1.00	1 872 (1 313–2 537)
<i>Base_Frec</i>	13 154 (9 293–18 241)	5.0%	0.453 (0.322–0.633)	1.00	1 778 (1 239–2 517)
<i>FMA8catch</i>	21 623 (16 312–29 671)	5.1%	0.32 (0.232–0.428)	1.00	2 958 (2 194–4 089)
<i>LowM</i>	14 567 (10 624–19 567)	4.4%	0.417 (0.309–0.566)	1.00	1 687 (1 236–2 329)
<i>PseudoFishery</i>	13 624 (9 587–20 087)	5.1%	0.381 (0.252–0.542)	1.00	1 841 (1 283–2 821)
<i>RecDevVector</i>	13 915 (9 683–19 529)	5.1%	0.373 (0.262–0.53)	1.00	1 881 (1 309–2 678)
<i>SigmaR11</i>	11 457 (8 482–15 625)	5.0%	0.457 (0.335–0.622)	1.00	1 541 (1 116–2 117)
<i>Steepness85</i>	13 976 (10 051–19 517)	4.8%	0.393 (0.282–0.551)	1.00	1 788 (1 248–2 516)
<i>TSurveySelect</i>	15 096 (11 046–20 961)	5.0%	0.33 (0.237–0.458)	1.00	2 192 (1 558–3 094)
<i>start1931</i>	13 212 (9 477–18 338)	5.0%	0.401 (0.283–0.57)	1.00	1 759 (1 222–2 527)



**Figure 7: Annual trend in fishing mortality relative to the  $U_{SB40\%}$  interim target biomass level for the base model. The line represents the median and the shaded area represents the 95% credible interval. The projection period (2024–2029) is in red (*RecF* option). The dashed line represents the interim target level.**

For all model options, estimates of current yield were derived for the stock based on the target fishing mortality rate ( $U_{SB40\%}$ ).  $U_{SB40\%}$  yields at 2023–24 biomass levels are about 1800 t, well above the level of current catch (626 t or 708 t), which includes commercial catch, other sources of mortality and recreational catch. The higher biomass and yield for the *FMA8catch* sensitivity reflects the addition of the southern SNA 8 catches encompassed in the model.

### Projections

Projections were conducted for the base model with two options for recreational catches: constant annual catch equivalent to the 2022–23 recreational harvest estimate (131 t) (*RecCatch*) or a constant recreational fishing mortality (the average of the last three recreational harvest estimates) (*RecF*). Stock projections were conducted for the 5-year period following the terminal year of the model (i.e., 2024–2028). Projections assumed future recruitments were resampled from the lognormal distribution around the geometric mean of recent recruitments. Commercial catches in the projection period were held constant at the current TACC of 450 t with an allowance for additional mortality of 45 t. There was no explicit allowance for customary catch.

The projections are strongly influenced by the high estimates of recent recruitment of the 2017 and 2018 year classes, resulting in an increase in total biomass during the projection period (Figure 6). Successive recent surveys and commercial age composition data have reaffirmed that these year classes are very strong although the magnitude of the year classes is uncertain.

For the base case (and all other model options), stock abundance is predicted to continue to increase in the projection period and fishing mortality remains well below the target level ( $F_{SB40\%}$  level) throughout the projection period (Table 13). Under the constant fishing mortality assumption, recreational catches are projected to increase in proportion to the overall increase in stock biomass.



**Table 13: Annual commercial (set at TACC, including an additional 10% unreported) and recreational catches (assumed or predicted) for the last three years of the model period and the five-year projection period (shaded) from the *RecF* and *RecCatch* models. The annual stock status ( $F_{year}/F_{SB40\%}$  and  $SB_{year}/SB_{2023}$ ) for the projection period is also presented for both model options (with associated 95% confidence intervals).**

Year	Commercial catch (t)	Recreational catch (t)		$SB_{year}/SB_{2023}$		$F_{year}/F_{SB40\%}$	
		<i>RecF</i>	<i>RecCatch</i>	<i>RecF</i>	<i>RecCatch</i>	<i>RecF</i>	<i>RecCatch</i>
2021–22	495	161	161				
2022–23	495	131	131				
2023–24	495	213	131				
2024–25	495	230 (123–353)	131	1.14 (1.10–1.18)	1.14 (1.11–1.18)	0.44 (0.34–0.57)	0.35 (0.26–0.51)
2025–26	495	237 (120–370)	131	1.24 (1.18–1.32)	1.25 (1.18–1.34)	0.41 (0.33–0.54)	0.33 (0.24–0.48)
2026–27	495	240 (112–382)	131	1.31 (1.22–1.42)	1.33 (1.23–1.45)	0.40 (0.31–0.53)	0.32 (0.23–0.46)
2027–28	495	242 (110–394)	131	1.36 (1.25–1.50)	1.38 (1.26–1.53)	0.39 (0.28–0.52)	0.30 (0.22–0.45)
2028–29	495	249 (111–405)	131	1.39 (1.26–1.56)	1.41 (1.28–1.58)	0.38 (0.26–0.51)	0.29 (0.21–0.44)

### Derivation of Reference Points

Substantial increases in annual recruitment suggested an increase in productivity, and possibly a regime shift, for SNA 7. Owing to the complexities associated with estimating  $SB_0$  under these circumstances, the Inshore Working Group made the decision to base the target reference point on exploitation rate instead of biomass as a proportion of  $SB_0$ . Consistent with international best practice the hard and soft limits were based on absolute biomass.

The default target accepted for SNA 7 was the exploitation rate that, if applied perfectly over the long term and assuming equilibrium recruitment, would produce a spawning biomass of 40% of that in the absence of fishing ( $F_{SB40\%}$ ;  $U=5\%$ ).

After considerable discussion on options for deriving biomass based hard and soft limits the WG agreed that the hard limit should be twice the 1987 spawning biomass estimate. This year followed a large decline in biomass following peak catch from the fishery and preceded a stable but very low biomass period that previous assessments had suggested was well below the default HSS hard limit of 10%B0. The 1987 tagging study biomass estimate is for vulnerable biomass and is fitted perfectly by the current model. The stock assessment spawning biomass estimate from the base case biomass trajectory is used for establishing the hard limit. The soft limit is twice the biomass of the hard limit. The 2024 May Plenary supported this approach for setting the hard limit, but recommended future review if new information becomes available.

### Qualifying comments

The 1987 tag biomass estimate is considered to be an underestimate of the total recruited biomass due to the relatively small proportion of older fish estimated to be in the tagged fish population. However, model testing, either excluding or increasing the tag biomass estimate, has indicated that the assessment is relatively insensitive to the tag biomass estimate, especially with the assumed level of precision (CV 30%).

For the earlier assessments, the main abundance indices were CPUE indices derived for the entire fishing season (October–April). Subsequent analyses revealed increasing avoidance of snapper by the trawl fleet since 2010, especially during the spawning season (October–December). For the current assessment, the main set of CPUE indices was derived for January–April. The degree of avoidance of snapper is likely to be considerably less during this period. Nonetheless, it is considered that the main CPUE indices are also likely to underestimate the extent of the increase in snapper abundance from 2010 onwards. There is, however, a general correspondence between the increase in the CPUE indices and the abundance indices from the *Kaharoa* trawl survey.

The increased avoidance of snapper has included changes in the configuration of the trawl gear over the last 3–5 years, including a reduction in the headline height to reduce the catch of snapper, particularly larger fish. The overall effect of this change in gear configuration has not been quantified but it may have resulted in a change in the selectivity of the trawl fishery, particularly for older fish. This effect has not been incorporated in the assessment modelling.

The CPUE indices are derived for the Tasman Bay and Golden Bay fishery only. There has been an expansion of the range of snapper over the last 10 years, especially southward along the west coast of the South Island. The recent WCSI inshore trawl survey indicated that about 19% of the snapper biomass was off the west coast of the South Island in March of 2023. The disproportional increase in the stock in this area is not reflected in the CPUE abundance indices.

The time-series of core area trawl survey biomass indices was included in the stock assessment. This component of the survey does not adequately monitor the younger (1–4 years old) snapper which are predominantly found within the shallower areas (<20 m) of Tasman Bay and Golden Bay not included in the core survey area.

The older fish in the population (greater than 10 years) do not appear to be fully available to the trawl survey, at least in some years. The trawl survey occurs during late summer and older fish may have already dispersed to areas outside the survey area, following spawning in November–December. There are significant catches of snapper taken within SNA 7 outside of the trawl survey area, i.e., in the deeper areas in the western approaches to Cook Strait. The distribution of older snapper may also extend into the southern areas of SNA 8 (South Taranaki Bight and Kapiti coast). The proportion of the older fish within the survey area may vary between years depending on the timing of the dispersal of larger snapper from Tasman Bay/Golden Bay. The most recent survey age composition was poorly fitted in the assessment models and, correspondingly, the selectivity function for the trawl survey is not well estimated, particularly for the older age classes.

Recent (2017) modifications of the trawl survey design to include the shallower areas of Tasman Bay/Golden Bay (SNA strata) have improved the utility of the survey for monitoring of SNA 7, particularly for younger (1–4 year old) fish. However, the limited number of observations (3 surveys) meant that there were insufficient data to reliably estimate the selectivity for snapper within the shallower area separately from the core area. Therefore, the inclusion of two sets of age compositions (core and core+SNA strata) in the assessment model duplicated the age composition data from the core survey area, effectively doubling the influence of these data in the assessment model. Down-weighting these two data sets in the model likelihood resulted in an overall improvement in stock status. The trawl survey age composition data were informative due to the lower proportion of older snapper from the 2020–21 survey, particularly relative to the age composition from the October–December trawl fishery.

The most recent trawl surveys have reaffirmed the presence of the very strong 2017 and 2018 year classes, while the more recent year classes (2019–2021) are relatively weak.

## Future research considerations

### Trawl surveys

- The modified WCSI RV *Kaharoa* survey (extra snapper strata in Tasman and Golden Bays) is monitoring the abundance and age composition of younger (1–4 year old) snapper enabling recent recruitments to be estimated in the stock assessment model and should be continued on the current biennial basis.
- Consideration around extending the current trawl survey to include the South Taranaki Bight and Kapiti area to improve monitoring of the snapper in the wider area that are potentially part of the same stock. Currently, this area is monitored by the WCNI trawl survey which is conducted in October–November and may not monitor the portion of snapper migrating to spawn in Tasman/Golden Bay. Correspondingly, some of those fish that spawned in Tasman/Golden Bay will not be monitored by the current March–April trawl survey.

- Explore alternative approaches to incorporating data from the new TB/GB snapper strata into the assessment, including accounting for selectivity differences.

#### Age sampling of commercial catches

- A cycle for two consecutive years in five is recommended for SNA 7 shed sampling. Sampling should be stratified by area and season (and possibly target species) to enable a comparison between the age structure of catches from the spawning and post spawning periods and potential differences in age structure related to fishing depth. Sampling of the west coast South Island trawl fishery should be maintained.
- The current programme of concurrent sampling of SNA 7 and SNA 8 (particularly the south Taranaki region) will enable further evaluation of the connectivity between the two QMAs. This should be continue in future.

#### CPUE analysis

- Improved spatial and seasonal modelling of CPUE data from SNA 7 and southern SNA 8 to improve understanding of movement between areas.
- Development of CPUE indices to account for changes in the spatial distribution of snapper (e.g. using VAST).
- Explore changes in fishing operations related to increased avoidance or preferential targeting of snapper to qualify the utility of CPUE indices for the monitoring of trends in snapper abundance.
- Investigate splitting the vessels with long time series into two or more pseudo vessels.

#### Recreational harvest estimates

- Recreational fishing has accounted for significant proportion of the total catch from SNA 7. In addition to the NPS, there should be ongoing sampling of the recreational catch of snapper from boat ramps; such data also need to be analysed in more detail. Boat ramp data may also provide the opportunity to collect additional size composition data from the recreational fishery, which could be used to derive age compositions.
- At presently there is only monitoring of the Nelson boat ramp and consideration should be given to monitoring additional ramps, e.g., Golden Bay and Motueka.

#### Assessment model structure

- Changes in stock abundance and age composition over the last decade appear to have expanded the stock distribution, with older fish now extending down the west coast of the South Island and probably into South Taranaki Bight. These changes may be influencing seasonal availability to the fishery and trawl survey, and more complex assessment model structure should be considered to model seasonal availability and selectivity (potentially interacting with density).
- Possibly develop a spatially structured model that includes SNA 7 and relevant parts of SNA 8.
- Develop an overfishing threshold that is higher than the exploitation rate target.

#### Stock structure

- Continue to review available data and explore additional data sources to inform stock structure assumptions.

#### Harvest Control Rule

- Develop a harvest control rule that requires the exploitation rate to decline below the target as spawning biomass approaches the soft limit, e.g., akin to the Harvest Strategy Standard default.

#### Environmental drivers of recruitment

- Further investigation should be conducted to identify correlations between snapper recruitment and key environmental variables to improve our understanding of snapper recruitment dynamics.

## Mean age at length

- Evaluate fit of VB functions to age data and if there are issues consider alternative means of representing mean age at length.

## Tagging study

- Undertake a tagging study (in association with SNA 8) to estimate biomass, seasonal movement, stock boundaries, and selectivity.

## 5. STATUS OF THE STOCKS


### Stock Structure Assumptions

New Zealand snapper are thought to comprise either seven or eight biological stocks based on the location of spawning and nursery grounds, differences in growth rates, age structure, and recruitment strength, and the results of tagging studies. These stocks are assumed to comprise three in SNA 1 (East Northland, Hauraki Gulf, and Bay of Plenty), two in SNA 2 (one of which may be associated with the Bay of Plenty stock), two in SNA 7 (Marlborough Sounds and Tasman/Golden Bay/west coast South Island), and one in SNA 8. Tagging studies reveal that limited mixing occurs between the three SNA 1 biological stocks, with the greatest exchange between the Bay of Plenty and Hauraki Gulf.

- **SNA 7**

The assessment is for the Tasman Bay, Golden Bay, and west coast South Island stock unit of SNA 7. The Marlborough Sounds is considered to support a separate stock of snapper within SNA 7, but catches, although minor, are included in the SNA 7 assessment model.

Stock Status		
Most Recent Assessment Plenary Publication Year	2024	
Intrinsic Productivity Level	Low	
Catch in most recent year of assessment	Year: 2023–24	Catch: 708 t (TACC 450 t, additional mortality 45 t, <i>F</i> based recreational estimate 213 t)
Assessment Runs Presented	Base case model	
Reference Points	Interim target ; $U_{SB40\%} = 5\%$ Soft Limit: Twice the biomass of hard limit. Hard Limit: Twice 1987 spawning biomass estimate. Overfishing threshold: $U_{SB40\%}$	
Status in relation to Target	Very Likely (> 90%) to be at or below the target.	
Status in relation to Limits	Soft Limit: Exceptionally Unlikely (< 1%) to be below Hard Limit: Exceptionally Unlikely (< 1%) to be below	
Status in relation to Overfishing	Overfishing is Very Unlikely (< 10%) to be occurring	

**Historical Stock Status Trajectory and Current Status**

Annual exploitation rate (relative to  $USB_{40\%}$ ) (left) and base model  $SSB$  trajectory (right) for the period since 1975 (dotted line indicates target  $USB_{40\%}$  exploitation rate). Projections ( $RecF$ ) are in red. The line represents the median and the shaded area represents the 95% credible interval. The red and orange dashed lines represent the hard and soft limits, respectively.

**Fisheries and Stock Trends**

Recent Trend in Biomass or Proxy	Biomass was at an historical low level in the early 2000s and increased substantially from 2009, initially due to the recruitment of several strong year classes. Biomass has continued to increase following recruitment of recent (2017 and 2018) very strong year classes.
Recent Trend in Fishing Intensity or Proxy	Fishing mortality declined steadily from 2006 to 2015, and has remained well below the overfishing threshold since then.
Other Abundance Indices	<ul style="list-style-type: none"> <li>- The Spring/Summer CPUE series (BT1), although assumed to be biased low and not used in the assessment after 2010, also shows a substantial increase.</li> <li>- CPUE from the west coast South Island trawl fishery has increased substantially over the last 10 years.</li> </ul>
Trends in Other Relevant Indicators or Variables	<ul style="list-style-type: none"> <li>- The increase in recreational catch estimates from 2005 onwards suggests that abundance has increased.</li> <li>- Increased abundance of snapper in areas adjacent to SNA 7, i.e South Taranaki Bight.</li> </ul>

**Projections and Prognosis**

Stock Projections or Prognosis	Projections (5 yr) are provided based on recent (2010–11 to 2019–20) average recruitment. Biomass is projected to continue to increase at the level of the current TACC and projected recreational catch.
Probability of Current Catch or TAC causing Biomass to remain below or to decline below Limits	Soft Limit: Exceptionally Unlikely (< 1%) Hard Limit: Exceptionally Unlikely (< 1%)
Probability of Current Catch or TAC causing Overfishing to continue or to commence	Exceptionally Unlikely (< 1%)

<b>Assessment Methodology and Evaluation</b>		
Assessment Type	Level 1 – Full Quantitative Stock Assessment	
Assessment Method	Age-structured Stock Synthesis model with MCMC estimation	
Assessment Dates	Latest assessment Plenary publication year: 2024	Next assessment: 2026
Overall assessment quality rank	1 – High Quality	
Main data inputs (rank)	<ul style="list-style-type: none"> <li>- Commercial catch history (1983 onwards)</li> <li>- Commercial catch history (1975-1983)</li> <li>Tagging biomass estimate</li> <li>- CPUE indices</li> <li>- Historical commercial age frequency</li> <li>- Recent commercial age frequency</li> <li>- Recreational catch history (2005 onwards)</li> <li>- Recreational catch history (preceding period)</li> <li>- <i>Kaharoa</i> WCSI trawl survey biomass indices (core area)</li> <li>-Trawl survey age compositions (2016, 2018, 2020, 2022)</li> <li>-Trawl survey length compositions (2008–2016)</li> </ul>	<ul style="list-style-type: none"> <li>1 – High Quality</li> <li>2 – Medium or Mixed Quality: catches are considered to be less reliable</li> <li>2 – Medium or Mixed Quality: whether the older ages are indexed by the tagging study is uncertain</li> <li>1 – High Quality</li> <li>2 – Medium Quality: needs to be better characterised by method of capture</li> <li>1 – High Quality</li> <li>1 – High quality</li> <li>2 – Medium or Mixed Quality: historical levels of recreational catch are assumed</li> <li>1 – High Quality</li> <li>1 – High Quality</li> <li>1 – High Quality</li> </ul>
Data not used (rank)	BT1 (October to December) CPUE index post 2010  West coast inshore bottom trawl CPUE	3 – Low Quality: potentially biased low due to recent avoidance of spawning aggregations 3 – Low Quality: potentially hypostable
Changes to Model Structure and Assumptions	<ul style="list-style-type: none"> <li>- Initialise model in 1975 under exploited conditions, including prior on initial equilibrium fishing mortality and initialising recruitment deviates (1960–1974).</li> <li>- Recruitment deviates estimated (from 1975) as simple deviates (i.e. not constrained to an average of one).</li> <li>- Updated recreational catch history incorporating recent recreational catch estimate (2022–23). Alternative assumptions for future recreational catches.</li> <li>- Increased weighting on early (1970s, early 1980s) BPT age compositions.</li> <li>- Increased age-at-maturity based on ovarian sampling from WCNI trawl survey. Individual sample sizes for each observation of BT1 and BT2 age composition series.</li> <li>- Parameterisation of trawl survey selectivity (time varying logistic).</li> </ul>	

Major Sources of Uncertainty	- Process error included for trawl survey biomass indices.
	- Strength of recent recruitment (2017 and 2018 year classes) - Reliability of CPUE indices, as could be biased low due to snapper avoidance since 2015. - Historical and projected levels of recreational catch - Availability of older (10+ yr) snapper to the trawl survey - Connectivity between SNA 7 and southern SNA 8

**Qualifying Comments**

The stock structure relationship with the South Taranaki Bight portion of SNA 8 is unclear.

**Fisheries Interactions**

Snapper target fisheries have a bycatch of flatfish, red cod, gurnard, tarakihi, and small amounts of barracouta and blue warehou. Snapper is taken as a bycatch of the inshore trawl fisheries operating within FMA 7, particularly within Tasman Bay and Golden Bay. Since 2013–14, most (> 80%) of the snapper catch has been taken as a bycatch of those fisheries.

**6. FOR FURTHER INFORMATION**

- Annala, J H; Sullivan, K J (Comps.) (1997) Report from the Fishery Assessment Plenary, May 1997: stock assessments and yield estimates. 381 p. (Unpublished report held by NIWA library, Wellington.)
- Bentley, N; Kendrick, T H (2015) The inshore fisheries of the Central (East) fisheries management area (FMA2): characterisation and catch-per-unit-effort analyses, 1989–90 to 2009–10 Draft New Zealand Fisheries Assessment Report for Research Project INS2009/03. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bernal-Ramírez, J H; Adcock, G J; Hauser, L; Carvalho, G R; Smith, P J (2003) Temporal stability of genetic population structure in the New Zealand snapper, *Pagrus auratus*, and relationship to coastal currents. *Marine Biology* 142(3): 567–574.
- Bian, R; McKenzie, J R; Davies, N M (2009) Determination of optimum frequency for SNA 8 snapper market sampling based on retrospective analysis. *New Zealand Fisheries Assessment Report* 2009/50. 15 p.
- Blackwell, R G; Gilbert, D J (2006) Age composition of commercial snapper landings in SNA 2, 2004–05. *New Zealand Fisheries Assessment Report* 2006/46. 18 p.
- Blackwell, R G; McKenzie, J R (2013) Age composition of commercial snapper landings in SNA 2, 2007–08. *New Zealand Fisheries Assessment Report* 2013/25. 32 p.
- Blackwell, R G; Stevenson, M L (1997) Trawl survey of juvenile snapper in Tasman and Golden Bays, July 1996 (KAH9608). *New Zealand Fisheries Data Report No.* 87.12 p.
- Boyd, R O; Gowing, L; Reilly, J L (2004) 2000–2001 national marine recreational fishing survey: diary results and harvest estimates. . Final Research Report for Ministry of Fisheries. (Unpublished report held by Fisheries New Zealand, Wellington.) 93 p.
- Boyd, R O; Reilly, J L (2002) 1999/2000 National marine recreational fishing survey: harvest estimates. Final Research Report for Ministry of Fisheries Research Project REC9803. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Bradford, E (1998) Harvest estimates from the 1996 national marine fishing surveys. New Zealand Fisheries Assessment Research Document 1998/16. 27 p. (Unpublished document held by NIWA library, Wellington.)
- Brothers, N; Duckworth, A R; Safina, C; Gilman, E L (2010) Seabird bycatch in pelagic longline fisheries is grossly underestimated when using only haul data. *PloS ONE* 5: e12491. doi: 10.1371/journal.pone.001249
- Bull, B; Francis, R I C C; Dunn, A; Gilbert, D J; Bian, R; Fu, D (2012) CASAL (C++ algorithmic stock assessment laboratory): CASAL User Manual v2.30.2012/03/21. *NIWA Technical Report* 135. 280 p.
- Bull, B; Francis, R I C C; Dunn, A; McKenzie, A; Gilbert, D J; Smith, M H (2004) CASAL (C++ algorithmic stock assessment laboratory): CASAL User Manual v2.06-2004/09/26. *NIWA Technical Report* 126. 261 p.
- Davies, N M (1997) Assessment of the west coast snapper (*Pagrus auratus*) stock (SNA 8) for the 1996–97 fishing year. New Zealand Fisheries Assessment Research Document 1997/12. 47 p. (Unpublished document held by NIWA library, Wellington.)
- Davies, N M (1999) Assessment of the SNA 1 and 8 stocks for the 1997–98 fishing year. New Zealand Fisheries Assessment Research Document 1999/19. 87 p. (Unpublished document held by NIWA library, Wellington.)
- Davies, N M; Gilbert, D J; McKenzie, J R (1999a) Assessment of the SNA 1 and 8 stocks for the 1998–99 fishing year. New Zealand Fisheries Assessment Research Document 1999/28. 82 p. (Unpublished document held by NIWA library, Wellington.)
- Davies, N M; McKenzie, J R; Gilbert, D J (2006) Assessment of the SNA 8 stock for the 2003–04 fishing year. *New Zealand Fisheries Assessment Report* 2006/9. 32 p.
- Davies, N M; McKenzie, J R; Gilbert, D J (2013) Assessment of the SNA 8 stock for the 2004–05 fishing year. *New Zealand Fisheries Assessment Report* 2013/28. 73 p.
- Davies, N M; McKenzie, J R; Gilbert, D J (1999b) Monte Carlo estimation of bias in Petersen mark-recapture estimates for snapper (*Pagrus auratus*). New Zealand Fisheries Assessment Research Document 1999/20. 52 p. (Unpublished document held by NIWA library, Wellington.)
- Davies, N M; Walsh, C; Hartill, B (1993) Estimating catch at age of snapper from west coast and Hauraki Gulf fisheries, 1992–93. Northern Fisheries Region Internal Report No. 17. 58 p. (Unpublished report held by Fisheries New Zealand, Auckland.)
- Dunn, M R; Hurst, R J; Renwick J; Francis, R I C C; Devine, J; McKenzie, A (2009) Fish abundance and climate trends in New Zealand. *New Zealand Aquatic Environment and Biodiversity Report No.* 31. 73 p.
- Fielder, D S; Bardsley, W J; Allan, G L; Pankhurst, P M (2005) The effects of salinity and temperature on growth and survival of Australian snapper, *Pagrus auratus* larvae. *Aquaculture* 250(1–2): 201–214.
- Fowler, A J; Jennings, P R (2003) Dynamics in 0+ recruitment and early life history for snapper (*Pagrus auratus*, Sparidae) in South Australia. *Marine and Freshwater Research* 54(8): 941–956.

- Francis, M P (1993) Does water temperature determine year class strength in New Zealand snapper (*Pagrus auratus*, Sparidae)? *Fisheries Oceanography* 2(2): 65–72.
- Francis, M P (1994) Growth of juvenile snapper, *Pagrus auratus*. *New Zealand Journal of Marine and Freshwater Research* 28(2): 201–218.
- Francis, M P; Evans, J (1992) Immigration of subtropical and tropical animals into north-eastern New Zealand. Paper presented at the Proceedings of the Second International Temperate Reef Symposium.
- Francis, M P; Langley, A; Gilbert, D (1997) Prediction of snapper (*Pagrus auratus*) recruitment from sea surface temperature. In: Hancock, D A; Smith, D C; Grant, A; Beumer J P (Eds) *Developing and sustaining world fisheries resources: the state of science and management*, pp. 67–71. 2<sup>nd</sup> World Fisheries Congress 28 Jul–2 Aug 1996, Brisbane, Australia. CSIRO Publishing.
- Francis, M P; Langley, A D; Gilbert, D J (1995) Snapper recruitment in the Hauraki Gulf. New Zealand Fisheries Assessment Research Document 1995/17. 26 p. (Unpublished document held by NIWA library, Wellington.)
- Francis, M P; Lyon, W S (2012) Review of commercial fishery interactions and population information for eight New Zealand protected fish species. (Unpublished NIWA client report WLG2012-64 prepared for the Department of Conservation, Wellington.) 67 p. Available at <https://www.doc.govt.nz/Documents/conservation/marine-and-coastal/marine-conservation-services/pop2011-03-protected-fish-review.pdf>
- Francis, M P; Paul, L J (2013) New Zealand inshore finfish and shellfish commercial landings, 1931–82. *New Zealand Fisheries Assessment Report 2013/55*. 136 p.
- Francis, R I C C (2011) Data weighting in statistical fisheries stock assessment models *Canadian Journal of Fisheries and Aquatic Sciences*. 68: 1124–1138.
- Francis, R I C C; McKenzie, J R (2015a) Assessment of the SNA 1 stocks in 2012. *New Zealand Fisheries Assessment Report 2015/75*.
- Francis, R I C C; McKenzie, J R (2015b) Assessment of the SNA 1 stocks in 2013. *New Zealand Fisheries Assessment Report 2015/76*.
- Froese, R; Pauly, D (2000) FishBase 2000: concepts, design and data sources. ICLARM, Los Banos, Laguna, Philippines. 344 p.
- Gilbert, D J (1994) A total catch history model for SNA 1. New Zealand Fisheries Assessment Research Document 1994/24. 16 p. (Unpublished document held by NIWA library, Wellington.)
- Gilbert, D J; McKenzie, J R (1999) Sources of bias in biomass estimates from tagging programmes in the SNA 1 snapper (*Pagrus auratus*) stock. New Zealand Fisheries Assessment Research Document 1999/16. 47 p. (Unpublished document held by NIWA library.)
- Gilbert, D J; McKenzie, J R; Davies, N M; Field, K D (2000) Assessment of the SNA 1 stocks for the 1999–2000 fishing year. *New Zealand Fisheries Assessment Report 2000/38*. 52 p.
- Gilbert, D J; Phillips, N L (2003) Assessment of the SNA 2 and Tasman and Golden Bays (SNA 7) snapper fisheries for the 2001–02 fishing year. *New Zealand Fisheries Assessment Report 2003/45*.
- Gilbert, D J; Sullivan, K J (1994) Stock assessment of snapper for the 1992–93 fishing year. New Zealand Fisheries Assessment Research Document 1994/3. 37 p. (Unpublished document held by NIWA library, Wellington.)
- Gilbert, D J; Sullivan, K J; Davies, N M; McKenzie, J R; Francis, M P; Stan, P J (1996) Population modelling of the SNA 1 stock for the 1995–96 fishing year. New Zealand Fisheries Assessment Research Document 96/15. 39 p. (Unpublished document held by NIWA library, Wellington.)
- Gilbert, D J; Taylor, P R (2001) The relationships between snapper (*Pagrus auratus*) year class strength and temperature for SNA 2 and SNA 7. *New Zealand Fisheries Assessment Report 2001/64*. 33 p.
- Godfriaux, B L (1969) Food of predatory demersal fish in Hauraki Gulf. 1. Food and Feeding habitats of snapper. *New Zealand Journal of Marine and Freshwater Research* 3: 518–544.
- Godfriaux, B L (1970) Food of predatory demersal fish in Hauraki Gulf. 3. Feeding relationships. *New Zealand Journal of Marine and Freshwater Research* 4: 325–336.
- Godfriaux, B L (1974) Feeding relationships between tarakihi and snapper in western Bay of Plenty, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 8: 589–609.
- Hamer, P A; Acevedo, S; Jenkins, G P; Newman, A (2011) Connectivity of a large embayment and coastal fishery: spawning aggregations in one bay source local and broadscale fishery replenishment. *Journal of Fish Biology* 78: 1090–1109.
- Harley, S J; Gilbert, D J (2000) Assessment of the Tasman and Golden Bays snapper fishery for the 1999–2000 fishing year. *New Zealand Fisheries Assessment Report 2000/28*. 42 p.
- Hartill, B; Bian, R; Armiger, H; Vaughan, M; Rush, N (2007) Recreational marine harvest estimates of snapper, kahawai, and kingfish in QMA 1 in 2004–05. *New Zealand Fisheries Assessment Report 2007/26*. 44 p.
- Hartill, B; Bian, R; Rush, N; Armiger, H (2013) Aerial-access recreational harvest estimates for snapper, kahawai, red gurnard, tarakihi and trevally in FMA 1 in 2011–12. *New Zealand Fisheries Assessment Report 2013/70*. 49 p.
- Hartill, B; Bian, R; Rush, N; Armiger, H (2019) Aerial-access recreational harvest estimates for snapper, kahawai, red gurnard, tarakihi and trevally in FMA 1 in 2017–18. *New Zealand Fisheries Assessment Report 2019/23*. 39 p.
- Hartill, B; Sutton, C (2011) Characterisation and catch per unit effort indices for the SNA 7 fishery. *New Zealand Fisheries Assessment Report 2011/53*. 55 p.
- Hartill, B; Watson, T; Cryer, M; Armiger, H (2007) Recreational marine harvest estimates of snapper and kahawai in the Hauraki Gulf in 2003–04. *New Zealand Fisheries Assessment Report 2007/25*. 55 p.
- Heinemann, A; Gray, A. (2024) National Panel Survey of Marine Recreational Fishers 2022–23. *New Zealand Fisheries Assessment Report 2024/51*. 122 p.
- Holdsworth, J C; Boyd, R O (2008) Size, condition and estimated release mortality of snapper (*Pagrus auratus*) caught in the SNA 1 recreational fishery, 2006–07. *New Zealand Fisheries Assessment Report 2008/53*. 37 p.
- Hurst, R J; Stevenson, M L; Bagley, N W; Griggs, L H; Morrison, M A; Francis, M P (2000) Areas of importance for spawning, pupping or egg-laying, and juveniles of New Zealand coastal fish. Final Research Report to the Ministry of Fisheries Research Project ENV1999-03. 271 p. (Unpublished draft NIWA Technical Report available at <https://fs.fish.govt.nz/Doc/22534/ENV1999-03%20Coastal%20Fish%20NZ%20Objective%201%20Final.pdf.ashx>.)
- Jones, E; Morrison, M; Parsons, D M; Paterson, C; Usmar, N; Bagley, N (2010) Fish communities (Chapter 13). Oceans 2020 Bay of Islands Survey report to LINZ prepared by NIWA. 98 p.
- Kendrick, T H; Bentley, N (2010) Fishery characterisation and catch-per-unit-effort indices for snapper in SNA 8, 1989–90 to 2007–08. *New Zealand Fisheries Assessment Report 2010/42*.
- Kendrick, T H; Francis, M P (2002) Fish assemblages in the Hauraki Gulf, New Zealand. *New Zealand Journal of Marine and Freshwater Research* 36: 699–717.
- King, M R (1985) Fish and shellfish landings by domestic fishermen, 1974–82. *Fisheries Research Division Occasional Publication: Data Series* 20. 96 p.
- King, M R (1986) Catch statistics for foreign and domestic commercial fishing in New Zealand waters, January–December, 1983. *Fisheries Research Division Occasional Publication: Data series* 21. 140 p.
- King, M R; Jones, D M; Fisher, K A; Sanders, B M (1987) Catch statistics for foreign and domestic commercial fishing in New Zealand waters, January – December 1984. *New Zealand Fisheries Data Report No. 30*. 150 p.
- Kirk, P D; Drummond, K L; Ryan, M (1988) Preliminary stock size analysis: Tasman/Golden Bay snapper tagging programme. New Zealand Fisheries Assessment Research Document 88/44. (Unpublished report held by Fisheries New Zealand, Wellington.)



- Langley, A D (1995a) Trawl survey of snapper and associated species in the Hauraki Gulf, October–November 1994 (KAH9411). *New Zealand Fisheries Data Report No. 61*. 35 p.
- Langley, A D (1995b) Trawl survey of snapper and associated species off the west coast of the North Island, October 1994 (KAH9410). *New Zealand Fisheries Data Report No. 65*. 35 p.
- Langley, A D (2010) Stock assessment of SNA 2 for 2010. *New Zealand Fisheries Assessment Report 2010/26*.
- Langley, A D (2013) An update of the analysis of SNA 7 trawl CPUE indices and other recent data from the SNA 7 fishery. *New Zealand Fisheries Assessment Report 2013/17*. 46 p.
- Langley, A D (2014) Updated CPUE analyses for selected South Island inshore finfish. *New Zealand Fisheries Assessment Report 2014/40*. 116 p.
- Langley, A D (2015) Stock assessment of SNA 7 for 2015. *New Zealand Fisheries Assessment Report 2015/42*. 67 p.
- Langley, A D (2016) An update of the analysis of longline Catch-Per-Unit-Effort indices for snapper in SNA 1. *New Zealand Fisheries Assessment Report 2016/17*. 71 p.
- Langley, A D (2017) Catch-Per-Unit-Effort indices for snapper in SNA 8. *New Zealand Fisheries Assessment Report 2017/45*. 40 p.
- Langley, A D (2018) Stock assessment of snapper in SNA 7. *New Zealand Fisheries Assessment Report 2018/25*. 67 p.
- Langley, A D (2020a) An update of the stock assessment of snapper in SNA 7. *New Zealand Fisheries Assessment Report 2020/09*. 67 p.
- Langley, A D (2020b) Stock assessment of snapper in SNA 8 for 2020. *New Zealand Fisheries Assessment Report 2020/20*. 87 p.
- Langley, A D (2021a) Stock assessment of snapper in SNA 7 for 2021. *New Zealand Fisheries Assessment Report 2021/83*. 93 p.
- Langley, A D (2021b) Stock assessment of snapper in SNA 8 for 2021. *New Zealand Fisheries Assessment Report 2021/38*. 86 p.
- Martino, J C; Fowler, A; Doubleday, Z A; Grammer, G L; Gillanders, B M (2019) Using otolith chronologies to understand long-term trends and extrinsic drivers of growth in fisheries. *Ecosphere 10(1)*: e02553.
- MacGibbon, D J (2019) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2019 (KAH1902). *New Zealand Fisheries Assessment Report 2019/64*. 87 p.
- MacGibbon, D J; Walsh, C; Buckthought, D; Bian, R (2022) Inshore trawl survey off the west coast South Island and in Tasman Bay and Golden Bay, March–April 2021 (KAH2103). *New Zealand Fisheries Assessment Report 2022/11*. 97 p.
- MacGibbon, D J; Walsh, C; Buckthought, D; Bian, R (2024) Inshore trawl survey off the west coast South Island and in Tasman Bay and Golden Bay, March–April 2023 (KAH2302). *New Zealand Fisheries Assessment Report 2024/06*. 91 p.
- Maunder, M N; Starr, P J (1995) Validating the Hauraki Gulf snapper pre-recruit trawl surveys and temperature recruitment relationship using catch at age analysis with auxiliary information. New Zealand Fisheries Assessment Research Document 1998/15. (Unpublished document held by NIWA library, Wellington.)
- McKenzie, J R (2000) Factors Affecting Mortality of small Snapper (*Pagrus auratus*) caught and released by the SNA 1 Longline Fishery. (Draft Fisheries Assessment Report held by NIWA Library, Wellington).
- McKenzie, J R (2012) An evaluation of a fully age-structured spatially disaggregated stock assessment model for the SNA 1 QMA. *New Zealand Fisheries Assessment Report 2012/38*. 120 p.
- McKenzie, J R; Diggles, B; Tubbs, L; Poortenaar, C; Parkinson, D; Webster, K; Miller, N (2006) An evaluation of a new type of plastic-coated PIT tag for tagging snapper (*Pagrus auratus*). *New Zealand Fisheries Assessment Report 2006/8*. 40 p.
- McKenzie, J R; Parsons, D M (2012) Fishery characterisations and catch-per-unit-effort indices for three sub-stocks of snapper SNA 1, 1989–90 to 2009–10. *New Zealand Fisheries Assessment Report 2012/29*. 112 p.
- McMahon, S J; Parsons, D M; Donelson, J M; Pether, S M; Munday, P L (2020a) Elevated CO<sub>2</sub> and heatwave conditions affect the aerobic and swimming performance of juvenile Australasian snapper. *Marine Biology 167(1)*: 1–12.
- McMahon, S J; Parsons, D M; Donelson, J M; Pether, S M; Munday, P L (2020b) Elevated temperature and CO<sub>2</sub> have positive effects on the growth and survival of larval Australasian snapper. *Marine Environmental Research 161*: 105054.
- Methot, R D (1990) Synthesis model: an adaptable framework for analysis of diverse stock assessment data. *International North Pacific Fisheries Commission Bulletin 50*: 259–275.
- Methot, R D (2005) Technical description of the Stock Synthesis II assessment program. NOAA Technical Memorandum SEDAR 16-AW-04.
- Methot, R D (2009) User manual for Stock Synthesis, model version 3.02C.
- Methot, R D; Wetzel, C R (2013) Stock Synthesis: A biological and statistical framework for fish stock assessment and fishery management. *Fisheries Research 142*: 86–99.
- Middleton, D A J; Abraham, E R (2007) The efficacy of warp strike mitigation devices: Trials in the 2006 squid fishery. Final Research Report for research project IPA2006/02. (Unpublished report held by Fisheries New Zealand, Wellington).
- Millar, R B; Akroyd, J M; Walshe, K A R (2001) Incidental mortality of snapper in SNA 1 and SNA 8. *New Zealand Fisheries Assessment Report 2001/78*. 36 p.
- Ministry of Fisheries (2008) Harvest Strategy Standard for New Zealand Fisheries. 25 p. Available online at: <https://fs.fish.govt.nz/Doc/16543/harveststrategyfinal.pdf.ashx>
- Morrison, M (1997) Trawl survey of snapper and associated species in the Bay of Plenty, February 1996 (KAH9601). *NIWA Technical Report 2*. 33 p.
- Morrison, M A (1998) Trawl survey of snapper and associated species off the west coast of the North Island, November 1996 (KAH9615). *NIWA Technical Report 33*. 48 p.
- Morrison, M A; Francis, M P (1997) Trawl survey of snapper and associated species in the Hauraki Gulf, October–November 1997 (KAH9720). *NIWA Technical Report 58*. 37 p.
- Morrison, M A; Jones, E G; Consalvey, M; Berkenbusch, K (2014a) Linking marine fisheries species to biogenic habitats in New Zealand: a review and synthesis of knowledge. *New Zealand Aquatic Environment and Biodiversity Report No. 130*. 156 p.
- Morrison, M A; Parkinson, D M (2001) Trawl survey of snapper and associated species off the west coast of the North Island, November 1999 (KAH9915). *NIWA Technical Report 100*. 51 p.
- Morrison, M A; Lowe, M L; Parsons, D M; Usmar, N R; McLeod, I M (2009) A review of land-based effects on coastal fisheries and supporting biodiversity in New Zealand. *New Zealand Aquatic Environment and Biodiversity Report No. 37*. 100 p.
- Morrison, M A; Lowe, M L; Grant, C M; Smith, P J; Carbines, G; Reed, J; Bury, S J; Brown, J (2014b) Seagrass meadows as biodiversity and productivity hotspots. *New Zealand Aquatic Environment and Biodiversity Report No. 137*. 147 p.
- Morrison, M A; Stevenson, M L (2001) Review of west coast North Island trawl survey time series, 1986–96. *NIWA Technical Report 107*. 55 p.
- Murphy, H M; Jenkins, G P; Hamer, P A; Swearer, S E (2013) Interannual variation in larval abundance and growth in snapper *Chrysophrys auratus* (Sparidae) is related to prey availability and temperature. *Marine Ecology Progress Series 487*: 151–162.
- Parker, S J; Parsons, D; Stevenson, M; Sutton, C; Walsh, C (2015) Landed catch sampling of snapper in SNA 7 in the 2013–14 fishing year. *New Zealand Fisheries Assessment Report 2015/61*. 21 p.
- Parsons, D M; Bian, R (in prep) Trawl surveys of the Hauraki Gulf and Bay of Plenty in 2019, 2020 and 2021 to estimate the abundance of juvenile snapper. Draft New Zealand Fisheries Assessment Report.
- Parsons, D M; Bian, R; McKenzie, J R; McMahon, S J; Pether, S; Munday, P L (2020) An uncertain future: Effects of ocean acidification and elevated temperature on a New Zealand snapper (*Chrysophrys auratus*) population. *Marine Environmental Research 161*: 105089.

- Parsons, D M; Morrison, M A; MacDiarmid, A B; Stirling, B; Cleaver, P; Smith, I W G; Butcher, M (2009) Risks of shifting baselines highlighted by anecdotal accounts of New Zealand's snapper (*Pagrus auratus*) fishery. *New Zealand Journal of Marine and Freshwater Research* 43: 965–983.
- Parsons, D M; Parker, S J; Stevenson, M; Sutton, C; Buckthought, D; Bian, R; McKenzie, J; Walsh, C (2018) Catch at-age of snapper in SNA 7 in the 2016–17 fishing year. *New Zealand Fisheries Assessment Report 2018/03*. 23 p.
- Paul, L J (1976) A study on age, growth and population structure of the snapper, *Chrysophrys auratus* in Hauraki Gulf. *New Zealand Fisheries Research Bulletin No. 13*. 63 p.
- Paul, L J (1977) The commercial fishery for snapper *Chrysophrys (Pagrus) auratus* in the Auckland region, New Zealand, from 1900 to 1971. *Fisheries Research Division Bulletin No 15*. 84 p.
- Powell, A W B (1937) Animal Communities of the Sea-bottom in Auckland and Manukau Harbours. *Transactions and Proceedings of the Royal Society of New Zealand* 66: 354–401.
- Ritchie, L; Saul, P; O'Sullivan, K. (1975) The wetfish report 1941–1970. *New Zealand Ministry of Agriculture and Fisheries Fisheries Technical Report 137*. 370 p.
- Schofield, M I; Langley, A D; Bentley, N; Middleton, D A J (2018a) Catch-per unit-effort (CPUE) analyses for SNA 2. *New Zealand Fisheries Assessment Report 2018/15*. 87 p.
- Schofield, M I; Langley, A D; Middleton, D A J (2018b) Catch-per unit-effort (CPUE) update for FMA 2 snapper (SNA 2). Report for Fisheries Inshore New Zealand. [https://www.inshore.co.nz/fileadmin/Documents/Science/SNA2\\_rapidCPUEupdate\\_2018.pdf](https://www.inshore.co.nz/fileadmin/Documents/Science/SNA2_rapidCPUEupdate_2018.pdf)
- Stevenson, M L; Hanchet, S M (2010) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2009. *New Zealand Fisheries Assessment Report 2010/11*. 77 p.
- Stevenson, M L; MacGibbon, D J (2018) Inshore trawl survey of the west coast South Island and Tasman and Golden Bays, March–April 2017 (KAH1703) *New Zealand Fisheries Assessment Report 2018/18*. 92 p.
- Sullivan, K J (1985) Snapper. In: Colman, J A; McKoy, J L; Baird, G G (Comps. and Eds.) (1985) Background papers for the 1985 Total Allowable Catch recommendations, pp. 187–214. (Unpublished report, held in NIWA library, Wellington.)
- Sullivan, K J; Hore, A J; Wilkinson, V H (1988) Snapper. In: Baird, G G; McKoy, J L Papers from the workshop to review fish stock assessments for the 1987–88 New Zealand fishing year, pp. 251–275. (Unpublished report, held in NIWA library, Wellington.)
- Sylvester, T (1995) Initial results of the Northern boat ramp survey. *Seafood News Zealand*, February 1995. pp. 11–13.
- Teirney, L D; Kilner, A R; Millar, R B; Bradford, E; Bell, J D (1997) Estimation of recreational harvests from 1991–92 to 1993–94. *New Zealand Fisheries Assessment Research Document 1997/15*. 43 p. (Unpublished document held by NIWA library, Wellington.)
- Vignaux, M (1993) Catch per unit of effort (CPUE) analysis of the SNA 8 snapper fishery. *New Zealand Fisheries Assessment Research Document 1993/2*. 12 p. (Unpublished document held by NIWA library, Wellington.)
- Wakefield, C B; Potter, I C; Hall, N G; Lenanton, R C; Hesp, S A (2017) Timing of growth zone formations in otoliths of the snapper, *Chrysophrys auratus*, in subtropical and temperate waters differ and growth follows a parabolic relationship with latitude. *ICES Journal of Marine Science* 74(1): 180–192.
- Walsh, C; Buckthought, D; Bian, R; Parsons, D; Patke, S; Smith, M (2019) Catch-at-age of snapper otoliths from 2019 West Coast South Island trawl survey. (Final Research Report for Fisheries New Zealand for Project SEA2019-05 held by Fisheries New Zealand.) 12 p.
- Walsh, C; Armiger, H; Bian, R; Buckthought, D; McKenzie, J (2017) Length and age composition of commercial snapper landings in SNA 8, 2015–16. *New Zealand Fisheries Assessment Report 2017/02*. 40 p.
- Walsh, C; Davies, N M (2004) Length and age composition of commercial landings in SNA 8, 2003–04. *New Zealand Fisheries Assessment Report 2004/56*. 18 p.
- Walsh, C; Davies, N M; Buckthought, D (2006a) Length and age composition of commercial snapper landings in SNA 8, 2005–06. *New Zealand Fisheries Assessment Report 2006/54*. 21 p.
- Walsh, C; Davies, N M; Rush, N; Buckthought, D; Smith, M (2006b) Age composition of commercial snapper landings in SNA 1, 2004–05. *New Zealand Fisheries Assessment Report 2006/39*. 34 p.
- Walsh, C; Davies, N M; Rush, N; Buckthought, D; Vaughn, M; Smith, M (2007) Length and age composition of commercial snapper landings in SNA 1, 2005–06. *New Zealand Fisheries Assessment Report 2007/01*. 30 p.
- Walsh, C; Davies, N M; Rush, N; Middleton, C; Smith, M; Newmarch, G (2006c) Length and age composition of commercial snapper landings in SNA 1, 2003–04. *New Zealand Fisheries Assessment Report 2006/7*. 46 p.
- Walsh, C; McKenzie, J; Arminger, H (2006d) Spatial and temporal patterns in snapper length and age composition and movement, west coast North Island, New Zealand. *New Zealand Fisheries Assessment Report 2006/6*. 59 p.
- Walsh, C; McKenzie, J M; Bian, R; Armiger, H; O'Maolagáin, C; Buckthought, D; Smith, M; Ferguson, H; Miller A (2012) Snapper catch-at-length and catch-at-age heterogeneity between spatial strata in SNA 2 bottom trawl landings, 2007–08 and 2008–09. *New Zealand Fisheries Assessment Report 2012/40*. 44 p.
- Walsh, C; McKenzie, J A; Buckthought, D; Armiger, H; Ferguson, H; Smith, M; Spong, K; Miller, A (2011) Age composition of commercial snapper landings in SNA 1, 2009–10. *New Zealand Fisheries Assessment Report 2011/54*.
- Wright, P; McClary, D; Boyd, R O (2004) 2000/2001 National Marine Recreational Fishing Survey: direct questioning of fishers compared with reported diary data. Final Research Report for Ministry of Fisheries Project REC2000-01: Objective 2. (Unpublished report held by Fisheries New Zealand, Wellington.)
- Wynne-Jones, J; Gray, A; Heinemann, A; Hill, L; Walton, L (2019) National Panel Survey of Marine Recreational Fishers 2017–2018. *New Zealand Fisheries Assessment Report 2019/24*. 104 p.
- Wynne-Jones, J; Gray, A; Hill, L; Heinemann, A (2014) National Panel Survey of Marine Recreational Fishers 2011–12: Harvest Estimates. *New Zealand Fisheries Assessment Report 2014/67*. 139 p.
- Zeldis, J R; Oldman, J; Ballara, S L; Richards, L A (2005) Physical fluxes, pelagic ecosystem structure, and larval fish survival in Hauraki Gulf, New Zealand. *Canadian Journal of Fisheries and Aquatic Sciences* 62(3): 593–610.
- Zeldis, J R; Francis, R I C C (1998) A daily egg production method estimate of snapper biomass in Hauraki Gulf, New Zealand. *ICES Journal of Marine Science* 55: 522–534.