

COCKLES (COC)

(*Austrovenus stutchburyi*)
Tuangi

1. INTRODUCTION

Commercial picking of cockles, *Austrovenus stutchburyi*, is carried out on Snake Bank, Whangarei Harbour (FMA 1), Pakawau Beach, Ferry Point and Tapu Bay in Tasman and Golden Bays (FMA 7), and Papanui and Waitati Inlets, Otago (FMA 3). Cockles were introduced into the QMS on 1 October 2002.

For assessment purposes, individual reports on the major fisheries have been produced separately:

1. Snake Bank, Whangarei Harbour, in COC 1A.
2. Papanui Inlet, Waitati Inlet, Purakanui and Otago Harbour, Otago Peninsula in COC 3A.
3. Nelson / Marlborough in COC 7A and COC 7B.

Information on the general biology of cockles is included below rather than being repeated in the reports for each fishery.

2. BIOLOGY

The cockle, *Austrovenus stutchburyi*, formerly known as *Chione stutchburyi*, is a shallow-burrowing suspension feeder of the family Veneridae. It is found in soft mud to fine sand on protected beaches and enclosed shores around the North and South Islands, Stewart Island and the Chatham Islands (Morton & Miller 1968). Suspension feeders such as *A. stutchburyi* tend to be more abundant in sediments with a larger grain size. They are also common in eelgrass (e.g., *Zostera* sp.), which often co-occurs with sand flats.

Cockles are found from the lowest high water neap tide mark to the lowest part of the shore; there is some evidence that they may extend to 20 m depth in some areas. Larcombe (1971) suggested that the upper limit is found where submergence is only 3.5 hours per day. *A. stutchburyi* is often a dominant species and densities as high as 4500 / m² have been reported in some areas. In Pauatahanui Inlet the biomass has been estimated at, 80% (5000 t) of the total intertidal biomass in the Inlet. Cockles may filter 1.6 million cubic metres of water on each tidal cycle at Pauatahanui, with a profound effect on water quality.

Sexes are separate and the sex ratio is usually close to 1:1. Maturity appears to be primarily a function of size rather than age, with sexual maturity occurring at a size of about 18 mm shell length. Spawning extends over spring and summer, and fertilisation is followed by a planktonic larval stage lasting about 3 weeks. Significant depression of larval settlement has been recorded for areas of otherwise suitable substrate from which all live cockles have been removed. This suggests the presence of some conditioning factor.

Quite extensive movements of juveniles and smaller animals have been documented, but individuals > 25 mm shell length remain largely sessile, moving only in response to disturbance. Small cockles grow faster than large cockles, but overall, growth is fastest during spring and summer. Growth is slower in the higher tidal ranges and in high density beds. Significant increases in growth rates have been observed for individuals remaining in areas that have been 'thinned out' by simulated harvesting.

Given that cockles recruit to the spawning biomass at ~18 mm shell length, but do not recruit to commercial or non-commercial fisheries until closer to 30 mm shell length, there is some protection for the stock against egg overfishing, especially as the Snake Bank and Papanui and Waitati Inlet stocks are probably not isolated as far as recruitment of juveniles is concerned. However, this generality should be treated with some caution, given that some population of adults seems to be required to stimulate settlement of spat.

COCKLES (COC)

Natural mortality arises from a number of sources. Birds are a major predator of cockles (up to about 23 mm shell length). However, they appear to have little impact on cockle abundance. Other predators include crabs and whelks. Cockles are also killed after being smothered by sediments shifted during storms or strong tides.

Experimental work on Snake Bank led to estimates of absolute mortality of 17–30% per annum, instantaneous mortality (M) of 0.19–0.35, with a midpoint of $M = 0.28$. The estimated mortality rates for cockles of > 30 mm shell length were slightly greater at 19–37% per annum, (M of 0.21–0.46 with a midpoint of 0.33). This higher estimate was caused by relatively high mortality rates for cockles of > 35 mm shell length and, as these are now uncommon in the population, $M = 0.30$ (range 0.20–0.40) was assumed for yield calculations.

Studies in 1992 and 1993 at Snake Bank to estimate growth rate by sequential length frequency analysis resulted in an estimate of $K = 1.10$. Such a rate indicates fairly rapid growth (~ 2 years) to the size of interest to both commercial and amateur fisheries (~ 30 mm). Further work at Snake Bank in 1996 using length frequency distributions from 1991–96 confirmed the rapid growth rate with an estimate of $K = 1.02$. This rate is much faster than shown in previous Snake Bank studies (Martin 1984) where up to 4 or 5 years could be taken for cockles to attain a size of 30 mm. The increased growth rate may be a result of the lower densities of cockles on the bank in recent years.

Growth in Papanui and Waitati Inlets, Purakanui and Otago Harbour (Bed 1) was examined by Breen *et al.* (1999). They found that notched clams did not exhibit significant growth when recovered after one year, and modes in the length frequency distributions did not shift when measured over four sampling periods within a year. They concluded that it was unlikely that average growth is really as slow as the results indicated, but there may be high interannual variability in growth.

The Snake Bank work also showed moderate differences among years in the level of recruitment of juveniles to the population. The variability of recruitment was estimated as $\sigma_R = 0.41$ using all available data (1983–1996) but as $\sigma_R = 0.31$ using data only from those years since the fishery has been considered to be fully developed (1991–96). Given the variability of most shellfish populations and the shortness of the time series, this is probably an underestimate of the real variability of recruitment in the Snake Bank population.

3. FOR FURTHER INFORMATION

- Annala JH., Sullivan KJ. 1996. Report from the mid-year Fishery Assessment Plenary, November 1996: stock assessments and yield estimates. Ministry of Fisheries.
- Annala JH., Sullivan KJ., O'Brien CJ., Smith NWMcL., and Grayling SM. 2003. Report from the Fishery Assessment Plenary, May 2003: stock assessments and yield estimates. Ministry of Fisheries.
- Bradford E. 1998. Harvest estimates from the 1996 national recreational fishing surveys. New Zealand Fisheries Assessment Research Document. 1998/16: 27p.
- Breen P A., Carbines GC., and Kendrick TH. 1999. Stock assessment of cockles in Papanui and Waitati Inlets, Otago Harbour, and Purakanui, Otago. Final Report for the Ministry of Fisheries research project COC9701 dated July 1999.
- Cameron T. 1997. Aspects of growth and condition in *Austrovenus stutchburyi* (Finlay 19270 (Bivalvia: Veneridae) at Waitati Inlet: Influence of shore height and distance from mouth. Unpublished MSc thesis, University of Otago, Dunedin, New Zealand.
- Cranfield HJ., Micheal KP., and Francis RICC. 1996). Growth rates of five species of subtidal clam on a beach in the South Island, New Zealand. Marine and Freshwater Research 47: 773-784.
- Francis RICC. (1984). An adaptive strategy for stratified random trawl surveys. New Zealand Journal of Marine and Freshwater Research 18: 59-71.
- Hilborn R. and Walters CJ. 1992. Quantitative Fisheries Stock Assessment: Choice, dynamics and uncertainty. Chapman and Hall, London.
- Irwin CR. 1999. The effects of harvesting on the reproductive and population biology of the New Zealand Littleneck Clam (*Austrovenus stutchburyi*) in Waitati Inlet. Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- McKinnon J. 1996. Studies of the age, growth and shell increment patterns in the New Zealand cockle (*Austrovenus stutchburyi*). Unpublished Msc thesis, University of Otago, Dunedin, New Zealand.
- Manly BFJ., Akroyd JM., and Walshe KAR. 2002. Two-phase stratified random surveys on multiple populations at multiple locations. New Zealand Journal of Marine and Freshwater Research 36: 581-591.
- Morton J., Miller M. 1973. The New Zealand Sea Shore. Collins, Auckland. 653p.
- Stewart B., Keogh J., Fletcher D., and Mladenov P. 1992. Biomass survey of the New Zealand littleneck clam (*Chione stutchburyi*) in Papanui and Waitati Inlets, Otago during 1991/1992. Marine Science and Aquaculture research Centre, University of Otago, Dunedin, New Zealand. 37p.
- Stewart B. 2004. Warrington treated Sewage Discharge: Waitati Inlet Ecological Survey, 2004. Prepared for DCC. 21p.
- Stewart B. 2006. Stock assessment of Cockles (*Austrovenus stutchburyi*) in Papanui and Waitati Inlets, Otago, 2004. Final Research Report for the Ministry of Fisheries Research Project COC2004/02. 54p.
- Stewart B. 2008. Stock assessment of clams (*Austrovenus stutchburyi*) in Waitati Inlet, Otago, 2007. Final Research Report for Southern Clams Ltd. 24p.

- Sukhatme PV. 1954. Sampling theory of surveys: with applications. New Delhi : Indian Society of Agricultural Statistics. 49p.
- Teirney LD., Kilner AR., Millar RE., Bradford E., and Bell J.D. 1997. Estimation of recreational catch from 1991–92 to 1993–94 New Zealand Fisheries Assessment Research Document 1997/15: 43p.
- Wildish K. 1984a. The cockle resource in the Otago region: 1. An initial study of the New Zealand cockle (*Chione stutchburyi*) resource in Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries.
- Wildish K. 1984b. The cockle resource in the Otago region: further analysis of results from the survey of *Chione stutchburyi* (the New Zealand cockle) populations at Papanui and Waitati Inlets. Report to the Ministry of Agriculture and Fisheries.
- Wing S., Irwin C., Granger G. 2002. Biomass survey and yield estimates for the New Zealand littleneck clam *Austrovenus stutchburyi* in Papanui and Waitati Inlets, Otago. Final Report for the Ministry of Fisheries Research Project COC2001/02. 52p.
- Zar JH. 1996. Biostatistical Analysis. Third Edition. Prentice Hall International Inc.