


ELEPHANT FISH (ELE)*(Callorhinchus milii)***1. FISHERY SUMMARY****(a) Commercial fisheries**

Reported landings since 1936 are shown in Tables 1 and 2. From the 1950s to the 1980s, landings of elephantfish of around 1000 t were common. Most of these landings were from the area now encompassed by ELE 3 but fisheries for elephantfish also developed on the south and west coasts of the South Island in the late 1950s and early 1960s, with average catches of around 70 t per year in the south (in the 1960s to the early 1980s) and 10-30 t per year on the west coast. Total annual landings of elephantfish dropped considerably in the early 1980s (between 1982-83 and 1994-96 they ranged between 500 and 700 t) but later increased to the point that they have annually exceeded 1000 t since the 1995-96 fishing season.

Table 1: Reported total landings of elephantfish for calendar years 1936 to 1982. Sources: MAF and FSU data.

Year	Landings (t)	Year	Landings (t)	Year	Landings (t)	Year	Landings (t)	Year	Landings (t)	Year	Landings (t)
1936	116	1944	270	1952	459	1960	1163	1968	862	1976	705
1937	184	1945	217	1953	530	1961	983	1969	934	1977	704
1938	201	1946	235	1954	853	1962	1156	1970	1128	1978	596
1939	193	1947	188	1955	802	1963	1095	1971	1401	1979	719
1940	259	1948	230	1956	980	1964	1235	1972	1019	1980	906
1941	222	1949	310	1957	1069	1965	1111	1973	957	1981	690
1942	171	1950	550	1958	1238	1966	1112	1974	848	1982	661
1943	220	1951	602	1959	1148	1967	934	1975	602	-	-

The TACC for ELE 3 has, with the exception of 2002-03, been consistently exceeded since 1986-87. The ELE 3 TACC was consequently increased to 500 t for the 1995-96 fishing year, and then increased twice more under an Adaptive Management Programme (AMP): initially to 825 t in October 2000 and then to 950 t in October 2002. This new TACC combined with the allowances for customary and recreational fisheries (5 t each), increased the new TAC for the 2002-03 fishing year in ELE 3 to 960 t. In ELE 3 fishing is seasonal, mostly occurring in spring and summer in inshore waters. Most of the recent increase in catch from the ELE 3 fishery has been taken as a bycatch of the RCO 3 trawl fishery (Raj & Voller, 1999). During 1989-90 to 1997-98, the level of elephantfish bycatch from the RCO 3 fishery increased from around 50 t to 300 t (Raj & Voller, 1999). There was also a steady increase in the level of ELE 3 bycatch from the FLA 3 trawl fishery, with catches increasing from around 50 t in 1994-95 to 150 t in 1997-98. The fishery in ELE 5 is mainly a trawl

fishery targeted at flatfish and to a lesser extent giant stargazer. Very little catch in ELE 5 is taken by target setnet fisheries. The ELE 5 TACC was increased from 71 t to 100 t under an AMP in October 2001. The TACC was further increased under the AMP to 120 t in October 2004.

Table 2: Reported landings (t) of elephantfish by Fishstock from 1983–84 to 2005–06 and actual TACCs (t) from 1986–87 to 2005–06.

Fishstock FMA (s)	ELE 1		ELE 2		ELE 3		ELE 5		ELE 7		ELE 10		Total	
	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC	Landings	TACC
1983–84*	<1	–	5	–	605	–	94	–	60	–	0	–	765	–
1984–85*	<1	–	3	–	517	–	134	–	50	–	0	–	704	–
1985–86*	<1	–	4	–	574	–	57	–	46	–	0	–	681	–
1986–87†	<1	10	2	20	506	280	48	60	29	90	0	10	584	470
1987–88†	<1	10	3	20	499	280	64	60	44	90	0	10	610	470
1988–89†	<1	10	1	22	450	415	49	62	43	100	0	10	543	619
1989–90†	<1	10	3	22	422	418	32	62	55	101	0	10	510	623
1990–91†	<1	10	5	22	434	422	55	71	59	101	0	10	553	636
1991–92†	<1	10	11	22	450	422	58	71	78	101	0	10	597	636
1992–93†	<1	10	5	22	501	423	39	71	61	102	0	10	606	638
1993–94†	<1	10	6	22	475	424	46	71	41	102	0	10	568	639
1994–95†	<1	10	5	22	580	424	60	71	39	102	0	10	684	639
1995–96†	<1	10	7	22	688	500	72	71	93	102	0	10	862	715
1996–97†	<1	10	9	22	734	500	74	71	94	102	0	10	912	715
1997–98†	<1	10	12	22	910	500	95	71	66	102	0	10	1082	715
1998–99†	<1	10	9	22	842	500	129	71	117	102	0	10	1098	715
1999–00†	<1	10	6	22	950	500	105	71	87	102	0	10	1148	715
2000–01†	2	10	7	22	956	825	153	71	90	102	0	10	1207	1040
2001–02†	<1	10	9	10	852	825	105	100	88	102	0	10	1053	1057
2002–03†	1	10	9	10	950	950	106	100	59	102	0	10	1125	1182
2003–04†	<1	10	10	10	984	950	102	100	42	102	0	10	1139	1182
2004–05†	<1	10	13	10	972	950	125	120	74	102	0	10	1184	1202
2005–06†	<1	10	14	10	1023	950	147	120	76	102	0	10	1260	1202

* FSU data.

† QMS data.

(b) Recreational fisheries

Catches of elephantfish by recreational fishers are low compared to those of the commercial sector. Recreational fishing surveys carried out by the Ministry of Fisheries in the early 1990s estimated the recreational catch of elephantfish in the South region of ELE 3 in 1991–92 at 3000 fish, 1000 fish in the central region of ELE 7 in 1992–93, and no catch was reported in the North region in 1993–94 (Teirney et al., 1997). The national diary survey of recreational fishers in 1996 estimated that recreational catches of elephantfish were less than 500 fish in ELE 2, 1000 fish in ELE 3 and less than 500 fish in ELE 7 (Bradford, 1998). Estimates from the 1999/2000 recreational survey were 1000 fish in ELE 2, 2000 fish in ELE 3 and less than 500 in ELE 7 (Boyd & Reilly, 2002). Owing to biases inherent to telephone vs. face-to-face interviews, the 1999/2000 estimate is regarded to be the most accurate.

(c) Maori customary fisheries

Quantitative information on the current level of Maori customary take is not available.

(d) Illegal catch

There are reports of discards of juvenile elephantfish by trawlers from some areas. However, no quantitative estimates of discards are available.

(e) Other sources of mortality

The significance of other sources of mortality has not been documented.

2. BIOLOGY

Elephantfish are uncommon off the North Island, occurring south of East Cape on the east coast and south of Kaipara on the west coast. They are most plentiful around the east coast of the South Island.

Males mature at a length of 50 cm fork length (FL) at an age of 3 years, females at 70 cm FL at 4 to 5 years of age. The maximum age cannot be reliably estimated, but appears to be at least 9 years and may be as high as 15 years. The M value of 0.35 used is based on unvalidated ageing work indicating a maximum age of 13 years. This results from use of the equation $M = \log_e 100/\text{maximum age}$, where maximum age is the age to which 1% of the population survives in an unexploited stock.

Mature elephantfish migrate to shallow inshore waters in spring and aggregate for mating. Eggs are laid on sand or mud bottoms, often in very shallow areas. They are laid in pairs in large yellow-brown egg cases. The period of incubation is at least 5–8 months, and juveniles hatch at a length of about 10 cm FL. Females are known to spawn multiple times per season. After egg laying the adults appear to disperse and are difficult to catch; however, juveniles remain in shallow waters for up to 3 years. During this time juveniles are vulnerable to incidental trawl capture, but are of little commercial value.

Biological parameters relevant to the stock assessment are shown in Table 3. Provisional Von Bertalanffy growth curves based on MULTIFAN are available for Pegasus Bay and Canterbury Bight in 1966–68 and 1983–88. Because the growth curves were based on a MULTIFAN analysis of length-frequency data, the ages of the larger fish were probably underestimated and the growth curves are only reliable to about 4–5 years. Fish appeared to grow faster in the 1980s than in the 1960s.

Table 3: Estimates of biological parameters for elephant fish.

Fishstock	Estimate		Source	
1. Natural mortality (M)				
All	0.35		Francis (1997)	
2. Weight = a (length)^b (Weight in g, length in cm fork length)				
ELE 3 both sexes combined	a = 9.1 x 10 ⁻³ b = 3.02		Gorman (1963)	
3. Von Bertalanffy Growth Function				
	Pegasus Bay 1966–68		Canterbury Bight 1966–68	
	Males	Females	Males	Females
K (yr ⁻¹)	0.231 ± 0.002	0.096 ± 0.001	0.089 ± 0.002	0.060 ± 0.001
L _∞ (cm)	74.7 ± 0.12	156.9 ± 1.38	141.5 ± 2.28	203.6 ± 3.2
t ₀ (yr)	-0.78 ± 0.008	-0.87 ± 0.006	-0.96 ± 0.008	-1.06 ± 0.009
	Pegasus Bay 1983–84		Canterbury Bight 1988	
	Males	Females	Males	Females
K (yr ⁻¹)	0.473 ± 0.009	0.195 ± 0.008	0.466 ± 0.008	0.224 ± 0.001
L _∞ (cm)	66.9 ± 0.52	113.9 ± 2.89	62.7 ± 0.23	94.1 ± 0.26
t ₀ (yr)	-0.24 ± 0.017	-0.53 ± 0.023	-0.38 ± 0.015	-0.69 ± 0.006

3. STOCKS AND AREAS

There are no new data that alter the stock boundaries given in previous assessment documents.

There is only limited information available to support existing stock boundaries. Results from tagging studies conducted during 1966–69 indicate that elephantfish tagged in the Canterbury Bight remained in ELE 3. Separate spawning grounds to maintain each ‘stock’ have not been identified. The boundaries used are related to the historical fishing pattern when this was a target fishery.

4. STOCK ASSESSMENT

There are no new data which would alter the yield estimates given in the 1996 Plenary Report. The yield estimates are based on commercial landings data only and have not changed since the 1988 Plenary Report.

(a) Estimates of fishery parameters and abundance

(i) Trawl survey biomass indices

Indices of relative biomass are available from recent trawl surveys (Table 4). These have not been used to estimate absolute biomass or yields as historically, these trawl surveys have given variable abundance and high CV's for elephantfish, and probably have not monitored their biomass very well. A pilot survey off the east coast of the South Island was undertaken in the summer of 1996–97 and was repeated in 1997–98, 1998–99, 1999–00 and 2000–01. This survey was initiated for several reasons, including a need to better survey elephantfish in ELE 3 in view of the recent TACC increase. In February 1999, the Inshore Fishery Assessment Working Group concluded that it was not clear whether the ECSI trawl survey was adequately sampling elephantfish, as the commercial fishery for this species included depths <10 m and the *Kaharoa* is unable to trawl in such areas. Subsequently, in 1999–00 and 2000–01 the commercial vessel *Compass Rose* carried out surveys (concurrently) with the *Kaharoa* in which it fished areas inside 10 m. In 2001 the Inshore FAWG recommended that the east coast South Island trawl survey be discontinued due to the extreme variability in the catchability of the target species. A workshop to review the monitoring of inshore finfish (May 2006) concluded that the ECSI winter survey series should be reinstated, as based on simulations using existing data, it was predicted to provide useful relative biomass estimates for many species (excluding elephantfish). The workshop concluded that ELE 3 relative biomass should be estimated using industry run “hybrid” surveys.

Table 4: Relative biomass indices (t) and coefficients of variation (CV) for elephant fish for east coast South Island (ECSI) – summer and winter, west coast South Island (WCSI) and the Stewart-Snares Island survey areas assuming areal availability, vertical availability and vulnerability equal 1.0. Biomass is only estimated outside 10 m depth except for COM9901 and CMP0001. Note: because trawl survey biomass estimates are indices, comparisons between different seasons (e.g., summer and winter ECSI) are not strictly valid.

Region	Fishstock	Year	Trip number	Biomass estimate	CV (%)
ECSI(winter)	ELE 3	1991	KAH9105	300	40
		1992	KAH9205	176	32
		1993	KAH9306	481	33
		1994	KAH9406	152	33
		1996	KAH9606	858	30
ECSI(summer)	ELE 3	1996–97	KAH9618	1127	31
		1997–98	KAH9704	404	18
		1998–99	KAH9809	1718	28
		1999-00	KAH9917	1097	25
		1999–00	COM9901	802	73
				+475 (inside 10m)	79
		2000-01	KAH0014	693	18
2000-01	CMP0001	1229	29		
		+84 (inside 10m)	23		
WCSI	ELE 7	1992	KAH9204	38	42
		1994	KAH9404	167	33
		1995	KAH9504	85	35
		1997	KAH9701	94	33
		2000	KAH0004	42	63
Stewart-Snares	ELE 5	1993	TAN9301	219	33
		1994	TAN9402	177	47
		1995	TAN9502	69	49
		1996	TAN9604	137	46

(ii) CPUE biomass indices

ELE 3 is monitored using standardized CPUE, based on non-zero catches recorded by bottom trawl fishery targeting RCO, as an index of relative abundance. The CPUE trend was updated to 2003/04 as part of the ELE 3 AMP in 2005 (SeaFIC, 2005).




Figure 1: Relative CPUE indices for ELE 3 using the loglinear non-zero model based on the bycatch in the RCO 3 target trawl fishery. Error bars are $\pm 2*SE$. The equivalent series from the 2002 RCO 3 target fishery is also shown as a thick grey line.

(b) Biomass Estimates

Estimates of current and reference absolute biomass are not available.

(c) Estimation of Maximum Constant Yield (MCY)

MCY was estimated from the equation $MCY = cY_{av}$ (Method 4). The value c was set equal to 0.7 based on the estimate of $M = 0.35$. Mean catches for the years 1983–84 to 1985–86 were used to estimate MCY because the fishery appeared to stabilise after an earlier period of decline.

(i) South–East (Coast) and South–East (Chatham Rise) (ELE 3)

$$MCY = 0.7 * 565.5 \text{ t} = 396 \text{ t (rounded to 400 t).}$$

(ii) Southland and Sub–Antarctic (ELE 5)

$$MCY = 0.7 * 94.9 \text{ t} = 66 \text{ t (rounded to 70 t).}$$

(iii) Challenger/Central (Plateau) (ELE 7)

$$MCY = 0.7 * 52.3 \text{ t} = 37 \text{ t (rounded to 40 t).}$$

The estimate of M is uncertain and recruitment variability may be low, so the estimate of c is uncertain. The MCY estimates are considered approximate and are probably conservative.

The level of risk to the stock by harvesting the population at the estimated MCY value cannot be determined.

(d) Estimation of Current Annual Yield (CAY)

CAY cannot be determined.

Yield estimates are summarised in Table 5.

Table 5: Yield estimates (t) for elephant fish.

Parameter	Fishstock	Estimate
MCY	ELE 3	400
	ELE 5	70
	ELE 7	40
	Total	510
CAY	All	Cannot be determined

(e) Other yield estimates and stock assessment results

No other yield estimates are available.

(f) Other Factors

The amount of quota allocated was below historic catch levels and has reduced elephantfish mainly to a trawl bycatch for inshore vessels. On the east coast of the South Island the availability of elephantfish since the start of the QMS appears to have been high, and many individual fishers have exceeded their quotas. As a result, deeming and bycatch trading of this species has increased.

Target fishing for elephantfish using set nets has decreased since the introduction of the QMS. The distribution of the target trawl species such as red cod, barracouta and flatfish influences the likelihood of fishers encountering elephantfish.

5. ANALYSIS OF ADAPTIVE MANAGEMENT PROGRAMMES (AMP)

The Ministry of Fisheries revised the AMP framework in December 2000. The AMP framework is intended to apply to all proposals for a TAC or TACC increase, with the exception of fisheries for which there is a robust stock assessment. In March 2002, the first meeting of the new Adaptive Management Programme Working Group was held. Two changes to the AMP were adopted:

- a new checklist was implemented with more attention being made to the environmental impacts of any new proposal;
- the annual review process was replaced with an annual review of the monitoring requirements only. Full analysis of information is required a minimum of twice during the 5 year AMP.

ELE 3

The ELE 3 TACC has been increased twice under AMP management: from 500 to 825 t in October 2000 and from 825 to 950 t in October 2002.

Full-term Review of ELE 3 AMP in 2007

In 2007 the AMP FAWG reviewed the performance of the AMP (Starr et al., 2007). The Working Group noted:

Fishery Characterization

- The ELE 3 TACC was increased from 500 t to 825 t in October 2000. An additional allowance of 5 t was made for customary and recreational use, bringing the total TAC to 830 t per year. The

TACC was again increased to 950 t in Oct 2002 in response to ongoing difficulties with limiting catches within the TACC.

- After an initial period of low catches from 1936 - 1950, ELE 3 catches increased rapidly, and catches over 1000 t per year (probably under-estimated) were common in the 1950s and 1960s. Catches then declined steadily to ~500 t by the time ELE 3 was introduced into the QMS in 1986/87.
- Since then, catches have increased rapidly, exceeding the TACC throughout the past two decades, and reaching similar levels to the 1950s by 1997/98. As a result of increasing TACCs under the AMP, the level of overcatch has decreased, and has averaged 3% over the past four fishing years.
- Management efforts to control this overcatch have likely affected the accuracy of reported and landed catch. In particular, anecdotal reports indicate that significant dumping of elephantfish probably occurred as a consequence of high deemed values for this fishery after 2000/01 and the elimination of the bycatch trade-off scheme in 1997/98. Deemed values were subsequently decreased from October 2005 onwards to encourage accurate reporting.
- 83% of the ELE 3 catch is taken by multi-species bottom trawls, and the remaining 17% by setnets. The trawl catch is primarily taken in the Canterbury Bight, with smaller catches in Pegasus Bay area. The setnet catch is mostly taken south of the Banks Peninsula in the Bight.
- The elephantfish fishery is quite seasonal, with trawl and setnet catches mainly being taken from October - March, mainly as a result of a summer fishery in area 22. Trawl catches have tended to extend further into the fishing year since 1997/98, particularly in other areas.
- ELE are caught in a range of target fisheries. The trawl catch is primarily made while targeting red cod, barracouta, elephantfish and flatfish, with a significant increase in targeting for elephantfish from 01/02 onwards. Setnet catches are made in the multispecies shark fishery targeting rig, school shark and elephantfish. Recreational catches are poorly estimated, but probably <5 t.

CPUE Analysis

- The 2001 Working Group concluded that the most suitable CPUE index for ELE 3 was provided by non-zero catch records in the red cod targeted trawl fishery. Additional analyses conducted since 2005 include two other fisheries: the flatfish targeted trawl fishery and the shark-targeted setnet fishery.
- CPUE for these fishery definitions were standardised using a lognormal model based on non-zero catches. In addition, a binomial model was used to investigate the effect of changing proportion of non-zero catches.
- The three standardised CPUE series provide fairly similar and consistent trends, all showing a steady increase in CPUE from 89/90 to somewhere around 99/00 - 01/02, followed by a decline in catch rates to the present time, possibly with some levelling off over the past two or three years.
- Offsets in the maximum catch rate peaks in these three fisheries may be related to the flatfish fishery taking smaller fish in shallower water, and the selectivity of the setnet fishery for larger (older) fish.
- Standardisation has little effect on the nominal CPUE from the flatfish fishery. However, it does have a strong effect on catch rates in the red cod fishery since 2002/03, converting the steep increase in unstandardised CPUE into a level trend, suggesting increasing effects of targeting on elephantfish and the effect of a trend towards longer tows in recent years.
- Standardisation of the setnet fishery converts the strongly fluctuating unstandardised data into a smoother trend quite similar to the other two series. The three series in combination provide a consistent picture of catch rates in this fishery.
- The Working Group noted fleet behaviour changes at the time of management changes in 00/01 - 01/02, with increases in factors such as trip length, tow duration and number of tows, which contribute to the increase in non-zero catches. The Working Group noted that these behaviour changes all confirm an increase in targeting on ELE after 2000.
- Vessels are therefore acting as proxies for effort, which does not enter this model. The Working Group again proposed that the binomial component of the models be accorded little weight.

- The progression in CPUE peaks across the fisheries is consistent with the biology of the species and what is known about mean sizes caught in these fisheries. In general, the Working Group concluded that they had more confidence in the CPUE trends prior to 2000, but less confidence in trends thereafter.

Trawl Surveys

- A series of winter trawl surveys were as conducted off the ECSI from 1991 - 1996, and a series of summer surveys from 1996/97 - 2000/01, for a range of species, including elephantfish.
- Results of these surveys were highly variable, suggesting unrealistically high inter-annual changes in abundance. These changes appear to have resulted more from annual changes in catchabilities of elephantfish and other species.
- Length-frequency distributions of fish in these surveys also showed high inter-annual variability, which is unlikely to reflect actual changes in the population structure.
- Estimates of abundance from these surveys widely straddle the CPUE abundance indices, alternately greatly exceeding or being well below CPUE estimates in alternate years.
- The 2001 FAWG meeting concluded that the summer ECSI survey was not reliably monitoring elephantfish, showing changing catchabilities that exceeded likely changes in abundance. The Working Group recommended that the summer trawl survey should be discontinued.

Logbook Programme

- A logbook programme was introduced into QMA3 in 1994/95 to cover the three main target shark species (SPO, SCH and ELE) in the setnet fishery. A bottom trawl logbook programme was initiated by the SEFMC in mid 2002, primarily as part of the AMP proposal for an increase TACC for ELE 3.
- The setnet logbook programme has achieved good levels of coverage, covering >40% of catch in 95/96 and from 98/99 - 01/02, with a peak of 63.5% coverage in 00/01. However, coverage decreased substantially to 11.6% in 04/05, and has since increased again to 21.4%. Apart from 97/98 (when coverage levels were minimal for some reason) the number of sharks sampled has generally exceeded 600 per year, with a peak of 1671 sharks sampled in 00/01.
- Setnet logbooks have provided good coverage by area and season, with particularly good coverage of the main Canterbury Bight fishing area. The programme has never achieved good coverage of the Kaikoura area, but ELE catches have always been small in that area.
- Bottom trawl logbook coverage has been poor (1.6% - 4.4%) in all years except 03/04, when a concerted effort achieved 19.6% coverage. Despite this poor coverage, 2000 - 2500 sharks have been sampled from 02/03 onwards, with 7239 being sampled in 03/04.
- Trawl logbook coverage has also focussed in the Canterbury Bight, and has never attained good seasonal coverage.
- These logbooks have provided good length-frequency samples, indicating similar length-frequency distributions in all years from 95/96 to 04/05, with consistently lower mean size for males (45 cm – 70 cm) than females (55 cm – 85 cm). The sex ratio is also skewed ~70% to females.
- There are no indications of any decline in mean size or decrease in proportions of larger sharks in the setnet fishery. The Working Group questioned the substantial increase in the apparent size of both males and females in 2005/06 samples. Did this result from a change in sampling procedures, misinterpretation of instructions, or a shift in the fishery to different depths or areas?
- In contrast, sex ratios in the trawl fishery averaged ~50:50, and length-frequencies show higher inter-annual variability, but with no clear apparent trends.

Effects of Fishing

- A general overview of the effects of setnet and trawl fishing along the South Island East Coast is presented under AMP reviews for SPO 3, SCH 3 and GUR 3.
- No additional information was presented on specific effects of the ELE 3 fishery.

Conclusions

- Continuing catch increases since 1989/90, sustained high catches, increasing CPUE after the low point of 1989/90 and absence of evidence of declines in mean size all indicate that the fishery is not having a substantial impact on the abundance of the ELE3 stock.
- However, there are indications of increased targeting on elephantfish, since 2000/01. There are also concerns that other fleet behavioural changes in response to management actions may have reduced the extent to which CPUE reflects abundance since 2000/01. There is lower confidence in recent CPUE estimates.
- Catches at the current TACC level are likely to be sustainable in the short to medium-term. However, it is not known whether current catches are sustainable in the long-term.

ELE 5

The ELE 5 TACC for was increased from 71 t to 100 t under an AMP in October 2001. The TACC was further increased under AMP Management to 120 t in October 2004.

Medium-term Review of ELE 5 AMP in 2007

The AMP FAWG reviewed Medium-term review of the ELE 5 AMP (Starr et al., 2007b). The Working Group noted:

Fishery Characterization

- The ELE 5 TACC was increased from 71 t to 100 t in October 2001. Allowances of an additional 5 t each for recreational and customary use brought the total TAC to 110 t. The TACC was again increased to 120 t in October 2004 as a result of ongoing difficulties with limiting catches within the TACC. An additional 16 t allowance for recreational and customary fishing brought the TAC to 136 t.
- Catches remained below the TACC until 1995/96, and then escalated rapidly to almost double the TACC level by 00/01. Following increase in the TACC under the AMP in 2001, catches dropped down and remained at about the TACC level from 02/03 - 04/05, attaining the new AMP TACC level in 04/05. Catches in 2005/06 have again exceeded the TACC by 27 t, but are still below the highest reported level of 154 t in 00/01.
- The ELE 5 fishery would have had similar problems to the ELE 3 fishery, where management efforts to control overcatch have likely affected the accuracy of reported and landed catch. In particular, anecdotal reports indicate that significant dumping of elephantfish probably occurred as a consequence of high deemed values after 2000/01.
- 87% of the ELE 5 catch is taken in the bottom trawl fishery, mainly as a bycatch in the flatfish or stargazer targeted fishery, a further 12% is caught in the multispecies shark-targeted (SPO, SCH, ELE) setnet fishery.
- Almost all the trawl and setnet ELE 5 landings come from the Western Foveaux Strait (area 30) and Eastern Stewart Island (area 25). The flatfish targeted trawl fishery occurs across both areas, whereas stargazer, and to some extent elephantfish targeting, occurs mainly in the western Strait. In the setnet fishery, more rig targeting occurs east of Stewart Island, with school shark and elephantfish targeting being more prevalent in the Western Strait.
- The setnet fishery is strongly seasonal, occurring mainly from November – February. The trawl fishery also used to be a predominantly summer fishery, but switched to a year-round fishery in 98/99.
- ELE 5 trawl catches are mostly made in a fairly narrow, shallow depth range from ~40 m – 100 m.

CPUE Analysis

- Previous standardised CPUE analyses have been conducted using data from the flatfish targeted trawl fishery. However, noting loss of data from key fishing months in earlier years, the 2006 Plenary report concluded that this was not a reliable index of abundance for ELE 5.
- With 4 years additional data, three different fishery definitions were used in standardised CPUE analyses this year: the flatfish targeted trawl fishery in area 30, the flatfish trawl fishery in area 25 and the multispecies shark-targeted (SPO, SCH, ELE) setnet fishery.
- CPUE for these fishery definitions were standardised using a lognormal model based on non-zero catches. In addition, a binomial model was used to investigate the effect of changing proportion of non-zero catches.
- The two bottom trawl standardised series showed different trends. Area 30 showed a strong peak in 94/95, followed by a decline to 98/99 and a stable trend thereafter at about historic CPUE levels. Area 25 shows a generally increasing trend to a peak in 03/04, with a decline thereafter. The setnet index is essentially stable at the average level across the period, but with a peak in 97/98.
- The overall impression of the three indices overlaid is of quite strongly fluctuating CPUE around a fairly stable long-term average level. It appears that these standardised series may be reflecting inter-annual changes in availability or targeting in these fishery components, rather than actual abundance.
- The Working Group noted a very strong seasonal signal in the trawl indices, with summer catches being 6 times the winter catches. This raises the question as to whether the summer fishery dominated index is an index of abundance, or just an index of strongly targeted effort on a dense near-shore summer aggregation. In contrast, the setnet fishery does not show a strong seasonal signal, possibly because catch levels for elephantfish are low and that this species is only taken as bycatch.
- Trends in the two trawl areas appear to contradict each other, with differences between the two areas difficult to understand. The Working Group raised questions regarding whether apparent changes in targeting reflected actual targeting changes, or simply changes in reporting of target, in which case stargazer would also be part of the fishery. The Working Group therefore requested that the analysis be repeated using all vessels which caught either ELE or STA in the flatfish fisheries in areas 25 and 30.
- The re-analysis of a mixed species target trawl CPUE was presented to the AMP FAWG at a subsequent meeting, based on data from a suite of common core vessels which fished in both Area 025 and 030. The analysis concentrated on targeted flatfish, stargazer and elephantfish, with each area analysed independently. Standardised CPUE indices in these two fisheries then showed similar trends, with the trend in stat area 30 being more optimistic. A scatter plot of the vessel coefficients also showed a reasonable level of consistency in these coefficients across the two analyses.

Trawl Surveys

- Four trawl surveys were conducted on the plateau extending southwest off the South Island from 1993 - 1996, targeted at the 6 main commercial species: barracouta, blue warehou, gemfish, ling, school shark and stargazer.
- The depth range of the surveys was generally deeper than optimum ranges for elephantfish, and the surveys are not expected to have provided reliable estimates of ELE abundance in the area. The estimates of ELE abundance from these surveys show high variance and no trend.

Logbook Programme

- A setnet logbook programme was introduced into FMA3 in 1994/95 for SPO, SCH and ELE, but primarily in support of the SPO 3 AMP. This programme was extended to area 5 in 1995/96. Most of the logbook sets in the Southland were targeting school shark.
- A trawl logbook program was implemented in 2002 in support of the ELE 3 AMP, and extended to FMA5 when the ELE 5 TACC was increased in October 2004.

- However, coverage has been poor, with sampling of ELE 5 only being reported in 95/96 and 98/99. Coverage of the setnet catch in these years was only 5.2% and 2.8% respectively, with 46 and 20 elephantfish being sampled.
- Despite the low coverage, setnet logbooks obtained reasonable representation of the main fishing areas in those 2 years, although good seasonal coverage was only achieved in 98/99.
- Coverage by the trawl logbooks has also been inadequate, increasing from 4.4% in 02/03 to 12.9% in 04/05, but decreasing again to only 2.3% in 05/06. The number of elephantfish sampled has been better, increasing from 411 to 706 fish per year by 04/05, but decreasing to 60 fish last year.
- Sex ratios in the setnet logbook samples showed about 80% females, and larger mean size for females. Size compositions were similar between the two years sampled, although sample numbers are too low to place any reliance on these.
- Trawl logbooks have achieved reasonable representation of areas and across the seasons in only one of the four years sampled.
- In marked contrast to the setnet fishery, trawl samples contained, on average, about 54% males. Size composition of males and females shows inter-annual variations, with a shift to larger fish in 2003/04. However, given the poor seasonal coverage, these differences are likely to result from seasonal sampling differences.
- Analysis of males by size in the trawl samples provided a good maturity ogive, indicating size-at-50% maturity at ~50 cm – 55 cm.

Effects of Fishing

- A general overview of the effects of setnet and trawl fishing along the South Island south coast is presented under the AMP reviews for SCH 5, which is essentially the same fishery as the ELE 5 fishery. No additional information was presented on specific effects of the ELE 5 fishery.
- The Working Group noted that logbook information indicated that some trawling and/or setnet activity occurred in the Foveaux Strait dredge oyster fishery area. The specific extent of overlap, and the potential effect of the SCH 5 fisheries on the oyster fishing area, should be evaluated and reported.
- The Working Group also noted that white sharks had recently been accorded protected status. Any available information on interactions with white sharks by e.g. the setnet fishery should be reported, and whether the industry has implemented any measures to reduce and report such interactions.

Conclusions

- The 2006 Plenary report notes that, at the time of the increases in TACC under the AMP, there was a “*reasonable probability that the stock was above the size that would support the MSY. However, it is not known if current catch levels and current TACCs are sustainable*”. The results of this year’s CPUE analysis do not provide any basis for changing those conclusions.

6. STATUS OF THE STOCKS

No estimates of current and reference biomass are available.

ELE 2

It is not known if recent catch levels or the current TACC are sustainable, or if they are at levels that will allow the stock to move towards a size that will support the maximum sustainable yield.

ELE 3

ELE 3 is currently being managed within an AMP that is monitoring a CPUE index derived from elephantfish landings in the red cod bottom trawl fishery. Before the introduction of the QMS, elephantfish off the south east coast of the South Island were considered severely overfished, and TACs were initially set at low levels to facilitate stock recovery. Based on the observed catches, and on the increasing biomass trend from the CPUE analysis, it now appears that this stock has rebuilt considerably since the mid-1980s. The TACC for ELE 3 was increased to 950 t for the 2002–03 fishing year under the AMP. Recent catch levels substantially exceeded the TACC and the MCY; but the MCY estimate is probably conservative. It cannot be determined if the current TACC of 950 t is sustainable in the long-term or if it will allow the stock to move towards the size that will support the maximum sustainable yield.

ELE 5

Before the introduction of the QMS, elephantfish in Southland were considered severely overfished, and TACs were initially set at low levels to facilitate stock recovery. The TACC for ELE 5 was increased twice within the AMP - first to 100 t in October 2001 and then to 120 t in October 2004 - as there was a reasonable probability that the stock was above the size that would support the MSY. However, it is not known if recent catch levels and current TACCs are sustainable or if they are at levels that will allow the stocks to move towards a size that will support the MSY.

ELE 7

In ELE 7 catches since 1987–88 have been above the MCY and below the TAC. It is not known if recent catch levels and current TACCs are sustainable or if they are at levels that will allow the stocks to move towards a size that will support the MSY.

TACCs and reported landings are summarised in Table 6.

Table 6: Summary of yields (t), TACCs (t), and reported landings (t) for elephant fish for the most recent fishing year.

Fishstock	QMA		MCY	2005–06 Actual TACC	2005–06 Reported landings
ELE 1	Auckland (East) (West)	1 & 9	–	10	<1
ELE 2	Central (East) (West)	2 & 8	–	10	14
ELE 3	South-East (Coast) (Chatham)	3 & 4	400	950	1023
ELE 5	Southland and Sub-Antarctic	5 & 6	70	120	147
ELE 7	Challenger	7	40	102	76
ELE 10	Kermadec	10	–	10	0
Total			510	1202	1260

7. FOR FURTHER INFORMATION

- Boyd, R.O., Reilly, J.L. (2002). 1999/2000 national marine recreational fishing survey: harvest estimates. *Draft New Zealand Fisheries Assessment Report*.
- Bradford, E. (1998). Harvest estimates from the 1996 national recreational fishing surveys. *N.Z. Fisheries Assessment Research Document 98/16* 27p.
- Coakley, A. (1971). The biological and commercial aspects of the elephantfish. *Fisheries Technical Report No. 76*. 29 p.
- Francis, M.P. (1996). Productivity of elephantfish – has it increased? *Seafood NZ Feb 96*: 22–25.
- Francis, M.P. (1997). Spatial and temporal variation in the growth rate of elephantfish (*Callorhynchus milii*). *N.Z. Journal of Marine and Freshwater Research 31*(9–23).
- Gorman, T.B.S. (1963). Biological and economic aspects of the elephantfish, *Callorhynchus milii* Bory, in Pegasus Bay and the Canterbury Bight. *Fisheries Technical Report No. 8*. 54 p.
- Langley, A.D. (2001). The analysis of ELE 3 catch and effort data from the RCO 3 target trawl fishery, 1989–90 to 1999–2000. *New Zealand Fisheries Assessment Report 2001/66*. 33 p.
- Lydon, G.J.; Middleton, D.A.J.; Starr, P.J. (2006). Performance of the ELE 3 and ELE 5 Logbook Programmes. AMP-WG-06/18. (Unpublished manuscript available from the NZ Seafood Industry Council, Wellington.)
- McClatchie, S.; Lester, P. (1994). Stock assessment of the elephantfish (*Callorhynchus milii*). *N.Z. Fisheries Assessment Research Document 94/6*. 17 p.
- Raj, L.; Voller, R. (1999). Characterisation of the south-east elephantfish fishery–1998. 55 p. (Report held by Ministry of Fisheries, Dunedin, New Zealand.)
- SeaFIC (2000). Proposal to the Inshore Fishery Assessment Working Group. Placement of the ELE 3 into Adaptive Management Programme dated 23 March 2000 (presented to the Inshore Fishery Assessment Working Group 28 March 2000). Copies held by MFish.
- SeaFIC (2002). Report to the Inshore Fishery Assessment Working Group: Performance of the ELE 3 Adaptive Management Programme (dated 25 February 2002). Copies held by MFish.
- SeaFIC (2003a). 2003 performance report: ELE 3 Adaptive Management Programme. AMP-WG-2003/06 3p. Copies held by MFish.
- SeaFIC (2003b). Report to the Adaptive Management Fishery Assessment Working Group: Performance of the ELE 5 Adaptive Management Programme and request for an additional increase in ELE 5. AMP-WG-2003/07 39 p. Copies held by MFish.
- SeaFIC (2005a). 2005 Report to the Adaptive Management Programme Fishery Assessment Working Group: Performance of the ELE 3 Adaptive Management Programme. AMP-WG-2005/16. Copies held by MFish.
- SeaFIC (2005b). 2005 Report to the Adaptive Management Programme Fishery Assessment Working Group: Performance of the ELE 5 Logbook Programme. AMP-WG-05/23. Copies held by MFish.
- SEFMC (2002a). (Southeast Finfish Management Company) 2002 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 3 Adaptive Management Programme (dated 25 February 2002). Copies held by MFish.
- SEFMC (2002b). (Southeast Finfish Management Company) 2002 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 5 Adaptive Management Programme (dated 25 February 2002). Copies held by MFish.
- SEFMC (2003). (Southeast Finfish Management Company) 2003 Report to the Inshore Fishery Assessment Working Group. Performance of the ELE 5 Adaptive Management Programme and request for an increase in ELE 5 (dated 13 Nov 2003). Copies held by MFish.
- Starr, P. J. (in prep.). Stock assessment of east coast South Island elephantfish (ELE 3). *Draft NZ Fisheries Assessment Report*. 32 p.
- Starr, P.J Kendrick, T.H.; Lydon, G.J.; and Bentley, N. (2007a). Report to the Adaptive Management Programme Fishery Assessment Working Group: Full-term review of the ELE 3 Adaptive Management Programme. AMP-WG-07/07. (Unpublished manuscript available from the Ministry of Fisheries, Wellington.). 104p.
- Starr, P.J Kendrick, T.H.; Lydon, G.J.; and Bentley, N. (2007b). Report to the Adaptive Management Programme Fishery Assessment Working Group: Two-year review of the ELE 5 Adaptive Management Programme. AMP-WG-07/10. (Unpublished manuscript available from the Ministry of Fisheries, Wellington.). 89p.
- Sullivan, K.J. (1977). Age and growth of the elephantfish *Callorhynchus milii* (Elasmobranchii: Callorhynchidae). *N.Z. Journal of Marine and Freshwater Research 11*: 745–753.
- Teirney, L.D.; Kilner, A.R.; Millar, R.E.; Bradford, E.; Bell, J.D. (1997). Estimation of recreational catch from 1991/92 to 1993/94. *N.Z. Fisheries Assessment Research Document 97/15*. 43 p.