


PACIFIC BLUEFIN TUNA (TOR)
(*Thunnus orientalis*)



1. FISHERY SUMMARY

Pacific bluefin tuna was introduced into the QMS on 1 October 2004 under a single QMA, TOR 1, with allowances, TACC, and TAC as follows:

<u>Fishstock</u>	<u>Recreational Allowance</u>	<u>Maori customary Allowance</u>	<u>Other mortality</u>	<u>TACC</u>	<u>TAC</u>
TOR 1	1	0.54	2.5	116	120

Pacific bluefin tuna were added to the Third Schedule of the 1996 Fisheries Act with a TAC set under s14 because Pacific bluefin tuna is a highly migratory species and it is not possible to estimate MSY for the part of the stock that is found within New Zealand fisheries waters.

Pacific bluefin tuna is believed to be a single Pacific-wide stock and is covered by two regional fisheries management organisations, the Western and Central Pacific Fisheries Commission (WCPFC), and the Inter-American Tropical Tuna Commission (IATTC). They will cooperate in the management of the Pacific bluefin tuna stock throughout the Pacific Ocean. Under the WCPFC Convention, New Zealand will be responsible for ensuring that the management measures applied within New Zealand fisheries waters are compatible with those of the Commission.

(a) Commercial fisheries

Pacific bluefin tuna was not widely recognised as a distinct species until the late 1990s. It was previously regarded as a sub-species of *Thunnus thynnus* (northern bluefin tuna, NTU). Prior to June 2001, catches of this species were either recorded as NTU or misidentified as southern bluefin tuna. Fishers have since become increasingly able to accurately identify TOR and, from June 2001, catch reports have rapidly increased. Catches of TOR may still be under reported to some degree as there is still some reporting against the NTU code. Recent genetic work suggests that true NTU (*Thunnus thynnus*) are not taken in the New Zealand fishery (see Biology section below for further details).

Pacific bluefin has been fished in the New Zealand EEZ since at least 1960, with some catch likely but undocumented prior to that time. New Zealand catches while increasing, are small compared to total stock removals.

Table 1: Reported total New Zealand landings (t) of Pacific bluefin tuna (includes landings attributed to NTU), 1991 – 2000 and total Pacific Ocean catches.

Year	NZ landings (t)	Total stock (t)	Year	NZ landings (t)	Total stock (t)
1991	1.5	13 876	1999	21.2	25 617
1992	0.3	12 962	2000	20.9	28 859
1993	5.6	9 617	2001	49.8	17 385
1994	1.9	14 913	2002	55.4	17 459
1995	1.8	25 842	2003	40.8	15 920
1996	4.2	21 978	2004	67.3	21 707
1997	14.3	22 086	2005	20.1	17 881
1998	20.4	13 918			

Source: NZ landings, for 1991-2002 Ministry of Fisheries Licensed Fish Receiver Reports and Solander Fisheries Ltd. 2003-2005 Ministry of Fisheries MHR data. Total Pacific landings for ISC members from ISC/06/PLENARY/14 prepared for ISC-6 March 2006. This covers most catches from this stock, but does not include South Pacific catches by coastal states in the South Pacific.

Catches from within New Zealand fisheries waters are very small compared to those from the greater stock in the Pacific Ocean (0.2% average for 1999-2002 of the Pacific wide catch). In contrast to New Zealand, where Pacific bluefin tuna are taken almost exclusively by longline, the majority of catches are taken in purse seine fisheries in the WCPO (Japan and Korea) and EPO (Mexico). Much of the fish taken by the Mexican fleet are on grown in sea pens.

Prior to the introduction to the QMS, the highest catches have been made in FMA 1 and FMA 2. While it is possible to catch Pacific bluefin as far south as 48° S, few catches are made in the colder southern FMAs. Although recent catches have occurred in FMA 7 fish have been in poor condition with little commercial value. Catches are almost exclusively by tuna longlines, typically as a bycatch of sets targeting bigeye tuna. Catches by fishing year and fleet are provided in Table 2.

Table 2: Reported catches or landings (t) of Pacific bluefin tuna by fleet and Fishing Year. NZ: New Zealand domestic and charter fleet, ET: catches from New Zealand flagged longline vessels outside these areas, JPNFL: Japanese foreign licensed vessels, KORFL: foreign licensed vessels from the Republic of Korea, and LFRR: Estimated landings from Licensed Fish Receiver Returns.

Fish Yr	TOR 1 (all FMAs)			LFRR/MHR	NZ ET
	JPNFL	NZ	Total		
1979/80	1.5		1.5		
1980/81	5.3		5.3		
1981/82	110.1		110.1		
1982/83	70.1		70.1		
1983/84	47		47		
1984/85	6		6		
1985/86	5.7		5.7		
1986/87	10.6		10.6	0.0	
1987/88	13.5		13.5	0.0	
1988/89	15.1		15.1	0.0	
1989/90	14.7		14.7	0.0	
1990/91	14.5		14.5	1.5	
1991/92	9.1		9.1	0.3	
1992/93	2.1		2.1	5.6	
1993/94	0.1		0.1	1.9	
1994/95			0	1.8	
1995/96			0	4.0	
1996/97		12.5	12.5	13.0	
1997/98		22.5	22.5	20.9	0.4
1998/99		20.6	20.6	17.9	0.1
1999/00		32.6	32.6	23.1	0.1
2000/01		43.9	43.9	51.8	1.0
2001/02		54.4	54.4	53.3	0.0
2002/03		41.6	41.6	39.8	0.0
2003/04		64.3	64.3	58.1	0.0
2004/05		21.6	21.6	21.1	0.0

(b) Recreational fisheries

Recreational fishers make occasional catches of Pacific bluefin tuna. In 2004 and 2005 a target recreational fishery developed off the west coast of the South Island during the hoki spawning ground fishery (August – September). Fish taken in this fishery have been submitted for various world

records for this species. Based on reports from this fishery described in various fishing magazines, catches have likely exceeded the allowance of 1t for this fishery. Due to the small numbers of fish landed, their large individual size, and reporting in the recreational fishing media it is feasible to derive a minimum estimate of recreational harvest of Pacific bluefin tuna. There is no information on the size of catch from the National Surveys of recreational fishers.

(c) **Maori customary fisheries**

There is no quantitative information available to allow the estimation of the harvest of Pacific bluefin tuna by customary fishers; however the Maori customary catch of Pacific bluefin is probably negligible because of the species seasonal and offshore distribution.

(d) **Illegal catch**

There is no known illegal catch of Pacific bluefin tuna in New Zealand fisheries waters.

(e) **Other sources of mortality**

There is likely to be a low level of shark damage and discard mortality of Pacific bluefin caught on tuna longlines that may be on the order of 1-2% assuming all tuna species are subject to equivalent levels of incidental mortality. There have been reports that some fish hooked in the target recreational fishery have been lost due to entanglement of the fishing line with trawl warps. The survival of these lost fish is not known. An allowance of 2.5 t has been made for other sources of mortality.

2. BIOLOGY

Pacific bluefin are epi-pelagic opportunistic predators of fish, crustaceans and cephalopods found within the upper few hundred meters of the water column. Individuals found in New Zealand fisheries waters are mostly adults. Adult Pacific bluefin occur broadly across the Pacific Ocean, especially the waters of the North Pacific Ocean.

There has been some uncertainty among fishers regarding bluefin tuna taken in New Zealand waters. Some fishers believe that three species of bluefin tuna are taken in New Zealand waters with some small catches of true “Northern” Atlantic tuna (*Thunnus thynnus*) in addition to Pacific and southern bluefin tuna. This belief is based on several factors include differences in morphology and the prices obtained for certain fish on the Japanese market.

To address this issue muscle tissue samples were taken from 20 fish for which there was uncertainty as to whether the fish was a Pacific bluefin tuna (*Thunnus orientalis*) or an Atlantic bluefin tuna. A further sample from a fish thought to be a southern bluefin tuna was also included. The tissue samples were sequenced for the COI region of DNA, and the sequences compared with COI sequences for the three species of tuna held in GenBank. All of the DNA sequences, except one, matched with sequences for Pacific bluefin tuna. The final sample was confirmed as a southern bluefin tuna. Therefore, based on DNA analysis, there is presently no evidence that Atlantic bluefin tuna are taken in New Zealand waters. Further tissue samples from fish thought by fishers to be NTU will be collected by scientific observers.

Adult Pacific bluefin reach a maximum size of 550 kg and lengths of 300 cm. Maturity is reached at 3 to 5 years of age and individuals live to 15+ years old. Spawning takes place between Japan and the Philippines in April, May and June, spreading to the waters off southern Honshu in July and to the Sea of Japan in August. Pacific bluefin of 270 to 300 kg produce about 10 million eggs but there is no information on the frequency of spawning. Juveniles make extensive migrations north and eastwards across the Pacific Ocean as 1-2 year old fish. Pacific bluefin caught in the southern hemisphere, including those caught in New Zealand waters, are primarily adults.

Natural mortality is assumed to vary from about 0.1 to 0.4 and to be age specific in assessments undertaken by the IATTC. A range of von Bertalanffy growth parameters have been estimated for

Pacific bluefin based on length frequency analysis, tagging and reading of hard parts. These are as follows:

Method	L infinity	k	t ₀
length frequencies	300.0		
scales	320.5	0.1035	- 0.7034
scales	295.4		
tagging	219.0	0.211	

The length weight relationship of Pacific bluefin based on observer data from New Zealand caught fish yields the following:

$$\text{whole weight} = 8.058 e^{0.015 \text{ length}} \quad R^2 = 0.895, n = 49 \text{ (weight is in kg and length is in cm).}$$

Although the sample size of genetically confirmed Pacific bluefin that has been sexed by observers is small (50 fish), the sex ratio in New Zealand waters is not significantly different from 1:1.

3. STOCKS AND AREAS

Pacific bluefin tuna constitutes a single Pacific-wide stock that is primarily distributed in the northern hemisphere.

4. STOCK ASSESSMENT

A stock assessment for Pacific bluefin tuna is under development by the International Scientific Committee for tuna and tuna-like species (ISC) and progress to date was reviewed at the 6th meeting of the ISC, March 2006. The ISC is a group of scientists primarily from countries bordering the North Pacific Ocean who provide advice to the Northern Committee of the WCPFC.

Results from a virtual population analysis (VPA) were compared with results from Stock Synthesis 2 (SS2). Both analyses are preliminary. It was noted that there were considerable uncertainties in the VPA assessment, and while not entirely comparable, the SS2 assessment suggests that the VPA assessment may be optimistic. Revisions to catch data and biological parameters will be considered as part of a revised assessment.

(a) Estimates of fishery parameters and abundance

None are available at present.

(b) Biomass estimates

Estimates of current and reference biomass are not available.

(c) Estimation of Maximum Constant Yield (MCY)

No estimates of MCY are available.

(d) Estimation of Current Annual Yield (CAY)

No estimates of CAY are available.

5. STATUS OF THE STOCKS

No reliable estimates of current biomass are available as stock assessments are under development. Total catches from the stock have not declined although there was a marked decline in abundance of adults over the period 1952-1986. The apparent absence of a spawner-recruit relationship is interpreted to mean that reductions in catches of large Pacific bluefin would not result in improved

recruitment. However, results of cohort analyses and yield-per-recruit analyses indicate that greater catches could be realised if catches of younger fish were reduced.

It is not known if recent catch levels are sustainable. There is no estimate of MSY and it is not possible to determine if recent catch levels will allow the stock to move towards a size that would support the maximum sustainable yield.

6. FOR FURTHER INFORMATION

- Anon. 1993. Annual Report of the Inter-American Tropical Tuna Commission. IATTC, La Jolla, California. 316 p.
- Anon. 1997. Annual Report of the Inter-American Tropical Tuna Commission. IATTC, La Jolla, California. 310 p.
- Anon. 2002. Annual Report of the Inter-American Tropical Tuna Commission. IATTC, La Jolla, California. 148 p.
- Bayliff, W. 1994. A review of the biology and fisheries for northern bluefin tuna, *Thunnus thynnus*, in the Pacific Ocean. *FAO Fisheries Technical Paper* 336/2: 244-294.
- ISC. Report of the sixth meeting of the International Scientific Committee for tuna and tuna like species in the North Pacific. 6th Meeting of the ISC 23-27 March, 2006, La Jolla, California, U.S.A.
- Collette, B. 1999. Mackerels, molecules, and morphology. *Proceedings of the 5th Indo-Pacific Fish Conference*, Noumea, 1997, pp 149–164.
- Smith, P.J. & M. McVeagh. 2006. DNA barcode identification of bluefin tuna samples from 2004-05. NIWA Client Report WLG2006-06, February 2006. Unpublished report held by the Ministry of Fisheries.
- Smith, P.J., L. Griggs & S. Chow. 2001. DNA identification of Pacific bluefin tuna (*Thunnus orientalis*) in the New Zealand fishery. *N.Z. Journal of Marine & Freshwater Research* 35:843–850.